

DESCRIPTION

INDOLE, INDAZOLE, AND BENZAZOLE DERIVATIVE

5 TECHNICAL FIELD

The present invention relates to a novel indole, indazole, and benzazole derivative, or a pharmaceutically acceptable salt thereof, which are useful as a medicament.

10 PRIOR ART

It has been known that β -adrenoceptor of sympathetic nerve has three subtypes, which are classified as $\beta 1$, $\beta 2$ and $\beta 3$ -receptors, and that these receptors distribute in specific tissues in the living body, and each exhibits inherent functions thereof.

15 For instance, $\beta 1$ -adrenoceptor mainly exists in the heart, and the stimulus mediated by $\beta 1$ -adrenoceptor may induce an increase in heart rate as well as the enhancement of cardiac contraction force. On the other hand, $\beta 2$ -adrenoceptors mainly exist in the smooth muscle of blood vessels, the bronchial tube and the uterus, and the stimulus
20 mediated by $\beta 2$ -adrenoceptor may dilate blood vessels and widen the airways, or inhibit the uterine contraction, respectively. In addition, it has been reported that $\beta 3$ -adrenoceptor mainly exists in the adipocytes, the gallbladder and the intestine, and further in the brain, the liver, the stomach, the prostate gland, etc. as well. The stimulus mediated by
25 $\beta 3$ -adrenoceptor may induce the increased lipolysis, the decreased intestinal motility, the increased glucose uptake, and the anti-depression effect, etc.

Further, it has been also reported recently that β 3-adrenoceptor mainly exists in the human urinary bladder, which is relaxed by a β 3-adrenoceptor-stimulating agent.

Hitherto, many β 1-adrenoceptor-stimulating agents and β 2-adrenoceptor-stimulating agents have been developed, and have been
5 used as cardiogenic agents, bronchodilators, or protectants for threatened abortion/threatened premature labor.

On the other hand, β 3-adrenoceptor-stimulating agents have been found to be useful as an agent for treatment or prophylaxis of
10 obesity, hyperglycemia, diseases caused by increased intestinal motility, frequent urination or urinary incontinence, depression, bilestone, or diseases caused by the increased biliary tract motility. At the present, the research and development of excellent β 3-adrenoceptor-stimulating agents have actively been done, and some compounds having β 3-
15 adrenoceptor-stimulating activity have been known (e.g., JP-A-11-255743), but none of these compounds has not been put on the market as a β 3-adrenoceptor-stimulating agent.

Under the above circumstance, it has highly been desired to develop a novel β 3-adrenoceptor-stimulating agent having an excellent
20 β 3-adrenoceptor-stimulating activity.

More preferably, it has been desired to develop a novel β 3-adrenoceptor-stimulating agent with excellent adrenoceptor selectivity wherein side effects such as cardiopalmus or the finger tremor caused by β 1 and/or β 2-adrenoceptor-stimulating activity are reduced by
25 having a more potent β 3-adrenoceptor-stimulating activity than β 1- and/or β 2-adrenoceptor-stimulating activities.


DISCLOSURE OF INVENTION

An object of the present invention is to provide a novel β 3-adrenoceptor-stimulating agent having an excellent β 3-adrenoceptor-stimulating activity, more preferably a novel β 3-adrenoceptor-stimulating agent with excellent adrenoceptor selectivity where side effects such as cardiopalmus or the finger tremor caused by β 1- and/or β 2-adrenoceptor-stimulating activities are reduced by having a more potent β 3-adrenoceptor-stimulating activity than β 1- and/or β 2-adrenoceptor-stimulating activities .

The present inventors have intensively studied in order to solve the above problems, and have found that indole, indazole and benzazole derivatives of the following formula (I) and a pharmaceutically acceptable salt thereof may exhibit an excellent β 3-adrenoceptor-stimulating activity, and they have finally accomplished the present invention.

Among the β 3-adrenoceptor-stimulating agents of the present invention having β 3-adrenoceptor-stimulating activity, the present inventors have further found the compounds of the following [2] as a compound having less β 1- and/or β 2-adrenoceptor-stimulating activities, which are explained below.

The present inventors have found that the compound disclosed in JP-A-11-255743 of the following formula:



does not exhibit $\beta 1$ - and/or $\beta 2$ -adrenoceptor-stimulating activities in the conventional system for evaluating $\beta 1$ - and/or $\beta 2$ -adrenoceptor-


stimulating activities (for example, by the method disclosed in Experiment 1 of the present invention), but exhibits $\beta 1$ - and/or $\beta 2$ -adrenoceptor-stimulating activities in the evaluation system using tissues (for example, by the method disclosed in Experiment 2 of the present invention). This fact is of a problem being pointed out by Sennitt et al. as a divergence in the results obtained with using cells and tissues (cf., J. Pharmacol. Exp. Ther. vol. 285, p. 1084-1095, (1998)).

Then, the present inventors have further studied, and have found that the compound of the following feature [2] wherein R^1 is a group of the formula: $-X-R^{1e}-C(=O)NR^{1a}R^{1b}$ or $-X-R^{1e}-C(=O)OR^{1a}$, X is a group of the formula: $-O-$ or $-S-$, and R^{1e} is a group of the formula: $-CR^{1f}R^{1g}-$ (R^{1f} and R^{1g} are independently a hydrogen atom, an optionally substituted lower alkyl group, an optionally substituted aryl group, or an optionally substituted aralkyl group, or both combine each other together with the carbon atom to which they bond and form an optionally substituted cycloalkane ring, provided that R^{1f} and R^{1g} are not simultaneously a hydrogen atom) exhibits a low $\beta 1$ - and/or $\beta 2$ -adrenoceptor-stimulating activities even in the evaluation system using tissues, as compared to the compounds where R^{1f} and R^{1g} in the same partial structure are simultaneously a hydrogen atom such as the compounds disclosed in the above JP-A-11-255743.


From the above, it is confirmed that the compound of the following feature [2] is an excellent compound as a medicament, because the side effects such as cardiopalmus or the finger tremor are reduced due to the low $\beta 1$ - and/or $\beta 2$ -adrenoceptor-stimulating activities.

The present invention concerns the following features.

[1] A compound of the formula (I):



wherein W is a group of the following formula (VIII) binding to any possible position on the Q;



Q is, together with W, a group of the formula: $-C(W)=C(R^{3A})-N(R^3)-$, $-C(R^{3A})=C(W)-N(R^3)-$, $-C(R^{3A})=C(R^{3B})-N(W)-$, $-C(W)=N-N(R^3)-$, $-C(R^{3A})=N-N(W)-$, $-N=C(W)-N(R^3)-$, $-N=C(R^{3A})-N(W)-$, $-C(W)=N-O-$, or $-C(W)=N-S-$;

R^{3A} and R^{3B} are independently a hydrogen atom or an optionally substituted lower alkyl group;

R^4 , R^5 , R^6 , and R^7 are independently a hydrogen atom or an optionally substituted lower alkyl group;

R^1 is an optionally substituted lower alkyl group, or a group of the formula: $-X-R^{1c}-C(=O)NR^{1a}R^{1b}$, $-X-R^{1c}-C(=O)OR^{1a}$ or $-X-R^{1d}$ (wherein X is a direct bond or a group of the formula: $-O-$, $-S-$, $-N(R^{1c})-$, $-N(R^{1c})C(=O)-$, $-C(=O)N(R^{1c})-$, $-N(R^{1c})SO_2-$, $-SO_2N(R^{1c})-$, or $-C(=O)NHSO_2-$, R^{1c} is a direct bond or an optionally substituted lower alkylene group, R^{1a} , R^{1b} , and R^{1c} are independently a hydrogen atom, an optionally substituted lower alkyl group, an optionally substituted aralkyl group, an optionally substituted cycloalkyl group, or an optionally substituted heterocyclic group, or R^{1a} and R^{1b} may combine each other, and with the adjacent nitrogen atom to which they bond,


form a saturated 3- to 8-membered cyclic amino group optionally having a group of the formula: -O- or -NH- within the ring (said saturated cyclic amino group being substituted or unsubstituted), R^{1d} is a hydrogen atom, an optionally substituted lower alkyl group, an optionally substituted phenyl group, or an optionally substituted cycloalkyl group (a -CH₂- moiety of said cycloalkyl group optionally being replaced by one or more groups of the formula: -O- or -N(R^{1a})-, which are the same or different));

R^2 is a hydrogen atom, a halogen atom, an optionally substituted lower alkyl group, an optionally substituted lower alkenyl group, an optionally substituted amino group, a hydroxy group, a lower alkoxy group, or R^1 and R^2 may combine each other and form a methylenedioxy group and said methylenedioxy group may optionally be substituted by a carboxyl group or a lower alkoxycarbonyl group;

R^3 is a hydrogen atom or an optionally substituted lower alkyl group, or R^1 and R^3 may combine each other and form a divalent group of the formula: -X-R^{1e}-C(=O)- (provided that the bond at the carbonyl side of the above formula binds to the atom to which R^3 of the compound of the formula (I) attaches);

Ar is a group of the following formula (IX), formula (X), or formula (XIII);

A group of Formula (IX):




(in which R^8 is a hydrogen atom, a halogen atom, a trifluoromethyl group, an optionally substituted lower alkyl group, a lower alkoxy group, a lower alkoxycarbonyl group, a carboxyl group, an optionally

substituted benzyloxy group, a hydroxy group, a nitro group, an optionally substituted lower alkylsulfonyl group, an optionally substituted benzenesulfonyl group, an optionally substituted lower alkylthio group, an optionally substituted lower alkylsulfinyl group, a mercapto group, a cyano group, an amino group, an optionally substituted lower alkanoylamino group, an optionally substituted mono- or di-lower alkylamino group, an optionally substituted lower alkylsulfonylamino group, or an optionally substituted benzenesulfonylamino group;

R^9 and R^{10} are independently a hydrogen atom, a halogen atom, an optionally substituted lower alkyl group, a lower alkoxy group, a lower alkoxycarbonyl group, a hydroxy group, an amino group or an optionally substituted mono- or di-lower alkylamino group, or two of R^8 , R^9 , and R^{10} may combine each other and form a methylenedioxy group, and said methylenedioxy group may optionally be substituted by a carboxyl group or a lower alkoxycarbonyl group, or two of R^8 , R^9 , and R^{10} may combine each other and form a group of the formula: $-NR^{8a}C(=O)CR^{8b}=CR^{8c}-$ (R^{8a} , R^{8b} , and R^{8c} being the same or different and each a hydrogen atom or an optionally substituted lower alkyl group);

provided that when R^1 is a group of the formula: $-O-CH_2-C(=O)OR^{1a}$ and all of R^4 , R^5 , R^9 , and R^{10} are a hydrogen atom, then R^8 is not a halogen atom or a trifluoromethyl group substituting on the 3-position);

A group of Formula (X):



(in which Z is an oxygen atom or a sulfur atom;


R¹¹ is a hydrogen atom, a lower alkyl group, or a group of the formula:
 -SO₂R¹⁴ or the formula: -NR¹⁵R¹⁶ (R¹⁴ is an optionally substituted lower
 5 alkyl group, an optionally substituted phenyl group, an optionally
 substituted aralkyl group, R¹⁵ and R¹⁶ are independently a hydrogen
 atom, an optionally substituted lower alkyl group, or an optionally
 substituted benzyl group);

R¹² is an oxygen atom, a sulfur atom or H₂;

10 R¹³ is an oxygen atom or H₂;

nn and mm are each 0 or 1); or

A group of Formula (XIII):




(in which R¹⁷ is a hydrogen atom, a halogen atom, or a cyano group),
 15 or a pharmaceutically acceptable salt thereof.

[2] The compound according to the above [1], wherein R¹ is a group
 of the formula: -X-R^{1e}-C(=O)NR^{1a}R^{1b} or the formula: -X-R^{1e}-C(=O)OR^{1a},
 or R¹ and R³ may combine each other and form a divalent group of the
 formula: -X-R^{1e}-C(=O)- (where X is a group of the formula: -O- or -S-
 20 and R^{1e} is a group of the formula: -CR^{1f}R^{1g}- (R^{1f} and R^{1g} are
 independently a hydrogen atom, an optionally substituted lower alkyl
 group, an optionally substituted cycloalkyl group, an optionally

substituted aryl group, or an optionally substituted aralkyl group, or both may combine each other, and with the carbon atom to which they bond, form an optionally substituted cycloalkane ring, provided that both R^{1f} and R^{1g} are not simultaneously a hydrogen atom)),

or a pharmaceutically acceptable salt thereof.

[3] The compound according to the above [1] or [2], which is a compound of the formula (I'):



wherein R¹, R², R³, R^{3A}, and W are as defined in the above [1],

or a pharmaceutically acceptable salt thereof.


[4] The compound according to the above [3], wherein R¹ binds to the 5-, 6- or 7-position of the indole ring of the compound of the formula (I'), and R² is a hydrogen atom,


or a pharmaceutically acceptable salt thereof.

[5] The compound according to the above [3], wherein R² is a group other than a hydrogen atom, and one of R¹ and R² binds to the 6-position of the indole ring of the compound of the formula (I'), and the other binds to the 7-position thereof,

or a pharmaceutically acceptable salt thereof.

[6] The compound according to any one of the above [1] to [5], wherein Ar is a group selected from the following substituents:





(wherein n is 0, 1, or 2),

or a pharmaceutically acceptable salt thereof.

[7] The compound according to any one of the above [1] to [6],
 wherein R^1 is a group of the formula: $-X-R^{1e}-C(=O)NR^{1a}R^{1b}$ or $-X-R^{1e}-C(=O)OR^{1a}$;

X is a direct bond or a group of the formula: $-O-$;

R^{1a} and R^{1b} , when it exists, are independently selected from

- (i) a hydrogen atom,
- (ii) an unsubstituted lower alkyl group,
- (iii) a lower alkyl group being substituted by one or more

substituents, which are the same or different, and said substituent(s) are selected from a carboxyl group, a lower alkoxycarbonyl group, an amino group, a hydroxy group, an alkoxy group, a mercapto group, an alkylthio group, a carbamoyl group, an indolyl group, a guanidino group, an imidazolyl group, and a phenyl group optionally being substituted by a hydroxy group, or

- (iv) a saturated 3- to 8-membered cyclic amino group which is formed by combining R^{1a} and R^{1b} together with the adjacent nitrogen atom to which they bond, and optionally has a group of the formula: $-O-$ or $-NH-$ within the ring (said saturated cyclic amino group being

unsubstituted, or optionally being substituted by a carboxyl group or a lower alkoxy carbonyl group), or
a pharmaceutically acceptable salt thereof.

[8] The compound according to the above [7], wherein R^1 is a group of the formula: $-X-R^{1e}-C(=O)NR^{1a}R^{1b}$ (where the moiety of the formula: $NR^{1a}R^{1b}$ is an amino acid or amino acid ester residue binding to the carbonyl group of the above formula at the N-terminus thereof, and R^{1a} binds to the nitrogen atom at the N-terminus thereof when R^{1a} and R^{1b} do not form a cyclic group), and


X and R^{1e} are a direct bond,
or a pharmaceutically acceptable salt thereof.

[9] The compound according to any one of the above [1] to [3], wherein R^1 is a group of the formula: $-C(=O)NR^{1a}R^{1b}$, R^{1a} and R^{1b} are independently a hydrogen atom or an optionally substituted lower alkyl group, or R^{1a} and R^{1b} may combine each other, and with the adjacent nitrogen atom to which they bond, form a saturated 3- to 8-membered cyclic amino group optionally having a group of the formula: $-O-$ or $-NH-$ within the ring (said saturated cyclic amino group being unsubstituted, or optionally being substituted by a carboxyl group or a lower alkoxy carbonyl group);

R^2 is a hydrogen atom, a halogen atom, an optionally substituted lower alkyl group, a hydroxy group, or a lower alkoxy group;

both of R^4 and R^5 are a hydrogen atom;

Ar is a group of the formula (XVI):



(in which R^{18} is a halogen atom or a trifluoromethyl group),

or a pharmaceutically acceptable salt thereof.

[10] The compound according to any one of the above [2] to [6], wherein R^1 is a group of the formula: $-X-CR^{1a}R^{1b}-C(=O)OR^{1a}$, or a pharmaceutically acceptable salt thereof.

5 [11] The compound according to any one of the above [2] to [6], wherein X is a group of the formula: $-O-$, or a pharmaceutically acceptable salt thereof.

[12] The compound according to any one of the above [1], [2], [6] to [11], wherein Q is, together with W, a group of the formula:
10 $-C(W)=C(R^{3A})-N(R^3)-$ or $-C(R^{3A})=C(W)-N(R^3)-$, or a pharmaceutically acceptable salt thereof.

[13] A pharmaceutical composition comprising as an active ingredient the compound as set forth in any one of the above [1] to [12], or a pharmaceutically acceptable salt thereof.

15 [14] An agent for treatment of obesity, hyperglycemia, frequent urination, urinary incontinence, depression, or bilestone, which comprises as an active ingredient the compound as set forth in any one of the above [1] to [12], or a pharmaceutically acceptable salt thereof.

[15] A method for treatment of obesity, hyperglycemia, frequent
20 urination, urinary incontinence, depression, or bilestone, which comprises administering to a patient in need an effective amount of the compound as set forth in any one of the above [1] to [12], or a pharmaceutically acceptable salt thereof.

[16] A use of the compound as set forth in any one of the above [1]
25 to [12], or a pharmaceutically acceptable salt thereof, in preparation of an agent for treatment of obesity, hyperglycemia, frequent urination, urinary incontinence, depression, or bilestone.

BEST MODE FOR CARRYING OUT THE INVENTION

Each group in the present invention is explained below.

Unless defined otherwise, the definition for each group should be applied to cases wherein said group is a part of another substituent.

The "substituted benzene", the "substituted phenyl group" and the "substituted aryl group" may have one or more substituents, and such substituents include, for example, a halogen atom, a C₁-C₈ haloalkyl group, a C₁-C₈ alkyl group, a C₂-C₈ alkenyl group, a C₁-C₈ alkoxy group, a hydroxy group, a nitro group, a cyano group, a mercapto group, a group of the formula: -S(O)_p(C₁-C₈ alkyl), a carboxyl group, a C₁-C₈ alkoxycarbonyl group, an aryloxycarbonyl group, an aralkyloxycarbonyl group, an optionally substituted amino group, an optionally substituted amido group, an optionally substituted urea group, an optionally substituted sulfonamido group, a group of the formula: -C(O)NHSO₂(C₁-C₈ alkyl), etc. (in the above formula, p is 0, 1 or 2, hereinafter the same).

The "aryl group" includes, for example, an aryl group having not more than 10 carbon atoms such as phenyl, 1- or 2-naphthyl, etc.

The aryl moiety of the "aralkyl group" includes, for example, an aryl group having not more than 10 carbon atoms such as phenyl, 1- or 2-naphthyl, etc., and the alkyl moiety thereof includes, for example, an alkyl group having not more than 5 carbon atoms such as methyl, ethyl, propyl, butyl, etc. Representative aralkyl group is, for example, benzyl group, 1- or 2-phenetyl group, etc.

The "substituted aralkyl group" may have one or more substituents on the aryl moiety and/or the alkyl moiety, and such

substituents include, for example, a halogen atom, a C₁-C₈ haloalkyl group, a C₁-C₈ alkyl group, a C₂-C₈ alkenyl group, a C₁-C₈ alkoxy group, a hydroxy group, a nitro group, a mercapto group, a group of the formula: -S(O)_p(C₁-C₈ alkyl), a carboxyl group, a C₁-C₈ alkoxycarbonyl group, an aryloxycarbonyl group, an aralkyloxycarbonyl group, an optionally substituted amino group, an optionally substituted amido group, an optionally substituted urea group, an optionally substituted sulfonamido group, a group of the formula: -C(O)NHSO₂(C₁-C₈ alkyl), etc.

The "alkyl group" includes "lower alkyl group". The "lower alkyl group" includes, unless defined otherwise, a straight chain or branched chain saturated C₁-C₈ hydrocarbon group, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, and sec-butyl, and further includes higher homologs and isomers such as n-pentyl, n-hexyl, 2-methylpentyl, etc.

The "substituted alkyl group", the "substituted alkenyl group", and the "substituted alkylene group" may have one or more substituents, and such substituents include, for example, a halogen atom, a C₁-C₈ alkoxy group, a C₁-C₈ alkoxycarbonyloxy group, a C₃-C₈ cycloalkyloxycarbonyloxy group, a hydroxy group, a mercapto group, a group of the formula: -S(O)_p(C₁-C₈ alkyl), a C₃-C₈ cycloalkyl group, an optionally substituted amino group, a carboxyl group, a C₁-C₈ alkoxy-carbonyl group, an aryloxycarbonyl group, an aralkyloxycarbonyl group, an optionally substituted heterocyclic group, an optionally substituted aryl group, an oxo group, etc.

The "alkenyl group" includes "lower alkenyl group". The "lower alkenyl group" includes a straight chain or branched chain alkenyl group having not more than 8 carbon atoms, for example, vinyl, allyl,

propenyl, 2-propenyl, butenyl, pentenyl, hexenyl, etc.

The "alkylene group" includes "lower alkylene group". The "lower alkylene group" includes a straight chain or branched chain alkylene group having not more than 8 carbon atoms, for example, methylene, ethylene, trimethylene, tetramethylene, pentamethylene, hexamethylene, etc.

The "optionally substituted alkoxy group" includes an optionally substituted alkyl group having an oxygen atom at the binding position thereof.

The "optionally substituted alkylthio group" includes an optionally substituted alkyl group having a sulfur atom at the binding position thereof.

The "optionally substituted alkylsulfinyl group" includes an optionally substituted alkyl group having a group of the formula: -SO- at the binding position thereof.

The "optionally substituted alkylsulfonyl group" includes an optionally substituted alkyl group having a group of the formula: -SO₂- at the binding position thereof.

The "optionally substituted alkoxycarbonyl group" includes an optionally substituted alkyl group having a group of the formula: -OC(=O)- at the binding position thereof via the oxygen atom side of the group of the formula: -OC(=O)-.

The "optionally substituted alkanoyl group" includes an optionally substituted alkyl group having a group of the formula: -C(=O)- at the binding position thereof.

The "optionally substituted alkanoylamino group" includes an optionally substituted alkyl group having a group of the formula:

-NHC(=O)- at the binding position thereof via the carbon atom side of the group of the formula: -NHC(=O)-, wherein the nitrogen atom is optionally substituted by a C₁-C₈ haloalkyl group, a C₁-C₈ alkyl group, a C₂-C₈ alkenyl group, a C₃-C₈ cycloalkyl group, a phenyl group, an aralkyl group, a heterocyclic group, etc.

The "optionally substituted alkylaminocarbonyl group" includes an optionally substituted alkyl group having a group of the formula: -C(=O)NH- at the binding position thereof via the nitrogen atom side of the group of the formula: -C(=O)NH-, wherein the nitrogen atom is optionally substituted by a C₁-C₈ haloalkyl group, a C₁-C₈ alkyl group, a C₂-C₈ alkenyl group, a C₃-C₈ cycloalkyl group, a phenyl group, an aralkyl group, a heterocyclic group, etc.

The "halogen atom" includes chlorine atom, bromine atom, fluorine atom, iodine atom, etc.

The "optionally substituted mono- or di-lower alkylamino group" includes an amino group where one or both of hydrogen atoms of the amino group are independently substituted by an optionally substituted alkyl group.

The "optionally substituted amido group" is a group of the formula: -NR¹⁹COR²⁰, where R¹⁹ is a hydrogen atom, a C₁-C₈ alkyl, etc., and R²⁰ is a C₁-C₈ haloalkyl group, a C₁-C₈ alkyl group, a C₂-C₈ alkenyl group, a C₃-C₈ cycloalkyl group, a phenyl group, an aralkyl group, a heterocyclic group, etc.

The "optionally substituted sulfonamido group" is a group of the formula: -NR²¹SO₂R²², where R²¹ is a hydrogen atom, C₁-C₈ alkyl, etc., and R²² is a C₁-C₈ haloalkyl group, a C₁-C₈ alkyl group, a C₂-C₈ alkenyl group, a C₃-C₈ cycloalkyl group, a phenyl group, an aralkyl

group, a heterocyclic group, etc.

The "optionally substituted alkylsulfonylamino group" is a group of the formula: $-NR^{23}SO_2R^{24}$, where R^{23} is a C_1 - C_8 haloalkyl group, a C_1 - C_8 alkyl group, a C_2 - C_8 alkenyl group, a C_3 - C_8 cycloalkyl group, a phenyl group, an aralkyl group, a heterocyclic group, etc., and R^{24} is a C_1 - C_8 alkyl group, etc.

The "substituted amino group" is an amino group where one or both of hydrogen atoms are independently substituted by a C_1 - C_8 alkyl group, a C_2 - C_8 alkenyl group, a C_1 - C_8 alkoxy group, a hydroxy group, etc.

The cycloalkyl group includes, for example, a 3- to 8-membered cycloalkyl group such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, etc.

The cycloalkane ring, which is formed by combining R^{1f} and R^{1g} of the group of the formula: $-CR^{1f}R^{1g}-$ together with the nitrogen atom to which they bond, includes, for example, a 3- to 8-membered cycloalkane ring wherein two hydrogen atoms on the same carbon atom are replaced by two bonds, such as cyclopropane, cyclobutane, cycloheptane, cyclohexane, cycloheptane, etc.

The "substituted cycloalkyl group" and the "substituted cycloalkane ring" may have one or more substituents, and such substituents are, in addition to an alkyl group and an aralkyl group, the same substituents for the "substituted alkyl group" as mentioned above.

The "heterocycle" includes a 5-membered or 6-membered aromatic heterocycle or a saturated or unsaturated aliphatic heterocycle, for example, a 5-membered or 6-membered heterocyclic group consisting of 1 to 4 heteroatoms selected from a nitrogen atom, an

oxygen atom and a sulfur atom and carbon atoms, such as pyridine ring, imidazole ring, pyrazine ring, pyrimidine ring, pyridazine ring, thiazole ring, isothiazole ring, isothiazoline ring, oxazole ring, isoxazole ring, isoxazoline ring, furan ring, thiophene ring, pyrrole ring, pyrroline ring, pyrrolidine ring, pyrazoline ring, imidazoline ring, tetrahydro-
 5 pyrane ring, tetrahydrofuran ring, tetrahydrothiophene ring, pyrrolidine ring, piperidine ring, etc.

The substituents of the "heterocycle" are 1 to 2 groups independently selected from a halogen atom, a C₁-C₈ alkyl group, a C₂-C₈ alkenyl group, a C₁-C₈ alkoxy group, a hydroxy group, a mercapto group, a group of the formula: -S(O)_p(C₁-C₈ alkyl), a carboxyl group, a C₁-C₈ alkoxycarbonyl group, an aryloxy carbonyl group, an aralkyloxy carbonyl group, an amino group, an alkylamino group, an aryl group, an aralkyl group, and an oxo group, if the substitution is available.

15 The "heterocyclic group" includes a group where the hydrogen atom of the above "heterocycle" is replaced by a bond, and the substituents of the "heterocyclic group" are the same substituents as those for the above "heterocycle".

20 The 3- to 8-membered saturated cyclic amino group optionally having a group of the formula: -O- or -NH- within the ring includes, for example, 1-pyrrolidinyl, piperidino, 1-piperazinyl, morpholino, etc.

25 The substituents of the 3- to 8-membered saturated cyclic amino group optionally having a group of the formula: -O- or -NH- within the ring includes, for example, the same substituents as those for the "heterocycle" as mentioned above.


The "amino acid residue" includes a group where the hydrogen atom at the N-terminus of an amino acid is replaced by a bond. The

“amino acid ester residue” includes a group where the hydrogen atom of the carboxyl group (at least one carboxyl group when two or more exist) of an amino acid residue is replaced by an alkyl group, an aryl group, or an aralkyl group.

5 The indolyl group includes 1-indolyl group and 2-indolyl group.


 The imidazolyl group includes 2-imidazolyl group, 4-imidazolyl group, and 5-imidazolyl group.

 When R^1 of the compound of the formula (I) is a group of the formula: $-X-R^{1e}-C(=O)OR^{1a}$, more concretely, such a group is a group of
10 the following formula:



(in which R^b is a hydrogen atom, an optionally substituted lower alkyl group, an optionally substituted lower alkoxy carbonyl group or a carboxyl group, R^{bb} is an optionally substituted lower alkoxy carbonyl
15 group or a carboxyl group, and m is an integer of 0-3)

 When R^1 of the compound of the formula (I) is a group of the formula: $-X-R^{1d}$, more concretely, such a group is a group of $-O(CH_2)_p-R^c$
(in which R^c is an optionally substituted lower alkanoyl group, a hydroxy group, a cyano group, an optionally substituted phenyl group,
20 an optionally substituted heterocyclic group, an optionally substituted mono- or di-lower alkylaminocarbonyl group, or a group of the following formula (Va):




(wherein R^A is a hydrogen atom or a lower alkyl group), and p is an
25 integer of 1 to 4).

The compounds of the present invention are prepared, for example, by the following Processes.

Process (a):

Among the present compounds, the compound wherein the substituents R^4 and R^5 in the formula (VIII) are a hydrogen atom, i.e., the compound of the following formula (I-a):




(in which R^1 , R^2 , R^6 , R^7 and Ar are as defined above, and Q is the same as defined for the compound of the formula (I) except that W is replaced by a group of the formula: $\text{ArCH(OH)CH}_2\text{NHC(R}^6\text{(R}^7\text{))CH}_2\text{-}$)

may be prepared by reacting a compound of the following formula (XVII):



(in which Ar is as defined above),

with a compound of the following formula (XVIII):



(in which R^1 , R^2 , R^6 , and R^7 are as defined above, and Q is the same as defined for the compound of the formula (I) except that W is replaced by a group of the formula: $\text{NH}_2\text{C(R}^6\text{(R}^7\text{))CH}_2\text{-}$), or a salt thereof.

This reaction is carried out in a suitable solvent or without a solvent. The solvent to be used should be selected according to the kinds of the starting compounds, etc., and includes, for example,

alcohols such as methanol, ethanol, isopropyl alcohol, ketones such as acetone, methyl ethyl ketone, halogenated hydrocarbons such as methylene chloride, chloroform, ethers such as diethyl ether, tetrahydrofuran, dioxane, aromatic hydrocarbons such as benzene, toluene, ethyl acetate, N,N-dimethylformamide, dimethylsulfoxide, etc., and these solvents may be used alone or in a mixture of two or more solvents.

Besides, when the compound of the formula (XVIII) is in the form of an acid addition salt, for example, a salt with an inorganic acid such as hydrochloride, hydrobromide, etc., or a salt with an organic acid such as oxalate, maleate, fumarate, etc., this reaction is carried out in the presence of a base.

Example of base is an alkali metal hydrogen carbonate such as sodium hydrogen carbonate, potassium hydrogen carbonate, an alkali metal carbonate such as sodium carbonate, potassium carbonate, or an organic base such as triethylamine, tributylamine, diisopropyl ethylamine, N-methylmorpholine, etc.

The reaction temperature may vary according the kinds of the starting compounds, etc., but this reaction is usually carried out at a temperature of from room temperature to about 150°C, preferably at a temperature of from about 25°C to about 100°C.

In this Process, when the starting compound of the formula (XVII) and the formula (XVIII) have an asymmetric carbon atom, the configuration based on said asymmetric carbon atom is maintained into the reaction product of the formula (I-a). That is, for example, the present compound in the (R,R)-configuration is obtained from the compound of the formula (XVII) in the R-configuration and the


compound of the formula (XVIII) in the R-configuration.

The optically active compound of the above formula (XVII) may be prepared according to the method disclosed in the literatures (the method of Bloom, J. D. et al. (J. Med. Chem., 35, 3081-3084 (1992)), or
 5 the method of Eliel, E. L. and Delmonte, D. W. (J. Org. Chem., 21, 596-597 (1956))).

The optically active compound of the above formula (XVIII) may be prepared according to the method disclosed in the literatures (the method of Repke, D. B. and Ferguson, W. J. (J. Heterocycl. Chem., 13,
 10 775-778 (1976))).

The starting compound of the above formula (XVIII) may be prepared, for example, by the method disclosed in the literature (J. Org. Chem., 25, 1548-1558 (1960)).

In addition, among the starting compounds of the above
 15 formula (XVIII), the compound having a partial structure of the following formula W':



(wherein R⁶ and R⁷ are as defined above)

being attached to the 3-position of the indole ring or the indazole ring
 20 may be prepared by the method disclosed in the literature (cf., J. Org. Chem., 51, 4294-4295 (1986), Tetrahedron Lett., 41, 4363-4366 (2000), etc.).


Further, among the starting compounds of the above formula (XVIII), the compound having the partial structure W' being attached to
 25 the 1-position of the indole ring, or to the 1- or 3-position of the benzimidazole ring, or to the 1-position of the indazole ring may be

prepared by the method disclosed in the literature (cf., J. Med. Chem., 40, 2003-2010 (1997), Aust. J. Chem., 46, 1177-1191 (1993), etc.).

Moreover, among the starting compounds of the above formula (XVIII), the compound having the partial structure W' being attached to the 2-position of the indole ring may be prepared by the method disclosed in the literature (cf., J. Heterocyclic. Chem., 36, 921-926 (1999), etc.).

Process (b):

As an alternative process to Process (a), the compound of the present invention of the formula (I) may be prepared by reacting a compound of the formula (XIX):



(wherein R⁴, R⁵, and Ar are as defined above, A is a protecting group for hydroxy group, B is a bromine atom or a iodine atom, and *1 means an asymmetric carbon atom)


with the compound of the above formula (XVIII), followed by removing the protecting group A.

The protecting group for hydroxy group is not necessarily specified, and may be any conventional protecting groups. For example, the protecting groups being usually removed selectively and easily are a benzyl group or a tert-butyldimethylsilyl group, etc. These protecting groups for hydroxy group are introduced by a conventional method. For example, a benzyl group may be introduced by reacting benzyl bromide in an amount of equimolar or 2 moles with sodium iodide in an amount of 1.1 mole in a solvent such as dimethylform-

amide in the presence of potassium carbonate at room temperature. A triethylsilyl group may be introduced by reacting with a silylating agent such as triethylsilyl chloride in an amount of 1.2 to 2 moles in a solvent such as pyridine at a temperature of from 0°C to 30°C for 1 to 3 hours.

5 The coupling reaction of the compound of the formula (XIX) with the compound of the formula (XVIII) is carried out by heating the compound of the formula (XVIII) in an amount of 1 to 1.5 moles to 1 mole of the compound of the formula (XIX) in the presence of an amine such as triethylamine or diisopropyl ethylamine as a trapping agent for
10 proton in a polar solvent such as dimethylformamide, dimethylacetamide or dimethylsulfoxide at a temperature of from room temperature to 90°C, preferably at 60°C for 5 to 10 hours. The deprotection is usually carried out by a hydrogenation with a catalyst such as palladium or nickel in a solvent such as methanol when the
15 protecting group A is a benzyl group. When the protecting group A is a benzyl group or methyl group, then the deprotection is usually carried out by treating with a Lewis acid such as boron tribromide in a solvent such as methylene chloride, etc. When the protecting group A is a triethylsilyl group, the deprotection is usually carried out by treating
20 with acetic acid and tetrabutylammonium fluoride in an amount of 3 to 5 times by mole in tetrahydrofuran at room temperature for 30 minutes to 5 hours.

The compound of the formula (XIX) may be prepared, for example, by reducing a compound of the formula (XX):




(wherein R⁴, R⁵, Ar, and B are defined above)

by the following methods, followed by protecting the hydroxy group.

Namely, the reduction of the compound of the formula (XX) is carried out by reducing with a reducing agent such as borane or sodium borohydride when the configuration (*1) of the hydroxy group of the desired compound of the formula (XIX) is racemic. The reaction is usually carried out in a solvent such as ethers (e.g., diethyl ether, tetrahydrofuran, etc.) or alcohols (e.g., methanol, ethanol, etc.) at a temperature of from 0°C to a boiling point of the solvent.

Further, with respect to the configuration *1 in the formula (XIX), in order to obtain an optical isomer R or S, the reaction is carried out using a chiral auxiliary of the following formula:



for example, by reducing the compound of the formula (XX) with boron in the presence of one of the above two chiral auxiliaries. The reduction is preferably carried out in a solvent such as tetrahydrofuran. The preparation and the reaction of these chiral auxiliaries are carried out according to the method disclosed in the literature (E. J. Corey et al., J. Org. Chem., vol. 56, p. 442, 1991).


When the bromine atom (bromine compound) should be converted into iodine atom after the reduction of the compound of the formula (XX), the compound obtained by the above reduction is additionally heated under reflux with a iodinating agent such sodium iodide in an amount of 3 to 10 moles to one mole of the bromine compound in a solvent such as acetone for 1 to 3 hours.

Then, by the method for protecting a hydroxy group as mentioned above, the hydroxy group is protected by a triethylsilyl group, etc. to give the compound of the formula (XIX).


The compound of the formula (XX) may be prepared by the method disclosed in the literature (A. A. Larsen et al., J. Med. Chem., 1967, vol. 10, p. 462 or C. Kaiser et al., J. Med. Chem., 1974, vol. 17, p. 49).

Process (c):


As an alternative method to Process (a) and Process (b), among the compounds of the present invention, the compound of the formula (I) wherein the substituent R^7 is a hydrogen atom, that is, the compound of the following formula (I-c):



(wherein R^1 , R^2 , R^4 , R^5 , R^6 , and Ar are as defined above, Q is the same as defined for the compound of the formula (I) except that W is replaced by a group of the formula: $\text{ArCH(OH)CR}^4(\text{R}^5)\text{NHCH(R}^6\text{)CH}_2\text{-}$) may be prepared by reacting a compound of the formula (XXII):



(wherein R^4 , R^5 , and Ar are as defined above) with a compound of the following formula (XXIII):



(wherein R^1 , R^2 , R^3 , and R^6 are as defined above, Q is the same as defined for the compound of the formula (I) except that W is replaced by a group of the formula: $R^6C(=O)CH_2-$) under reduction conditions.

5 In this reaction, the method of "reaction under reduction conditions" means that the compound of the formula (XXII) and the compound of the formula (XXIII) are reacted in the presence of a reducing agent which may reduce only an imine moiety formed in course of the reaction without affecting the carbonyl group, or in the
10 presence of a catalyst.

 The reducing agent used in this reaction includes, for example, sodium cyanoborohydride, and the catalyst includes, for example, palladium, platinum oxide, etc.

 This reaction is carried out in the presence of a reducing agent
15 or a catalyst in a suitable solvent. The solvent is preferably alcohols such as methanol, ethanol, etc. The reaction temperature is usually in the range of about 20 to about 80°C when a reducing agent is used. When a catalyst is used, the reaction temperature is usually in the range of about 10°C to about 25°C.


20 The compound of the formula (XXII) used as a starting compound is prepared by the optical resolution of a commercially available enantiomer mixture by a conventional method, or by the method disclosed in the literature (cf., J. Med. Chem., vol. 20, no. 7, p. 978-981 (1977)).

25 Further, the starting compound of the formula (XXIII) may be prepared by the method disclosed in the literatures (J. Org. Chem., vol. 55, no. 4, p. 1390-1394 (1990), Chem. Pharm. Bull., vol. 45, no. 5, p.

932-935 (1997), Heterocycles, vol. 32, no. 10, p. 1923-1931 (1991)).

Process (d):

As an alternative to Process (a), Process (b), and Process (c), the compound of the formula (I-a) may be prepared by reducing a compound of the following formula (XXIV):




(wherein R^1 , R^2 , R^6 , R^7 , and Ar are as defined above; Q is the same as defined for the compound of the formula (I) except that W is replaced by a group of the formula: $\text{ArCH}(\text{OH})\text{C}(=\text{O})\text{NHCR}^6(\text{R}^7)\text{CH}_2-$).

This process is carried out in the presence of a reducing agent in a solvent. The reducing agent to be used in this Process includes, for example, diborane, lithium aluminum hydride and an alkoxy complex thereof or a transition metal salt thereof, or a sodium borohydride with aluminum chloride, boron trifluoride, phosphorous oxychloride or a carboxylic acid (e.g., acetic acid, trifluoroacetic acid), etc. The solvent includes, for example, ethers (e.g., diethyl ether, tetrahydrofuran, dimethoxyethane, dioxane, diglyme, etc.). The reaction temperature may vary according to the kinds of the reducing agent, etc., and it is usually in the range of from about 0°C to about 160°C .

In this reaction, the configuration based on the asymmetric carbon atom of the starting compound of the formula (XXIV) is maintained into the product of the reaction.

The compound of the formula (XXIV), which is the starting compound of the above reaction, may be prepared, for example, by

reacting a compound of the formula (XXV):



(wherein Ar is the same as defined above)

with the compound of the above formula (XVIII) or a salt thereof.

5 The reaction of the compound of the formula (XXV) with the compound of the formula (XVIII) is carried out in the presence of a condensing agent such as N,N'-dicyclohexylcarbodiimide, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, N,N'-carbonyl-diimidazole, N,N'-carbonyldisuccinimide, 1-ethoxycarbonyl-2-ethoxy-
10 1,2-dihydroquinoline, diphenylphosphorylazide, propanephosphoric anhydride, etc. When N,N'-dicyclohexylcarbodiimide or 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride is used as a condensing agent, N-hydroxysuccinimide, 1-hydroxybenzotriazole, etc. may be added to the reaction system.

15 This reaction is carried out in a suitable solvent. The solvent includes, for example, the same solvents exemplified for Process (a). Further, the compound of the formula (XVIII) may be used in the form of an acid addition salt, as mentioned in Process (a). In these cases, the reaction is carried out in the presence of an organic base such as
20 triethylamine, tributylamine, diisopropylethylamine, N-methyl-morpholine. The reaction temperature is usually in the range of about 20°C to about 50°C.


 The configuration based on the asymmetric carbon atom of the compound of the formula (XXV) and the compound of the formula
25 (XVIII) is maintained into the product of this reaction (i.e., the compound of the formula (XXIV)).

The optically active compound of the formula (XXV) may be prepared according to the method disclosed in the literature (cf., the method of Collet, A. and Jacques, J. (Bull. Soc. Chim. France, 3330-3334 (1973))).

5 The optically active compound of the above formula (XVIII) may be prepared, for example, according to the method disclosed in JP-B-63-22559.

Process (e):

10 As an alternative to Process (a), Process (b), Process (c) and Process (d), the compound of the formula (I-a) may be prepared by reduction of a compound of the formula (XXVI):




15 (wherein R^1 , R^2 , R^6 , R^7 , and Ar are as defined above, Q is the same as defined in the compound of the formula (I) except that W is replaced by a group of the formula: $\text{ArCH(OH)CR}^4(\text{R}^5)\text{NHC(=O)CH}_2-$).

20 This process is carried out in a solvent in the presence of a reducing agent. The reducing agent to be used in this process includes, for example, diborane, lithium aluminum hydride and an alkoxy complex thereof or a transition metal salt thereof, or a sodium borohydride with aluminum chloride, boron trifluoride, phosphorous oxychloride or a carboxylic acid (e.g., acetic acid, trifluoroacetic acid), etc. The solvent includes, for example, ethers (e.g., diethyl ether, tetrahydrofuran, dimethoxyethane, dioxane, diglyme, etc.). The reaction temperature may vary according to the kinds of the reducing agent, etc., and it is usually in the range of about 0°C to about 160°C .

25

In this process, the configuration based on the asymmetric carbon atom of the starting compound of the formula (XXIV) is maintained into the product of the reaction.

The compound of the formula (XXVI), which is a starting compound of the above reaction, may be prepared by reacting a compound of the formula (XXVII):



(wherein R¹ and R² are as defined above, and Q is the same as defined for the compound of the formula (I) except that W is replaced by a group of the formula: HOC(=O)CH₂-) with the compound of the above formula (XXII) or a salt thereof.


The reaction of the compound of the formula (XXVII) with the compound of the formula (XXII) is carried out in the presence of a condensing agent such as N,N'-dicyclohexylcarbodiimide, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, N,N'-carbonyldiimidazole, N,N'-carbonyldisuccinimide, 1-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline, diphenylphosphorylazide, propanephosphoric anhydride, etc. When N,N'-dicyclohexylcarbodiimide or 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride is used as a condensing agent, N-hydroxysuccinimide, 1-hydroxybenzotriazole, etc. may be added to the reaction system.

This reaction is carried out in a suitable solvent. The solvent includes, for example, the same solvents exemplified for Process (a). Further, the compound of the formula (XVIII) may be used in the form of an acid addition salt, as mentioned in Process (a). In these cases,

the reaction is carried out in the presence of an organic base such as triethylamine, tributylamine, diisopropylethylamine, N-methyl-morpholine. The reaction temperature is usually in the range of about 20°C to about 50°C.

5 The configuration based on the asymmetric carbon atom in the compound of the formula (XXVII) and the compound of the formula (XXII) is maintained into the product of this reaction (i.e., the compound of the formula (XXVI)).

10 Among the compounds of the formula (XXV), the compound of the formula (XXV) wherein the partial structure W" of the following

formula:  attaches to the 3-position of the indole ring or the benzisoxazole ring may be prepared by the method disclosed in the literature (cf., J. Biological Chemistry, 23, 12685-12689 (1985), J. Heterocycl. Chem., 6, 279-283 (1969)).


15 In addition, among the starting compounds of the above formula (XXV), the compound of the formula (XXV) wherein the partial structure W" attaches to the 2-position of the benzimidazole ring may be prepared by the method disclosed in the literature (cf., Bull. Soc. Chim. Belg, 100, 277-286 (1991), etc.).

20 Further, among the starting compounds of the above formula (XXV), the compound of the formula (XXV) wherein the partial structure W" attaches to the 3-position of the benzisothiazole ring may be prepared by the method disclosed in the literature (cf., J. Chem. Soc. Perkin Trans 1, 3006 (1972), etc.).

25 Process (f):

 Among the compounds of the present invention, the compound

of the formula (I) wherein R^1 is a group of the formula: $-X-R^{1e}-C(=O)-NR^{1a}R^{1b}$ or $-X-R^{1e}-C(=O)OR^{1a}$ (X is a group of the formula: $-O-$ or $-S-$) may be prepared by reacting a compound of the formula (XXVII):



5 (wherein A^1 is a hydrogen atom or a protecting group for hydroxy group; A^2 is a hydrogen atom or a protecting group for amino group; X, R^2 , R^4 , R^5 , R^6 , R^7 and Ar are as defined above; Q is the same as defined for the compound of the formula (I) except that W is replaced by a group of the formula: $ArCH(OA^1)CR^4(R^5)NA^2CR^6(R^7)CH_2-$) with a compound of the
10 formula: $YR^{1e}-C(=O)NR^{1a}R^{1b}$ or $YR^{1e}-C(=O)OR^{1a}$ (wherein Y is an alcoholic reactive residue, and R^{1e} , R^{1a} , R^{1b} are as defined above), followed by removing the protecting group, if necessary.

The "alcoholic reactive residue" includes, for example, a halogen atom, a lower alkylsulfonyloxy group such as methanesulfonyloxy, ethanesulfonyloxy, etc., and an arylsulfonyloxy group such as
15 benzenesulfonyloxy, p-toluenesulfonyloxy, etc.

The "protecting group for hydroxy group" may be any conventional ones, and the protecting groups being easily and selectively removed includes, for example, a benzyl group, a t-butyl-
20 dimethylsilyl group, a triethylsilyl group, etc.

The "protecting group for amino group" may be any conventional ones, and includes, for example, a methoxycarbonyl group, an ethoxycarbonyl group, a 2,2,2-trichloroethoxycarbonyl group, a tert-butoxycarbonyl group, a benzyloxycarbonyl group, a vinyloxycarbonyl
25 group, a 9-fluorenylmethoxycarbonyl group, a formyl group, an acetyl

group, a trifluoroacetyl group, a benzoyl group, a phthalimide group, a p-toluenesulfonyl group, a benzenesulfonyl group, a methanesulfonyl group, and a benzyl group, etc., and preferable one is a tert-butoxy-carbonyl group.


5 The reaction temperature may vary according to the kinds of the starting compounds, and it is usually in the range of from about 50°C to about 200°C. The solvent includes, for example, aromatic hydrocarbons such as benzene, toluene, etc., ketone such as acetone, methyl ethyl ketone, etc., ethers such as tetrahydrofuran, dioxane, etc.,
10 alcohols such as ethanol, isopropanol, etc., acetonitrile, N,N-dimethyl-formamide, 1,3-dimethyl-2-imidazolidinone, etc., and these solvents may be used alone or in a mixture of two or more solvents.

 This reaction is preferably carried out in the presence of a base, and the base includes, for example, an inorganic base such as an alkali
15 metal carbonate (e.g., sodium carbonate, potassium carbonate, etc.), an alkali metal hydrogen carbonate (e.g., sodium hydrogen carbonate, potassium hydrogen carbonate, etc.), an alkali metal hydroxide (e.g., sodium hydroxide, potassium hydroxide, etc.), or an organic base such as triethylamine, tributylamine, N-methylmorpholine, etc. When the
20 compound wherein Y is a chlorine atom or bromine atom is used, the reaction may smoothly proceed by addition of an alkali metal iodide (e.g., sodium iodide, potassium iodide, etc.), or a tetraalkylammonium halide (e.g., tetra-n-butylammonium chloride, etc.) into the reaction system.

25 Process (g):

 Among the compounds of the present invention, the compound of the formula (I) wherein R¹ is a group of the formula: -X-R^{1c}-C(=O)N-

$R^{1a}R^{1b}$ may be prepared by condensing a compound of the formula (XXVIII):



(wherein X, R^{1e} , A^1 , A^2 , R^2 , R^4 , R^5 , R^6 , R^7 and Ar are as defined above; Q is the same as defined for the compound of the formula (I) except that W is replaced by a group of the formula: $ArCH(OA^1)CR^4(R^5)NA^2CR^6(R^7)-CH_2-$)

with a compound of the formula: $HNR^{1a}R^{1b}$ (wherein R^{1a} , R^{1b} are as defined above), followed by removing the protecting group, if necessary.

In this condensation reaction, the organic solvent to be used may be any inert organic solvents, for example, halogenated hydrocarbons (e.g., carbon tetrachloride, dichloromethane, 1,2-dichloroethane, etc.), ethers (e.g., diethyl ether, tetrahydrofuran, 1,4-dioxane, etc.), and N,N-dimethylformamide, etc.

The condensing agent to be used in the reaction includes, for example, condensing agents usually used in the peptide-bond formation (e.g., dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DIPC), N-ethyl-N'-3-dimethylaminopropylcarbodiimide (EDC=WSCI) and a hydrochloride thereof (WSCI·HCl), benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate salt (BOP), diphenylphosphoryl azide (DPPA), etc.), or carbonyl diimidazole (CDI), 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), triphenylphosphine-carbon tetrachloride, diethyl cyanophosphate, diphenyl phosphoroazide, etc.

In order to accelerate the reaction speed of the condensing

reaction or to avoid any side reactions, an additive such as N-hydroxy-succinimide (HONSu), 1-hydroxybenztriazole (HOBt), 3,4-dihydro-3-hydroxy-4-oxo-1,2,3-benzotriazine (HOOBt), etc. may be used. The reaction is preferably carried out at a temperature of from room
5 temperature to a refluxing temperature, for 30 minutes to 24 hours.

Process (h):

Among the compounds of the present invention, the compound of the formula (I) wherein R^1 is a group of the formula: $-X-R^{1e}-C(=O)-NR^{1a}R^{1b}$ may also be prepared by activating the carboxyl group of the
10 compound of the above formula (XXVIII) to give an acid halide, condensing said acid halide with a compound of the formula: $HNR^{1a}R^{1b}$ (wherein R^{1a} and R^{1b} are as defined above), following by removing the protecting group.

The compound of the formula (XXVIII) is converted into an acid
15 halide by reacting the compound of the formula (XXVIII) with triamide hexamethyl phosphite - carbon tetrachloride, triphenylphosphine-carbon tetrachloride, thionyl chloride, phosphorus oxychloride, phosphorus pentachloride, oxalyl chloride, etc. in a solvent. The reaction is usually carried out at room temperature for 30 minutes to 2 hours with stirring.
20 The solvent used herein includes, for example, tetrahydrofuran, dichloromethane, N,N-dimethylformamide, toluene, benzene, etc.

The organic solvent used in the condensation reaction may be any inert organic solvents, for example, halogenated hydrocarbons (e.g., carbon tetrachloride, dichloromethane, 1,2-dichloroethane, etc.), ethers
25 (e.g., diethyl ether, tetrahydrofuran, 1,4-dioxane, etc.), N,N-dimethylformamide, etc. The reaction may be carried out in the presence of a base. The reaction temperature is usually in the range of about -10°C


to 120°C, preferably in the range of about 0°C to 100°C. The reaction is usually carried out for 1 to 48 hours, preferably for 1 to 24 hours.

The base may be, for example, alkylamines (e.g., triethylamine, etc.), cyclic amines (e.g., N-methylmorpholine, pyridine, etc.), aromatic

amines (e.g., N,N-dimethylaniline, N,N-diethylaniline, etc.), alkali metal carbonates (e.g., sodium carbonate, potassium carbonate, etc.), alkali metal hydrogen carbonates (e.g., sodium hydrogen carbonate, potassium hydrogen carbonate, etc.).

Process (i):

Among the present compounds, the compound of the formula (I) wherein R¹ and R³ combine each other and form a divalent group of the formula: -X-R^{1e}-C(=O)- may be prepared by subjecting a compound of the formula (XXIX):



(wherein X, R^{1e}, A¹, A², R², R⁴, R⁵, R⁶, R⁷ and Ar are as defined above, R²⁰ is a lower alkyl group, Q is the same as defined for the compound of the formula (I) except that W is replaced by a group of the formula: ArCH(OA¹)CR⁴(R⁵)NA²CR⁶(R⁷)CH₂-) to intramolecular cyclization reaction, followed by removing the protecting group, if necessary.


This reaction is preferably carried out in the presence of a base, and the base includes, for example, an alkali metal carbonate (e.g., sodium carbonate, potassium carbonate, etc.), an alkali metal hydrogen carbonate (e.g., sodium hydrogen carbonate, potassium hydrogen carbonate, etc.), an alkali metal hydroxide (e.g., sodium hydroxide, potassium hydroxide, etc.), and an organic base such as triethylamine,

tributylamine, N-methylmorpholine, etc.

The reaction temperature may vary according to the kinds of the starting compound to be used, and it is usually in the range of about 50°C to about 200°C. The solvent includes, for example, aromatic hydrocarbons (e.g., benzene, toluene, etc.), ketones (e.g., acetone, methyl ethyl ketone, etc.), ethers (e.g., tetrahydrofuran, dioxane, etc.), alcohols (e.g., ethanol, isopropanol, etc.), acetonitrile, N,N-dimethylformamide, 1,3-dimethyl-2-imidazolidinone, etc., and these solvents may be used alone or in a mixture of two or more thereof.

Process (j):

Among the present compounds, the compound of the formula (I) wherein R^1 and R^3 combine each other and form a divalent group of the formula: $-X-R^{1e}-C(=O)-$ may be prepared by subjecting a compound of the formula (XXX):



(wherein X, R^{1e} , A^1 , A^2 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 and Ar are as defined above, Q is the same as defined for the compound of the formula (I) except that W is replaced by a group of the formula: $ArCH(OA^1)CR^4(R^5)NA^2CR^6(R^7)-CH_2-$)

to intramolecular cyclization reaction, followed by removing the protecting group, if necessary.

In this condensation reaction, the organic solvent may be an organic inert solvent, for example, halogenated hydrocarbons (e.g., carbon tetrachloride, dichloromethane, 1,2-dichloroethane, etc.), ethers (e.g., diethyl ether, tetrahydrofuran, 1,4-dioxane, etc.), N,N-dimethyl-

formamide, etc.

The condensing agent to be used in the reaction includes, for example, condensing agents usually used in the peptide-bond formation (e.g., dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DIPC),
5 N-ethyl-N'-3-dimethylaminopropylcarbodiimide (EDC=WSCI) and a hydrochloride thereof (WSCI·HCl), benzotriazol-1-yl-tris(dimethylamino)-phosphonium hexafluorophosphate salt (BOP), diphenylphosphoryl azide (DPPA), etc., or carbonyldiimidazole (CDI), 2-ethoxy-1-ethoxy-carbonyl-1,2-dihydroquinoline (EEDQ), triphenylphosphine-carbon
10 tetrachloride, diethyl cyanophosphate, diphenyl phosphorazide, etc.

In order to accelerate the reaction speed of the condensing reaction or to avoid any side reactions, an additive such as N-hydroxy-succinimide (HONSu), 1-hydroxybenzotriazole (HOBt), 3,4-dihydro-3-hydroxy-4-oxo-1,2,3-benzotriazine (HOOBt), etc. may be used.

15 The reaction is preferably carried out at a temperature of from room temperature to a refluxing temperature, for 30 minutes to 24 hours.

The compound of the present invention of the above formula (I) may be converted into a pharmaceutically acceptable salt thereof by a
20 conventional method. The pharmaceutically acceptable salt includes, for example, an acid addition salt with an inorganic acid such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, nitric acid, phosphoric acid, etc., an acid addition salt with an organic acid such as formic acid, acetic acid, methanesulfonic acid, benzene-
25 sulfonic acid, p-toluenesulfonic acid, propionic acid, citric acid, succinic acid, tartaric acid, fumaric acid, butyric acid, oxalic acid, malonic acid, maleic acid, lactic acid, malic acid, carbonic acid, glutamic acid,

aspartic acid, etc., a salt with an inorganic base such as sodium salt, potassium salt, calcium salt, etc., a salt with an organic base such as triethylamine, piperidine, morpholine, pyridine, lysine, etc.

5 The compound of the present invention includes solvates with a pharmaceutically acceptable solvent such as water or ethanol.

The compound of the present invention obtained in the above-mentioned Processes may be isolated and purified by a conventional isolation method such as recrystallization, a conventional purification method using chromatography, solvent extraction method, or
10 reprecipitation.

The product obtained in any one of Processes may be in the form of an acid addition salt thereof or a free base, due to the reaction conditions. These products may be converted into a desired acid addition salt or a free base by a conventional method.

15 When the compound of the present invention or the starting compound obtained in any one of Processes are in the form of a racemic mixture or a mixture of diastereomers, each stereoisomer may be separated by a conventional method, for example, by the method disclosed in EP-A-455006.

20 In addition to the specific protecting groups as exemplified in Processes as explained above, when each starting compound has a reactive group such as a carboxyl group, a hydroxy group or an amino group, the desired compound may be obtained by protecting these groups in advance by a suitable protecting group, and removing these
25 protecting groups after the reaction. The protection or deprotection can be carried out by a method disclosed in the literature: Green, T. W. and Wuts, P. G. M., PROTECTIVE GROUPS IN ORGANIC SYNTHESIS,

JOHN WILEY & SONS, INC. (1999), which should be selected according to the kinds of the protecting group to be removed.

The compound of the present invention may be administered either parenterally or orally when used as a medicament. Namely, the present compounds may be administered orally in the form of a conventional administration form such as powders, granules, tablets, capsules, syrups, and suspensions, or administered parenterally in the form of an injection form such as solutions, emulsions, suspension, etc. Further, the present compounds may also be administered rectally in the form of a suppository. The above-mentioned pharmaceutical preparations can be formulated, for example, in a conventional manner by mixing the present compound with conventional pharmaceutically acceptable carriers, excipients, binding agents, stabilizers, or diluents. When the present compound is used in the form of an injection form, the injection form may further contain, for example, pharmaceutically acceptable buffering agents, solubilizing agents, and isotonic agents. The dosage and the frequency of administration of the present compounds may vary, for example, according to the diseases to be treated, the conditions, ages, body weights of the patients and the administration form, etc., but the present compounds can usually be administered in a dose of 0.1 to 2000 mg per day in adult, preferably in a dose of 1 to 200 mg per day in adult, once a day, or divided into several (e.g., 2 - 4) dosage units.

EXAMPLES

The present invention will be illustrated in more detail by Reference Examples, Examples and Experiments, but should not be

construed to be limited thereto.

Reference Example 1

(R)-2-Chloro-1-pyridin-3-ylethanol

To a solution of (-)-B-chlorodiisopinocampheylborane [(-)-DIP-
5 Cl] (25 g, 77.9 mmol) in tetrahydrofuran (90 ml) are added with stirring
3-(2-chloroacetyl)pyridine hydrochloride (Can. J. Chem., vol. 61, p. 334
(1983)) (3.0 g, 15.6 mmol) and triethylamine (2.39 ml, 17.2 mmol) at
-25°C, and the reaction mixture is stirred at -25°C for 3 days. To this
mixture is added water (300 ml), and the mixture is warmed to room
10 temperature. To the mixture is added ethyl acetate, and the organic
phase is separated. The aqueous phase is neutralized with a saturated
aqueous sodium hydrogen carbonate solution, and extracted 6 times
with ethyl acetate. The combined organic phase is dried over sodium
sulfate, and concentrated under reduced pressure to give yellow oil,
15 which is purified by silica gel column chromatography (methanol/
chloroform = 1/20) to give the title compound (R)-2-chloro-1-pyridin-3-
ylethanol (2.02 g, yield: 82 %) as pale yellow oil.

¹H-NMR (CDCl₃) δ: 2.75 (1H, d, J=3.4Hz), 3.67 (1H, dd, J=8.5,
11.3Hz), 3.78 (1H, dd, J=3.5, 11.3Hz), 4.96-5.00 (1H,m), 7.33 (1H, dd,
20 J=4.9, 7.9Hz), 7.75-7.78 (1H, m), 8.59 (1H, dd, J=1.6, 4.8Hz), 8.64 (1H,
d, J=2.2Hz).

Reference Example 2

(R)-(Pyridin-3-yl)oxirane

To a solution of (R)-2-chloro-1-pyridin-3-ylethanol (2.0 g, 12.7
25 mmol) in acetonitrile (100 ml) is added potassium carbonate (7.02 g).
The mixture is refluxed for 2.5 hours, and cooled to room temperature.
The mixture is filtered, and the filtrate is distilled off under reduced

pressure. The resultant is purified by silica gel column chromatography (methanol/chloroform = 1/100) to give the title compound (R)-(pyridin-3-yl)oxirane (1.46 g, yield: 94 %) as pale yellow oil.

¹H-NMR (CDCl₃) δ: 2.83 (1H, dd, J=2.5, 5.3Hz), 3.21 (1H, dd, J=4.1, 5.3Hz), 3.90 (1H, dd, J=2.6, 4.0Hz), 7.29 (1H, dd, J=5.1, 8.3Hz), 7.53-7.56 (1H, m), 8.57 (1H, dd, J=1.6, 4.8Hz), 8.60 (1H, d, J=2.0Hz).

Reference Example 3

N,N-Diethyl-2-(1H-indol-7-yloxy)acetamide

To a solution of 7-hydroxyindole (2.65 g, 20 mmol) in acetone (20 ml) are added potassium carbonate (3.32 g), chloroacetic acid diethylamide (3.29 g, 22 mmol) and potassium iodide (0.33 g), and the mixture is refluxed for 4 hours. After cooling with ice, the insoluble materials are removed by filtration, and the filtrate is concentrated under reduced pressure to remove the solvent. To the residue are added chloroform (40 ml) and water (20 ml), and the mixture is stirred. The chloroform layer is separated, and dried over anhydrous magnesium sulfate. The solvent is evaporated under reduced pressure, and isopropyl ether (100 ml) is added to the residue. The precipitated crystals are collected by filtration, and dried to give the title compound N,N-diethyl-2-(1H-indol-7-yloxy)acetamide (4.37 g, yield: 89 %) as white crystals.

¹H-NMR (CDCl₃) δ: 1.18 (3H, t, J=7.1Hz), 1.23 (3H, t, J=7.2Hz), 3.36 (2H, q, J=7.2Hz), 3.45 (2H, q, J=7.1Hz), 4.81 (2H, s), 6.52 (1H, dd, J=2.2, 3.0Hz), 6.68 (1H, d, J=7.6Hz), 6.99 (1H, t, J=7.8Hz), 7.22 (1H, t, J=2.7Hz), 7.32 (1H, d, J=7.9Hz), 9.59 (1H, s).

Reference Example 4

(R)-2-[3-(2-Aminopropyl)-1H-indol-7-yloxy]-N,N-diethylacetamide

To a suspension of N-(9-fluorenylmethoxycarbonyl)-D-alanine (8.17 g, 26.3 mmol), methylene chloride (88 ml) and N,N-dimethylformamide (0.14 ml) is added dropwise oxalyl chloride (2.44 ml, 28 mmol) at room temperature with stirring, and the mixture is further stirred for one hour. The reaction solution is concentrated to dryness under reduced pressure to give an oil containing 9-fluorenylmethoxycarbonyl-D-Ala-Cl. A solution of N,N-diethyl-2-(1H-indol-7-yloxy)acetamide (4.31 g, 17.5 mmol) in methylene chloride (44 ml) is cooled to -20°C, and thereto is added dropwise a 3M solution of methyl magnesium bromide in diethyl ether (20.5 ml, 61.4 mmol). After the addition, the mixture is further stirred at -15°C for one hour. This reaction solution is cooled to -20 to -15°C, and thereto is added dropwise a solution of N-(9-fluorenylmethoxycarbonyl)-D-alanyl chloride in methylene chloride (53 ml) obtained above, and the mixture is further stirred at -20°C for 5.5 hours. The mixture is warmed to about -5°C, and poured into 2N hydrochloric acid. The organic layer is collected, washed with water (25 ml), and dried over anhydrous magnesium sulfate. The solvent is evaporated under reduced pressure to give an oil (13.3 g) containing (R)-[2-(7-diethylcarbamoylmethoxy-1H-indol-3-yl)-1-methyl-2-oxoethyl]-carbamic acid 9H-fluoren-9-ylmethyl ester. To a solution of the obtained oil in a mixture of acetonitrile (70 mL) and 2-propanol (4.0 ml) is added sodium borohydride (1.98 g, 52 mmol) in portions at room temperature with stirring, and the mixture is refluxed for 5 hours. The reaction solution is cooled to room temperature, and thereto is added dropwise methanol (120 ml). The reaction mixture is concentrated to dryness under reduced pressure. To the residue are added ethyl acetate and water, and the mixture is stirred. The organic layer is

separated, washed with a saturated brine, and dried over anhydrous sodium sulfate. The solvent is evaporated under reduced pressure, and the residue is purified by silica gel column chromatography (methanol/a saturated ammonia solution in chloroform = 1/20) to give the desired (R)-2-[3-(2-aminopropyl)-1H-indol-7-yloxy]-N,N-diethyl-
 5 acetamide (1.33 g, yield: 25 %).

¹H-NMR (CDCl₃) δ: 1.16 (3H, d, J=6.3Hz), 1.17 (3H, t, J=7.1Hz), 1.23 (3H, t, J=7.1Hz), 2.63 (1H, dd, J=8.2, 14.2Hz), 2.86 (1H, dd, J=4.4, 14.1Hz), 3.24-3.29 (1H, m), 3.36 (2H, q, J=7.1Hz), 3.45 (2H, q, J=7.1Hz),
 10 4.81 (2H, s), 6.68 (1H, d, J=7.6Hz), 6.99 (1H, t, J=7.8Hz), 7.06 (1H, d, J=1.8Hz), 7.29 (1H, d, J=8.0Hz), 9.36 (1H, s).

Example 1

N,N-Diethyl-2-{3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)-propyl]-1H-indol-7-yloxy}acetamide

15 A solution of (R)-(pyridin-3-yl)oxirane (97 mg, 0.80 mmol) and (R)-2-[3-(2-aminopropyl)-1H-indol-7-yloxy]-N,N-diethylacetamide (243 mg, 0.80 mmol) in water (0.8 mL) and ethanol (8 mL) is refluxed for 5 hours. After concentrated, the resulting residue is purified by preparative TLC (SiO₂, 20 x 20 cm, 0.5 mm, ⁱPrOH/saturated ammonia
 20 solution in chloroform = 1/10) to give the desired N,N-diethyl-2-{3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yloxy}-acetamide (70 mg, yield: 21 %).

¹H-NMR (CDCl₃) δ: 1.13 (3H, d, J=6.2Hz), 1.17 (3H, t, J=7.2Hz), 1.23 (3H, t, J=7.1Hz), 2.67 (1H, dd, J=9.4, 12.2Hz), 2.77-2.86 (2H, m),
 25 2.89 (1H, dd, J=3.5, 12.2Hz), 3.02-3.08 (1H, m), 3.35 (2H, q, J=7.1Hz), 3.44 (2H, q, J=7.1Hz), 4.57 (1H, dd, J=3.5, 9.3Hz), 4.80 (2H, s), 6.66 (1H, d, J=1.5Hz), 6.99 (1H, t, J=7.9Hz), 7.02 (1H, d, J=1.5Hz), 7.22-7.27

(2H, m), 7.65-7.68 (1H, m), 8.49 (1H, dd, J=1.6, 4.8Hz), 8.54 (1H, d, J=2.1Hz), 9.58 (1H, brs).

Example 2

{3-[(2R)-2-((2R)-2-Hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yloxy}acetic acid · 2 trifluoroacetate

A solution of N,N-diethyl-2-{3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yloxy}acetamide (30 mg, 0.071 mmol) and potassium hydroxide (120 mg, 2.1 mmol) in water (1 mL)/ethanol (1 mL) is stirred at 60°C for 6 hours. After cooling, to the mixture is added acetic acid (0.112 mL), and the mixture is concentrated. The residue is purified by preparative reversed phase HPLC (octadecylsilyl, trade name: Combiprep, ODS-A (YMC), inner diameter: 50 x 20 mm, particle size: 5 µm, pore size: 120 angstrom (hereinafter, this column is used in preparative reversed phase HPLC column), 0.05 % trifluoroacetic acid/water-0.035 % trifluoroacetic acid/acetonitrile) to give the desired {3-[(2R)-2-((2R)-2-hydroxy 2-pyridin-3-yl-ethylamino)propyl]-1H-indol-7-yloxy}acetic acid · 2 trifluoroacetate (24 mg, yield: 57 %).

¹H-NMR (CD₃OD) δ: 1.33 (3H, d, J=6.5Hz), 3.02 (1H, dd, J=9.1, 14.2Hz), 3.25-3.33 (2H, m), 3.40 (1H, dd, J=3.2, 12.8Hz), 3.65-3.70 (1H, m), 4.80 (2H, s), 5.22 (1H, dd, J=3.2, 9.9Hz), 6.62 (1H, d, J=7.6Hz), 6.97 (1H, t, J=7.9Hz), 7.20 (1H, s), 7.24 (1H, d, J=7.8Hz), 7.80 (1H, dd, J=5.7, 8.0Hz), 8.50 (1H, d, J=8.1Hz), 8.79 (1H, d, J=5.1Hz), 8.86 (1H, brs).

Reference Example 5

(R)-2-Bromo-1-[4-(phenylmethoxy)-3-[(methylsulfonyl)amino]phenyl]-ethanol

Under nitrogen atmosphere, a solution of 2-bromo-1-[4-(phenylmethoxy)-3-[(methylsulfonyl)amino]phenyl]ethanone (J. Med. Chem., vol.

23, no. 7, p. 738 (1980)) (977 mg, 2.45 mmol) and (R)-5,5-diphenyl-2-methyl-3,4-propano-1,3,2-oxazaborolidine (170 mg, 0.61 mmol) in tetrahydrofuran is stirred for 30 minutes. The mixture is cooled to -30°C, and thereto is added a 2M borane-dimethylsulfide complex in tetrahydrofuran (1.84 mL, 3.68 mmol), and the mixture is stirred at -25°C for 2 days. The reaction solution is warmed to room temperature, and ethyl acetate and a saturated aqueous ammonium chloride solution are added thereto, and the mixture is stirred. The organic layer is collected and dried over anhydrous sodium sulfate. The solvent is evaporated under reduced pressure, and the residue is purified by silica gel column chromatography (ethyl acetate/chloroform = 1/50) to give the desired (R)-2-bromo-1-[4-(phenylmethoxy)-3-[(methylsulfonyl)-amino]phenyl]ethanol (881 mg, yield: 90 %).

¹H-NMR (CDCl₃) δ: 2.64 (1H, d, J=3.3Hz), 2.93 (3H, s), 3.53 (1H, dd, J=8.7, 10.4Hz), 3.62 (1H, dd, J=3.5, 10.4Hz), 4.89 (1H, ddt, J=3.3, 3.5, 8.7Hz), 5.12 (2H, s), 6.82 (1H, brs), 7.00 (1H, d, J=8.5Hz), 7.18 (1H, dd, J=2.1, 8.5Hz), 7.36-7.44 (5H, m), 7.55 (1H, d, J=2.1Hz).

Reference Example 6

(R)-2-Iodo-1-[4-(phenylmethoxy)-3-[(methylsulfonyl)amino]phenyl]-ethanol

A mixture of (R)-2-bromo-1-[4-(phenylmethoxy)-3-[(methylsulfonyl)amino]phenyl]ethanol (880 mg, 2.2 mmol) and NaI (3.66 g, 24.4 mmol) is refluxed in acetone (35 mL) for one hour. After the mixture is filtered, the filtrate is concentrated, and the residue is separated into ethyl acetate and water. The organic phase is washed with a 25 % (w/w) aqueous sodium bisulfite solution, and dried over magnesium sulfate to give the desired (R)-2-iodo-1-[4-(phenylmethoxy)-3-[(methyl-

sulfonyl)amino]phenyl]ethanol (938 mg, yield: 95 %).

¹H-NMR (CDCl₃) δ: 2.46 (1H, d, J=3.7Hz), 2.94 (3H, s), 3.39 (1H, dd, J=8.4, 10.3Hz), 3.48 (1H, dd, J=3.8, 10.3Hz), 4.78 (1H, ddt, J=3.7, 3.8, 8.4Hz), 5.11 (2H, s), 6.82 (1H, brs), 7.00 (1H, d, J=8.4Hz), 7.17 (1H, dd, J=2.1, 8.4Hz), 7.36-7.44 (5H, m), 7.54 (1H, d, J=2.1Hz).

Reference Example 7

(R)-N-[2-Benzyloxy-5-(2-iodo-1-triethylsilyloxyethyl)phenyl]methanesulfonamide

To a solution of (R)-2-iodo-1-[4-(phenylmethoxy)-3-[(methylsulfonyl)amino]phenyl]ethanol (938 mg, 2.1 mmol), imidazole (393 mg, 5.77 mmol) and 4-dimethylaminopyridine (22.5 mg, 0.184 mmol) in dimethylformamide (10 mL) is added triethylsilyl chloride (0.38 mL, 2.2 mmol) with stirring. One hour thereafter, the reaction is complete, and the reaction solution is diluted with EtOAc (150 mL) and heptane (15 mL). The organic phase is washed with water, a saturated aqueous copper sulfate solution and a saturated brine, and dried over magnesium sulfate. The filtrate is concentrated under reduced pressure to give a solid, which is dissolved in ethyl acetate. The mixture is diluted with heptane, and the precipitated crystals are collected by filtration. The collected solid is washed with heptane and dried in vacuo to give the desired (R)-N-[2-benzyloxy-5-(2-iodo-1-triethylsilyloxyethyl)phenyl]methanesulfonamide (926 mg, yield: 79 %).

¹H-NMR (CDCl₃) δ: 0.52-0.62 (6H, m), 0.90 (9H, t, J=7.9Hz), 2.91 (3H, s), 3.31 (1H, dd, J=5.3, 10.1Hz), 3.34 (1H, dd, J=6.8, 10.1Hz), 4.73 (1H, dd, J=5.3, 6.8Hz), 5.10 (2H, s), 6.80 (1H, brs), 6.98 (1H, d, J=8.4Hz), 7.13 (1H, dd, J=2.1, 8.4Hz), 7.36-7.44 (5H, m), 7.53 (1H, d, J=2.1Hz).

Reference Example 8

2-(3-((2R)-2-[2-(4-Benzoyloxy-3-methanesulfonylaminophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indol-7-yloxy)-N,N-diethylacetamide

A solution of (R)-N-[2-benzoyloxy-5-(2-iodo-1-triethylsilyloxy-ethyl)phenyl]methanesulfonamide (185 mg, 0.33 mmol), (R)-2-[3-(2-aminopropyl)-1H-indol-7-yloxy]-N,N-diethylacetamide (100 mg, 0.33 mmol) and diisopropylethylamine (287 μ L, 1.65 mmol) in tetrahydrofuran (2 ml) is stirred at 110°C for 18 hours in an autoclave. After cooling, to the mixture is added ethyl acetate, and the mixture is washed with a saturated brine, dried over sodium sulfate, and concentrated. The residue is dissolved in dichloromethane (2 mL), and thereto is added trifluoroacetic acid (0.5 mL), and the mixture is stirred at room temperature for one hour. After concentrated, the resulting residue is dissolved in ethyl acetate, washed with a saturated aqueous sodium hydrogen carbonate solution, dried over sodium sulfate, and concentrated. The residue is purified by silica gel column chromatography (methanol/a saturated ammonia solution in chloroform = 1/20) to give the desired 2-(3-((2R)-2-[2-(4-benzoyloxy-3-methanesulfonylaminophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indol-7-yloxy)-N,N-diethylacetamide (144 mg, yield: 70 %).

$^1\text{H-NMR}$ (CDCl_3) δ : 1.12 (3H, d, $J=6.2\text{Hz}$), 1.17 (3H, t, $J=7.1\text{Hz}$), 1.23 (3H, t, $J=7.2\text{Hz}$), 2.65 (1H, dd, $J=8.9, 12.1\text{Hz}$), 2.76-2.89 (3H, m), 2.88 (3H, s), 2.98-3.05 (1H, m), 3.35 (2H, q, $J=7.2\text{Hz}$), 3.44 (2H, q, $J=7.1\text{Hz}$), 4.48 (1H, dd, $J=3.9, 9.1\text{Hz}$), 4.80 (2H, s), 5.09 (2H, s), 6.67 (1H, d, $J=7.6\text{Hz}$), 6.91-7.02 (3H, m), 7.11 (1H, dd, $J=2.0, 8.5\text{Hz}$), 7.24-7.28 (1H, m), 7.35-7.44 (6H, m), 9.39 (1H, brs).

Example 3

N,N-Diethyl-2-(3-((2R)-2-((2R)-2-hydroxy-2-(4-hydroxy-3-methanesulfonylaminophenyl)ethylamino)propyl)-1H-indol-7-yloxy)acetamide

To a solution of 2-(3-((2R)-2-[2-(4-benzyloxy-3-methanesulfonylaminophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indol-7-yloxy)-N,N-diethylacetamide (110 mg, 0.177 mmol) in methanol (10 mL) is added 10 % palladium on carbon (50 % wet, 30 mg), and the mixture is stirred under atmospheric hydrogen atmosphere for 12 hours. The catalyst is filtered through celite, and the filtrate is concentrated. The resulting crude product is crystallized from diisopropyl ether to give the desired N,N-diethyl-2-(3-((2R)-2-((2R)-2-hydroxy-2-(4-hydroxy-3-methanesulfonylaminophenyl)ethylamino)propyl)-1H-indol-7-yloxy)acetamide (80 mg, yield: 85 %).

¹H-NMR (CD₃OD) δ: 1.12 (3H, d, J=6.2Hz), 1.17 (3H, t, J=7.0Hz), 1.25 (3H, t, J=7.1Hz), 2.58 (1H, dd, J=6.7, 11.4Hz), 2.70-2.80 (3H, m), 2.80 (3H, s), 2.93-2.99 (1H, m), 3.47 (2H, q, J=6.9Hz), 3.59 (2H, q, J=7.0Hz), 4.51 (1H, t, J=6.8Hz), 4.90 (2H, s), 6.56-6.60 (2H, m), 6.75 (1H, dd, J=2.1, 8.3Hz), 6.86-6.92 (2H, m), 7.10 (1H, d, J=8.0Hz), 7.17 (1H, d, J=2.0Hz).

Example 4

(3-((2R)-2-((2R)-2-Hydroxy-2-(4-hydroxy-3-methanesulfonylamino-phenyl)ethylamino)propyl)-1H-indol-7-yloxy)acetic acid · trifluoroacetate

A solution of N,N-diethyl-2-(3-((2R)-2-((2R)-2-hydroxy-2-(4-hydroxy-3-methanesulfonylaminophenyl)ethylamino)propyl)-1H-indol-7-yloxy)acetamide (10 mg, 0.016 mmol) and potassium hydroxide (60 mg, 1.1 mmol) in water (0.5 mL)/ethanol (0.5 mL) is stirred at room temperature under argon atmosphere for one day. The mixture is purified by acetic acid (0.10 mL) preparative HPLC to give the desired

(3-((2R)-2-[(2R)-2-hydroxy-2-(4-hydroxy-3-methanesulfonylamino-phenyl)ethylamino]propyl)-1H-indol-7-yloxy)acetic acid · trifluoroacetate (3 mg, yield: 32 %).

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.5Hz), 2.92 (3H, s), 2.99 (1H, dd, J=9.0, 14.3Hz), 3.16-3.18 (2H, m), 3.22 (1H, dd, J=5.8, 14.8Hz), 3.60-3.64 (1H, m), 4.79 (2H, s), 4.82-4.88 (1H, m), 6.61 (1H, d, J=7.7Hz), 6.88 (1H, d, J=8.3Hz), 6.97 (1H, t, J=7.9Hz), 7.10 (1H, dd, J=2.1, 8.4Hz), 7.18 (1H, s), 7.22 (1H, d, J=8.0Hz), 7.38 (1H, d, J=2.1Hz).

Reference Example 9

2-[3-((2R)-2-{2-[4-(tert-Butyldimethylsilyloxy)phenyl]-(2R)-2-(triethylsilyloxy)ethylamino}propyl)-1H-indol-7-yloxy]-N,N-diethylacetamide (R)-1-(tert-Butyldimethylsilyloxy)-4-(2-iodo-1-triethylsilyloxy-ethyl)benzene (J. Org. Chem., vol. 56, p. 442 (1991)) (65 mg, 0.13 mmol) and (R)-2-[3-(2-aminopropyl)-1H-indol-7-yloxy]-N,N-diethylacetamide (80 mg, 0.26 mmol) and ¹Pr₂NEt (115 μL) are dissolved in tetrahydrofuran (2 mL), and the mixture is stirred at 110°C for 10 hours. After cooling, the mixture is concentrated and the residue is purified by silica gel column chromatography (chloroform → a saturated ammonia solution in chloroform) to give the desired 2-[3-((2R)-2-{2-[4-(tert-butyl-dimethylsilyloxy)phenyl]-(2R)-2-(triethylsilyloxy)ethylamino}propyl)-1H-indol-7-yloxy]-N,N-diethylacetamide (42 mg, yield: 48 %).

¹H-NMR (CDCl₃) δ: 0.18 (6H, s), 0.31 (6H, q, J=7.5Hz), 0.70 (9H, t, J=7.9Hz), 0.97 (9H, s), 1.18 (3H, t, J=7.2Hz), 1.23 (3H, t, J=7.2Hz), 1.38 (3H, d, J=6.4Hz), 2.92-3.14 (4H, m), 4.81 (2H, s), 4.96-4.99 (1H, m), 6.69 (1H, d, J=7.7Hz), 6.76 (2H, d, J=8.4Hz), 7.03 (1H, t, J=7.9Hz), 7.12 (2H, d, J=8.6Hz), 7.20 (1H, s), 7.27-7.30 (1H, m), 9.90 (1H, brs).

Example 5

N,N-Diethyl-2-(3-((2R)-2-((2R)-2-hydroxy-2-(4-hydroxyphenyl)ethylamino)propyl)-1H-indol-7-yloxy)acetamide

A solution of 2-[3-((2R)-2-(2-[4-(tert-butyldimethylsilyloxy)-phenyl)-(2R)-2-(triethylsilyloxy)ethylamino]-propyl)-1H-indol-7-yloxy]-N,N-diethylacetamide (40 mg, 0.060 mmol) and TBAF (261 mg, 1 mmol) in tetrahydrofuran (1 mL) is stirred at room temperature for 8 hours. After concentrated, the resultant is purified by preparative TLC (thickness: 0.5 mm, methanol/(a saturated ammonia solution in chloroform) = 1/10), extracted with saturated aqueous sodium hydrogen carbonate solution/chloroform. The organic layer is dried over sodium sulfate, and concentrated to give the desired N,N-diethyl-2-(3-((2R)-2-((2R)-2-hydroxy-2-(4-hydroxyphenyl)ethylamino)propyl)-1H-indol-7-yloxy)acetamide (15 mg, yield: 55 %).

¹H-NMR (CDCl₃) δ: 1.17 (3H, d, J=6.1Hz), 1.20 (3H, t, J=7.2Hz), 1.30 (3H, t, J=7.2Hz), 2.47 (1H, dd, J=8.1, 10.4Hz), 2.55 (1H, dd, J=10.1, 14.2Hz), 2.81 (1H, dd, J=3.8, 14.2Hz), 2.86-2.91 (1H, m), 3.10 (1H, dd, J=5.9, 10.5Hz), 3.36 (2H, q, J=7.1Hz), 3.44-3.52 (2H, m), 4.54 (1H, dd, J=6.1, 7.7Hz), 4.81 (1H, d, J=14.7Hz), 4.85 (1H, d, J=14.7Hz), 4.96-4.99 (1H, m), 6.44 (1H, d, J=7.6Hz), 6.45 (2H, d, J=8.5Hz), 6.66 (1H, s), 6.78 (2H, d, J=8.5Hz), 6.93 (1H, t, J=7.9Hz), 7.15 (1H, d, J=8.0Hz), 9.10 (1H, brs)

Example 6

(3-((2R)-2-((2R)-2-Hydroxy-2-(4-hydroxyphenyl)ethylamino)propyl)-1H-indol-7-yloxy)acetic acid

A solution of N,N-diethyl-2-(3-((2R)-2-((2R)-2-hydroxy-2-(4-hydroxyphenyl)ethylamino)propyl)-1H-indol-7-yloxy)acetamide (12 mg, 0.027 mmol) and potassium hydroxide (60 mg, 1.1mmol) in water (0.5

mL) and ethanol (0.5 mL) is stirred at 60°C for 6 hours. After cooling, the mixture is concentrated and acidified by addition of water (30 mL) and acetic acid. The resultant is purified by reversed phase column chromatography (COSMOSIL 75C18-OPN (Nacalai Tesque) (octadecyl-silica gel, particle diameter: 75 μ m, hereinafter, the same), water \rightarrow water/methanol = 10/1) to give the desired (3-((2R)-2-[(2R)-2-hydroxy-2-(4-hydroxyphenyl)ethylamino]propyl]-1H-indol-7-yloxy)acetic acid (7 mg, yield: 68 %).

$^1\text{H-NMR}$ (CD_3OD) δ : 1.27 (3H, d, $J=6.5\text{Hz}$), 2.95 (1H, dd, $J=8.1$, 14.3Hz), 3.02 (1H, dd, $J=3.9$, 12.4Hz), 3.08-3.13 (2H, m), 3.48-3.53 (1H, m), 4.48 (2H, s), 4.76 (1H, dd, $J=3.8$, 9.4Hz), 6.57 (1H, d, $J=7.6\text{Hz}$), 6.72 (2H, d, $J=8.6\text{Hz}$), 6.92 (1H, t, $J=7.9\text{Hz}$), 7.09 (1H, s), 7.13 (2H, d, $J=8.5\text{Hz}$), 7.14 (1H, d, $J=7.8\text{Hz}$).

Reference Example 10

tert-Butyl (R)-[2-(7-diethylcarbamoylmethoxy-1H-indol-3-yl)-1-methylethyl]carbamate

A mixture of (R)-2-[3-(2-aminopropyl)-1H-indol-7-yloxy]-N,N-diethylacetamide (0.30 g, 1 mmol) and di-tert-butyl dicarbonate (0.44 g, 2 mmol) and potassium carbonate (0.28 g, 2 mmol) in water (40 mL) and ethyl acetate (40 mL) is stirred at room temperature for 2 hours. The mixture is separated, and the organic layer is washed with a saturated brine, dried over magnesium sulfate, and concentrated to give crystals. The resulting crystals are filtered, washed with hexane, and dried under reduced pressure to give tert-butyl (R)-[2-(7-diethylcarbamoylmethoxy-1H-indol-3-yl)-1-methylethyl]carbamate (0.35 g, yield: 87 %).

$^1\text{H-NMR}$ (CDCl_3) δ : 1.10 (3H, d, $J=6.6\text{Hz}$), 1.17 (3H, t, $J=7.1\text{Hz}$),

1.23 (3H, d, J=7.1Hz), 1.44 (9H, s), 2.84 (1H, dd, J=6.6, 14.1Hz), 2.94 (1H, dd, J=6.0, 14.3Hz), 3.35 (2H, q, J=7.1Hz), 3.45 (2H, q, J=7.1Hz), 4.00 (1H, brs), 4.45 (1H, brs), 4.80 (2H, s), 6.66 (1H, d, J=7.7Hz), 6.99 (1H, t, J=7.8Hz), 7.03 (1H, d, J=2.0Hz), 7.30 (1H, d, J=8.0Hz), 9.58 (1H, brs).

Reference Example 11

(R)-[3-(2-(tert-Butoxycarbonylamino)propyl)-1H-indol-7-yloxy]acetic acid

A solution of tert-butyl (R)-[2-(7-diethylcarbamoylmethoxy-1H-indol-3-yl)-1-methylethyl]carbamate (0.35 g, 0.87 mmol) and potassium hydroxide (0.39 g, 7 mmol) in water (4 mL) and ethanol (6 mL) is stirred at 60°C for 8 hours. After cooling, the mixture is concentrated and acidified by addition of water (30 mL) and acetic acid. The resultant is extracted with chloroform, washed with water, dried over magnesium sulfate, and concentrated under reduced pressure to give the desired (R)-[3-(2-(tert-butoxycarbonylamino)propyl)-1H-indol-7-yloxy]acetic acid (0.30 g, yield: 100 %).

¹H-NMR (CD₃OD) δ: 1.10 (3H, d, J=6.3Hz), 1.43 (9H, s), 2.78 (1H, dd, J=7.5, 14.2Hz), 2.92 (1H, dd, J=4.6, 13.6Hz), 3.84-3.93 (1H, m), 4.79 (2H, s), 6.59 (1H, d, J=7.6Hz), 6.93 (1H, t, J=7.8Hz), 7.06 (1H, s), 7.26 (1H, d, J=8.0Hz).

Reference Example 12

tert-Butyl (R)-{2-[7-(2-methanesulfonylamino-2-oxoethoxy)-1H-indol-3-yl]-1-methylethyl}carbamate

A solution of (R)-[3-(2-(tert-butoxycarbonylamino)propyl)-1H-indol-7-yloxy]acetic acid (0.30 g, 0.86 mmol) and carbonyldiimidazole (0.21 g, 1.3 mmol) in dimethylformamide (10 mL) is stirred at room temperature for 3 days. To the mixture are added methanesulfonamide

(0.16 g, 1.7 mmol) and 1,8-diazabicyclo[5.4.0]undeca-7-ene (0.26 mL, 1.7 mmol), and the mixture is stirred at 50°C for 4 hours. After cooling, to the mixture is added water (0.1 mL), and the mixture is concentrated. The residue is purified by silica gel column chromatography (methanol/
 5 chloroform = 1/10 → 1/3), and the resultant is dissolved in ethyl acetate, washed with a 0.05N hydrochloric acid (x 2) and saturated brine, and further dried over magnesium sulfate and concentrated to give the desired tert-butyl (R)-{2-[7-(2-methanesulfonylamino-2-oxoethoxy)-1H-indol-3-yl]-1-methylethyl}carbamate (0.14 g, yield: 38 %).

10 ¹H-NMR (CD₃OD) δ: 1.07 (3H, d, J=6.3Hz), 1.40 (9H, s), 2.75 (1H, dd, J=7.4, 14.2Hz), 2.89 (1H, dd, J=3.8, 13.3Hz), 3.17 (3H, s), 3.83-3.88 (1H, m), 4.64 (2H, s), 6.56 (1H, d, J=7.7Hz), 6.89 (1H, t, J=7.9Hz), 7.03 (1H, s), 7.22 (1H, d, J=7.9Hz).

Reference Example 13

15 (R)-N-{2-[3-(2-Aminopropyl)-1H-indol-7-yloxy]acetyl}methanesulfonamide · trifluoroacetate

A solution of tert-butyl (R)-{2-[7-(2-methanesulfonylamino-2-oxoethoxy)-1H-indol-3-yl]-1-methylethyl}carbamate (0.14 g, 0.33 mmol) and trifluoroacetic acid (1 mL) in methylene chloride (4 mL) is stirred at
 20 room temperature for 8 hours. The mixture is concentrated to give the desired (R)-N-{2-[3-(2-aminopropyl)-1H-indol-7-yloxy]acetyl}methanesulfonamide · trifluoroacetate (0.13 g, yield: 90 %).

25 ¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.6Hz), 2.98-3.06 (2H, m), 3.31 (3H, s), 3.55-3.60 (1H, m), 4.80 (2H, s), 6.64 (1H, d, J=7.7Hz), 6.99 (1H, t, J=7.9Hz), 7.18 (1H, s), 7.24 (1H, d, J=7.6Hz)

Example 7

N-[2-(3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-

1H-indol-7-yloxy)acetyl]methanesulfonamide · trifluoroacetate

A mixture of (R)-N-{2-[3-(2-aminopropyl)-1H-indol-7-yloxy]-acetyl]methanesulfonamide · trifluoroacetate (40 mg, 0.091 mmol), (R)-(+)-3-chlorostyren oxide (24 μ L, 0.18 mmol) and ⁱPr₂NEt (24 μ L, 0.137 mmol) in acetonitrile (1 mL) is refluxed for 8 hours. After cooling, the mixture is concentrated and purified by preparative TLC (thickness: 0.5 mm, methanol/(a saturated ammonia solution in chloroform) = 1/3) and preparative reversed phase HPLC (trifluoroacetic acid/water/acetonitrile) to give the desired N-[2-(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indol-7-yloxy)acetyl]methanesulfonamide · trifluoroacetate (9 mg, yield: 21 %).

¹H-NMR (CD₃OD) δ : 1.31 (3H, d, J=6.5Hz), 3.01 (1H, dd, J=9.1, 14.3Hz), 3.16 (1H, dd, J=10.1, 12.7Hz), 3.21-3.28 (2H, m), 3.31 (3H, s), 3.62-3.67 (1H, m), 4.79 (2H, s), 4.93 (1H, dd, J=3.3, 9.9Hz), 6.64 (1H, d, J=7.7Hz), 6.99 (1H, t, J=7.9Hz), 7.21 (1H, s), 7.27 (1H, d, J=7.9Hz), 7.30-7.38 (3H, m), 7.44 (1H, s).

Example 8

N-(2-{3-[(2R)-2-((2R)-2-Hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yloxy}acetyl]methanesulfonamide · 2 trifluoroacetate

A mixture of (R)-N-{2-[3-(2-aminopropyl)-1H-indol-7-yloxy]-acetyl]methanesulfonamide · trifluoroacetate (40 mg, 0.091 mmol), (R)-(pyridin-3-yl)oxirane (48 mg, 0.36 mmol) and ⁱPr₂NEt (24 μ L, 0.137 mmol) in acetonitrile (1 mL) is refluxed for 3 hours. After cooling, the mixture is concentrated and purified by preparative reversed phase HPLC (trifluoroacetic acid/water/acetonitrile) to give the desired N-(2-{3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yloxy}acetyl]methanesulfonamide · 2 trifluoroacetate (3 mg, yield: 3 %).

¹H-NMR (CD₃OD) δ: 1.32 (3H, d, J=6.5Hz), 3.02 (1H, dd, J=9.1, 14.2Hz), 3.26-3.33 (5H, m), 3.38 (1H, dd, J=3.4, 12.8Hz), 3.65-3.70 (1H, m), 4.80 (2H, s), 5.18 (1H, dd, J=3.2, 9.9Hz), 6.64 (1H, d, J=7.6Hz), 6.99 (1H, t, J=7.9Hz), 7.22 (1H, s), 7.27 (1H, d, J=8.0Hz), 7.90 (1H, dd, J=5.5, 7.8Hz), 8.39 (1H, d, J=8.1Hz), 8.74 (1H, d, J=4.0Hz), 8.81 (1H, s).

Reference Example 14

1H-Indole-7-carboxylic acid diethylamide

A solution of 1H-indole-7-carboxylic acid (0.48 g, 3 mmol), diethylamine (0.93 mL, 9 mmol), 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (1.74 g, 9 mmol), 1-hydroxybenzotriazole (1.23 g, 9 mmol), triethylamine (1.26 mL, 12 mmol) in dimethylformamide (30 mL) is stirred at room temperature for 4 hours. The mixture is diluted with ethyl acetate, washed with 1N HCl, water, a saturated aqueous sodium hydrogen carbonate solution, water, and a saturated brine, dried over magnesium sulfate, and concentrated to give the desired 1H-indole-7-carboxylic acid diethylamide (0.64 g, yield: 99 %).

¹H-NMR (CDCl₃) δ: 1.27 (6H, t, J=7.1Hz), 3.56 (4H, q, J=7.1Hz), 6.56 (1H, dd, J=2.2, 3.2Hz), 7.09 (1H, t, J=7.5Hz), 7.23 (1H, dd, J=0.7, 7.3Hz), 7.25-7.27 (1H, m), 7.70 (1H, d, J=7.9Hz), 9.06 (1H, brs).

Reference Example 15

(R)-[2-(7-Diethylcarbamoyl-1H-indol-3-yl)-1-methyl-2-oxoethyl]carbamic acid 9H-fluoren-9-ylmethyl ester

To a suspension of N-(9-fluorenylmethoxycarbonyl)-D-alanine (21.61 g, 69.4 mmol), methylene chloride (230 mL) and N,N-dimethylformamide (0.37 mL) is added dropwise oxalyl chloride (6.46 mL, 74.0 mmol) at room temperature with stirring, and the mixture is further stirred for 2 hours. The reaction solution is concentrated to dryness

under reduced pressure to give an oil containing N-(9-fluorenylmethoxycarbonyl)-D-alanyl chloride. A solution of 1H-indole-7-carboxylic acid diethylamide (10.0 g, 46.3 mmol) in methylene chloride (120 ml) is cooled to -20°C, and thereto is added dropwise a 3M solution of methyl magnesium bromide/diethyl ether (54 ml, 162 mmol). After the addition, the mixture is further stirred at -20°C for 2 hours. This reaction solution is cooled to -25 – -15°C, and thereto is added dropwise a solution of the above obtained N-(9-fluorenylmethoxycarbonyl)-D-alanyl in methylene chloride (140 ml), and the mixture is further stirred at -20°C for 2 hours. The mixture is warmed to about -5°C, and poured into 2N hydrochloric acid. The organic layer is separated, washed with water (25 ml) and dried over anhydrous magnesium sulfate. The solvent is evaporated under reduced pressure, and the residue is purified by silica gel column chromatography (ethyl acetate/hexane = 1/2 → 5/1) to give the desired (R)-[2-(7-diethylcarbamoyl-1H-indol-3-yl)-1-methyl-2-oxoethyl]carbamic acid 9H-fluoren-9-ylmethyl ester (10.24 g, yield: 43 %).

¹H-NMR (CDCl₃) δ: 1.22-1.30 (6H, m), 1.38 (3H, d, J=7.0Hz), 3.56 (brs, 4H), 4.22 (1H, t, J=7.2Hz), 4.36 (2H, d, J=7.2Hz), 4.96-4.99 (1H, m), 6.01 (1H, d, J=6.2Hz), 7.27-7.33 (4H, m), 7.39 (2H, t, J=7.5Hz), 7.59-7.62 (3H, m), 7.75 (2H, d, J=7.5Hz), 8.42 (1H, dd, J=2.5, 6.3Hz), 10.71 (1H, brs).

Reference Example 16

(R)-3-(2-Aminopropyl)-1H-indole-7-carboxylic acid diethylamide

A solution of (R)-[2-(7-diethylcarbamoyl-1H-indol-3-yl)-1-methyl-2-oxoethyl]carbamic acid 9H-fluoren-9-ylmethyl ester (10.24 g, 20.1 mmol) in a mixture of acetonitrile (134 mL) and 2-propanol (7.7 ml)

is added sodium borohydride (3.80 g, 100 mmol) in portions at room temperature with stirring, and the mixture is refluxed for 4 hours. The reaction solution is cooled to room temperature, and thereto is added dropwise methanol (200 ml). The reaction mixture is concentrated to dryness under reduced pressure. To the residue are added ethyl acetate and water, and the mixture is stirred. The organic layer is separated, washed with a saturated brine, and dried over anhydrous sodium sulfate. The solvent is evaporated under reduced pressure, and the residue is purified by silica gel column chromatography (methanol/(a saturated ammonia solution in chloroform) = 1/100) to give the desired (R)-3-(2-aminopropyl)-1H-indole-7-carboxylic acid diethylamide (3.60 g, yield: 66 %).

¹H-NMR (CDCl₃) δ: 1.14 (3H, d, J=6.3Hz), 1.24 (6H, t, J=6.9Hz), 2.60-2.65 (1H, dd, J=8.1, 14.1Hz), 2.82-2.87 (1H, dd, J=4.9, 14.2Hz), 3.19-3.27 (1H, m), 3.51-3.53 (4H, m), 7.01 (1H, s), 7.04-7.08 (1H, m), 7.18 (1H, d, J=7.3Hz), 7.64 (1H, d, J=6.3Hz), 9.35 (1H, brs)

Example 9

3-{{(2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl}-1H-indole-7-carboxylic acid diethylamide

A mixture of (R)-3-(2-aminopropyl)-1H-indole-7-carboxylic acid diethylamide (1.30 g, 4.76 mmol) and (R)-(+)-3-chlorostyrene oxide (1.21 mL, 9.52 mmol) in acetonitrile (15 mL) is refluxed for 10 hours. After cooling, the mixture is concentrated and the residue is purified by silica gel column chromatography (a saturated ammonia solution in chloroform) to give the desired 3-{{(2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl}-1H-indole-7-carboxylic acid diethylamide (0.79 g, yield: 39 %).

¹H-NMR (CDCl₃) δ: 1.10 (3H, d, J=6.2Hz), 1.25 (6H, t, J=7.0Hz), 2.63 (1H, dd, J=9.2, 12.0Hz), 2.78-2.85 (3H, m), 2.98-3.03 (1H, m), 3.51-3.56 (4H, m), 4.49 (1H, dd, J=3.4, 9.0Hz), 7.01 (1H, s), 7.07 (1H, t, J=7.7Hz), 7.14-7.25 (4H, m), 7.32 (1H, s), 7.62 (1H, d, J=7.9Hz), 9.17 (1H, brs).

Example 10

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid

A solution of 3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid diethylamide (175 mg, 0.41 mmol) in 1,4-dioxane (7 mL) and 6N hydrochloric acid (7 mL) is stirred with heating at 150°C for 7 hours in a closed system. After cooling, the mixture is evaporated under reduced pressure to remove dioxane, and the resultant is neutralized with an aqueous Na₂CO₃ solution. The precipitates are filtered, and purified by reversed phase column (octadecylsilyl, methanol/water = 1/1 → 3/1) to give the desired 3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid (45 mg, yield: 30 %).

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.4Hz), 3.02 (1H, dd, J=8.1, 14.3Hz), 3.11-3.21 (3H, m), 3.53-3.59 (1H, m), 7.07 (1H, t, J=7.8Hz), 7.22-7.34 (4H, m), 7.42 (1H, s), 7.68 (1H, d, J=7.9Hz), 7.78 (1H, d, J=7.4Hz).

Example 11

Methyl (2S)-2-[(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carbonyl)amino]-4-methylpentanoate trifluoroacetate

A solution of 3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethyl-

amino]propyl}-1H-indole-7-carboxylic acid (22 mg, 0.059 mmol), L-Leu-OMe · HCl (107 mg, 0.59 mmol), 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (113 mg, 0.59 mmol), 1-hydroxybenztriazole (80 mg, 0.59 mmol), triethylamine (164 μ L, 1.18 mmol) in dimethylformamide (2 mL) is stirred at room temperature for 18 hours. The mixture is diluted with ethyl acetate, washed with a saturated aqueous sodium hydrogen carbonate solution, water, and a saturated brine, dried over sodium sulfate, and concentrated. The residue is purified by silica gel column chromatography (a saturated ammonia solution in chloroform) and preparative reversed phase HPLC (octadecylsilyl, trifluoroacetic acid/acetonitrile/water) to give the desired methyl (2S)-2-[(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carbonyl)amino]-4-methylpentanoate · trifluoroacetate (23 mg, yield: 63 %).

$^1\text{H-NMR}$ (CD_3OD) δ : 0.98 (3H, d, $J=6.2\text{Hz}$), 1.00 (3H, d, $J=6.3\text{Hz}$), 1.31 (3H, d, $J=6.5\text{Hz}$), 1.69-1.88 (3H, m), 3.04 (1H, dd, $J=9.2, 14.1\text{Hz}$), 3.14-3.30 (3H, m), 3.63-3.69 (1H, m), 3.73 (3H, s), 4.75 (1H, dd, $J=4.4, 10.0\text{Hz}$), 4.95 (1H, dd, $J=3.1, 9.9\text{Hz}$), 7.16 (1H, t, $J=7.7\text{Hz}$), 7.30-7.37 (4H, m), 7.45 (1H, s), 7.72 (1H, d, $J=7.5\text{Hz}$), 7.82 (1H, d, $J=7.9\text{Hz}$).

Examples 12 to 23

The compounds of the following Examples 12 to 23 are obtained in a similar manner to Example 11.

Example 12

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid dibutylamide · trifluoroacetate

$^1\text{H-NMR}$ (CD_3OD) δ : 0.61 (3H, brs), 1.02 (5H, brs), 1.30 (3H, d,

J=6.5Hz), 1.46 (4H, brs), 1.73 (2H, brs), 3.01 (1H, dd, J=9.3, 14.2Hz), 3.17 (1H, dd, J=10.1, 12.5Hz), 3.25-3.31 (6H, m), 3.57 (2H, brs), 3.66 (1H, m), 4.95 (1H, dd, J=3.1, 10.0Hz), 7.11 (2H, m), 7.26 (1H, s), 7.34 (3H, m), 7.47 (1H, m), 7.68 (1H, dd, J=1.7, 7.2Hz).

5 Example 13

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid (3-methylbutyl)amide · trifluoroacetate

¹H-NMR (CD₃OD) δ: 0.97 (6H, d, J=6.6Hz), 1.31 (3H, d, J=6.5Hz), 1.55 (2H, dt, J=7.3, 8.7Hz), 1.69 (1H, qq, J=6.7Hz), 3.04 (2H, dd, J=9.1, 14.3 Hz), 3.27 (2H, m), 3.45 (1H, t, J=14.9Hz), 3.66 (1H, m), 4.93 (1H, dd, J=3.3, 8.0Hz), 7.13 (1H, t, J=7.7Hz), 7.34 (4H, m), 7.45 (1H, m), 7.60 (1H, m), 7.79 (1H, m).

Example 14

15 (3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indol-7-yl)-piperidin-1-ylmethanone · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.5Hz), 1.41-1.65 (6H, m), 3.02 (1H, dd, J=9.2, 14.2Hz), 3.17 (1H, dd, J=10.1, 12.5Hz), 3.27 (2H, m), 3.30-3.50 (2H, m), 3.66-3.88 (2H, m), 4.94 (1H, dd, J=3.1, 10.0Hz), 7.15 (2H, m), 7.27 (1H, s), 7.34 (3H, m), 7.47 (1H, m), 7.70 (1H, dd, J=2.1, 6.9Hz).

Example 15

(3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indol-7-yl)-morpholin-4-ylmethanone · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 3.02 (1H, dd, J=9.2, 14.2Hz), 3.17 (1H, dd, J=10.1, 12.5Hz), 3.25-3.35 (4H, m), 3.66-3.76 (6H, m), 4.94 (1H, dd, J=3.1, 10.0Hz), 7.17 (2H, m), 7.27 (1H, s), 7.34 (3H, m), 7.47 (1H, m), 7.72 (1H, dd, J=1.3, 7.7Hz).

Example 16

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid ((1S)-1-hydroxymethyl-3-methylbutyl)amide · trifluoroacetate

5 $^1\text{H-NMR}$ (CD_3OD) δ : 0.97 (6H, d, $J=6.6\text{Hz}$), 1.45 (3H, d, $J=9.2\text{Hz}$), 1.49 (1H, m), 1.51 (1H, m), 1.56 (1H, m), 3.04 (2H, dd, $J=9.1$, 14.3Hz), 3.27 (2H, m), 3.61 (3H, d, $J=5.6\text{Hz}$), 3.66 (1H, m), 4.30 (1H, tt, $J=4.7\text{Hz}$), 4.93 (1H, dd, $J=3.2$, 10.2Hz), 7.14 (1H, t, $J=7.7\text{Hz}$), 7.34 (4H, m), 7.45 (1H, m), 7.65 (1H, m), 7.79 (1H, m).

10 Example 17

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxamide · trifluoroacetate

15 $^1\text{H-NMR}$ (CD_3OD) δ : 1.31 (3H, d, $J=6.5\text{Hz}$), 3.04 (1H, dd, $J=9.1$, 14.3 Hz), 3.17 (2H, m), 3.25-3.35 (3H, m), 3.66 (1H, m), 4.94 (1H, dd, $J=3.1$, 10.0Hz), 7.13 (1H, t, $J=7.6\text{Hz}$), 7.45 (4H, m), 7.67 (1H, m), 7.82 (1H, dd, $J=1.3$, 7.7Hz).

Example 18

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid methanamide · trifluoroacetate

20 $^1\text{H-NMR}$ (CD_3OD) δ : 1.31 (3H, d, $J=6.5\text{Hz}$), 2.96 (3H, s), 3.04 (2H, dd, $J=9.1$, 14.3Hz), 3.27-3.35 (3H, m), 3.65 (3H, m), 4.93 (1H, dd, $J=3.2$, 10.2Hz), 7.13 (1H, t, $J=7.7\text{Hz}$), 7.34 (4H, m), 7.45 (1H, m), 7.57 (1H, m), 7.79 (1H, m).

Example 19

25 3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid dimethanamide · trifluoroacetate

$^1\text{H-NMR}$ (CD_3OD) δ : 1.31 (3H, d, $J=6.5\text{Hz}$), 2.95 (3H, brs), 3.04

(2H, dd, J=9.1, 14.3Hz), 3.10-3.35 (6H, m), 3.76 (1H, m), 4.93 (1H, dd, J=3.2, 10.2Hz), 7.13 (1H, t, J=7.7Hz), 7.27 (1H, m), 7.45 (3H, m), 7.47 (1H, m), 7.71 (1H, m).

Example 20

5 3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid ethylamide · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.25 (3H, t, J=7.2Hz), 1.31 (3H, d, J=6.5Hz), 3.04 (1H, dd, J=9.1, 14.3Hz), 3.16-3.35 (3H, m), 3.46 (2H, q, J=7.2Hz), 3.76 (1H, m), 4.93 (1H, m), 7.13 (1H, t, J=7.7Hz), 7.27 (3H, m), 7.45 (1H, m), 7.60 (1H, m), 7.79 (1H, m).

Example 21

Ethyl (2S)-2-[(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carbonyl)amino]propionate · trifluoroacetate

15 ¹H-NMR (CD₃OD) δ: 1.27 (3H, t, J=7.2Hz), 1.31 (3H, d, J=6.5Hz), 1.52 (3H, d, J=7.3Hz), 3.04 (1H, dd, J=9.1, 14.3Hz), 3.16-3.35 (3H, m), 3.66 (1H, m), 4.18 (2H, q, J=7.2Hz), 4.66 (1H, q, J=7.3Hz), 4.93 (1H, d, J=3.7Hz, 9.8Hz), 7.16 (1H, t, J=7.7Hz), 7.27 (3H, m), 7.45 (1H, m), 7.73 (1H, m), 7.81 (1H, m).

Example 22

20 Ethyl 3-[(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carbonyl)amino]propionate · trifluoroacetate

25 ¹H-NMR (CD₃OD) δ: 1.23 (3H, t, J=7.1Hz), 1.31 (3H, d, J=6.5Hz), 2.68 (3H, d, J=6.9Hz), 3.04 (1H, dd, J=9.1, 14.3Hz), 3.16-3.35 (3H, m), 3.66 (1H, m), 3.69 (2H, t, J=6.9Hz), 4.14 (2H, q, J=7.1Hz), 4.93 (1H, d, J=3.3Hz, 9.8Hz), 7.14 (1H, t, J=7.7Hz), 7.27 (3H, m), 7.45 (1H, m), 7.58 (1H, m), 7.81 (1H, m).

Example 23

Ethyl [(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carbonyl)amino]acetate · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.27 (3H, t, J=7.1Hz), 1.29 (3H, d, J=6.5Hz), 3.04 (1H, dd, J=9.1, 14.3Hz), 3.16-3.35 (3H, m), 3.66 (1H, m), 3.69 (2H, t, J=6.9Hz), 4.15 (2H, s), 4.22 (2H, q, J=7.1Hz), 4.93 (1H, d, J=3.3Hz, 9.8Hz), 7.14 (1H, t, J=7.7Hz), 7.27 (3H, m), 7.45 (1H, m), 7.66 (1H, m), 7.81 (1H, m).

Example 24

(2S)-2-[(3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carbonyl)amino]-4-methylpentanoic acid · trifluoroacetate

A solution of methyl (2S)-2-[(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carbonyl)amino]-4-methylpentanoate · trifluoroacetate (20 mg, 0.0326 mmol) and potassium hydroxide (60 mg, 1.1 mmol) in water (0.5 mL) and methanol (2 mL) is stirred at room temperature for 3 hours. The mixture is acidified with 1N hydrochloric acid, and purified by preparative reversed phase HPLC (octadecylsilyl, trifluoroacetic acid/acetonitrile/water) to give the desired (2S)-2-[(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carbonyl)amino]-4-methylpentanoic acid · trifluoroacetate (18 mg, yield: 92 %).

¹H-NMR (CD₃OD) δ: 0.98-1.01 (6H, m), 1.31 (3H, d, J=6.5Hz), 1.72-1.89 (3H, m), 3.05 (1H, dd, J=9.0, 14.3Hz), 3.14-3.31 (3H, m), 3.64-3.67 (1H, m), 4.74 (1H, dd, J=4.1, 10.2Hz), 4.94 (1H, dd, J=3.3, 9.9Hz), 7.16 (1H, t, J=7.7Hz), 7.29-7.40 (4H, m), 7.45 (1H, s), 7.72 (1H, d, J=7.4Hz), 7.82 (1H, d, J=7.9Hz).

Examples 25 to 27

The compounds of the following Examples 25 to 27 are obtained

in a similar manner to Example 24.

Example 25

Ethyl (2S)-2-[(3-[(2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl]-1H-indole-7-carbonyl)amino]propionate · trifluoroacetate

5 $^1\text{H-NMR}$ (CD_3OD) δ : 1.31 (3H, d, $J=6.5$ Hz), 3.04 (1H, dd, $J=9.1$, 14.3 Hz), 3.16-3.35 (3H, m), 3.66 (1H, m), 4.18 (2H, q, $J=7.2$ Hz), 4.67 (1H, q, $J=7.4$ Hz), 4.93 (1H, dd, $J=3.4$, 9.7Hz), 7.16 (1H, t, $J=7.7$ Hz), 7.27 (3H, m), 7.45 (1H, m), 7.72 (1H, m), 7.81 (1H, m).

Example 26

10 3-[(3-[(2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl]-1H-indole-7-carbonyl)amino]propionic acid · trifluoroacetate

$^1\text{H-NMR}$ (CD_3OD) δ : 1.31 (3H, d, $J=6.5$ Hz), 2.67 (3H, d, $J=6.9$ Hz), 3.04 (1H, dd, $J=9.1$, 14.3Hz), 3.16-3.35 (3H, m), 3.66 (1H, m), 3.69 (2H, t, $J=6.9$ Hz), 4.93 (1H, dd, $J=3.3$, 9.8Hz), 7.14 (1H, t, $J=7.7$ Hz), 7.27 (3H, m), 7.45 (1H, m), 7.58 (1H, m), 7.81 (1H, m).

Example 27

[(3-[(2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl]-1H-indole-7-carbonyl)amino]acetic acid · trifluoroacetate

20 $^1\text{H-NMR}$ (CD_3OD) δ : 1.31 (3H, d, $J=6.5$ Hz), 3.04 (1H, dd, $J=9.1$, 14.3Hz), 3.16-3.35 (3H, m), 3.66 (1H, m), 3.69 (2H, t, $J=6.9$ Hz), 4.15 (2H, s), 4.93 (1H, dd, $J=3.3$, 9.8Hz), 7.14 (1H, t, $J=7.7$ Hz), 7.27 (3H, m), 7.45 (1H, m), 7.66 (1H, m), 7.81 (1H, m).

Example 28

25 3-[(2R)-2-[(2R)-2-Hydroxy-2-pyridin-3-ylethylamino]propyl]-1H-indole-7-carboxylic acid diethylamide

The title compound is prepared from (R)-3-(2-aminopropyl)-1H-indole-7-carboxylic acid diethylamide in a similar manner to Example 1.

¹H-NMR (CDCl₃) δ: 1.13 (3H, d, J=6.2Hz), 1.27 (6H, t, J=7.1Hz), 2.65 (1H, dd, J=9.4, 12.2Hz), 2.83-2.90 (3H, m), 3.02-3.07 (1H, m), 3.55 (4H, q, J=7.0Hz), 4.55 (1H, dd, J=3.5, 9.3Hz), 7.06-7.10 (2H, m), 7.21-7.26 (2H, m), 7.63-7.68 (2H, m), 8.49 (1H, dd, J=1.6, 4.8Hz), 8.54 (1H, d, J=2.1Hz), 9.03 (1H, brs).

Reference Example 17

3-{{(2R)-2-[(5R)-5-(3-Chlorophenyl)-2-oxooxazolidin-3-yl]propyl}-1H-indole-7-carboxylic acid diethylamide

To a solution of 3-{{(2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl}-1H-indole-7-carboxylic acid diethylamide · hydrochloride (47 mg, 0.1 mmol) and triethylamine (139 μL, 1 mmol) in tetrahydrofuran (3 mL) is added triphosgene (12 mg, 0.04 mmol), and the mixture is stirred at room temperature for 4 hours. To the mixture is added a saturated aqueous sodium hydrogen carbonate solution, and the mixture is extracted with ethyl acetate. The organic layer is washed with a saturated brine, dried over sodium sulfate, and concentrated. The residue is purified by silica gel column chromatography (methanol/chloroform = 1/50) to give 3-{{(2R)-2-[(5R)-5-(3-chlorophenyl)-2-oxooxazolidin-3-yl]propyl}-1H-indole-7-carboxylic acid diethylamide (45 mg, yield: 100 %).

¹H-NMR (CDCl₃) δ: 1.25-1.30 (9H, m), 2.90-3.01 (2H, m), 3.26 (1H, dd, J=6.8, 8.6Hz), 3.54 (4H, q, J=7.0Hz), 3.83 (1H, t, J=6.8Hz), 4.38-4.44 (1H, m), 5.36 (1H, dd, J=6.8, 8.9Hz), 6.93 (1H, d, J=7.5Hz), 7.02 (1H, d, J=2.2Hz), 7.07 (1H, t, J=7.5Hz), 7.15 (1H, m), 7.18-7.26 (4H, m), 7.66 (1H, d, J=7.9Hz), 8.92 (1H, brs).

Example 29

3-{{(2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl}-1-

methyl-1H-indole-7-carboxylic acid diethylamide

A solution of 3-((2R)-2-((5R)-5-(3-chlorophenyl)-2-oxooxazolidin-3-yl)propyl)-1H-indole-7-carboxylic acid diethylamide (10 mg, 0.022 mmol), methyl iodide (21 μ L, 0.33 mmol), potassium carbonate (90 mg, 0.66 mmol) in acetone (2 mL) is refluxed for 24 hours. After cooling, the insoluble materials are removed by filtration, and the filtrate is concentrated under reduced pressure. The residue is purified by preparative TLC (silica gel, 0.5 mm, 20 x 20 cm, methanol/chloroform = 1/30) to give 3-((2R)-2-((5R)-5-(3-chlorophenyl)-2-oxooxazolidin-3-yl)propyl)-1-methyl-1H-indole-7-carboxylic acid diethylamide (3 mg). A solution of 3-((2R)-2-((5R)-5-(3-chlorophenyl)-2-oxooxazolidin-3-yl)propyl)-1-methyl-1H-indole-7-carboxylic acid diethylamide (3 mg, 0.0064 mmol) and potassium hydroxide (0.6 g) in water (1 mL) and ethanol (1 mL) is stirred at 70°C for 7 hours. After cooling, ethanol is removed by distillation, and water is added thereto. The mixture is extracted with ethyl acetate, and the organic layer is washed with a saturated aqueous sodium hydrogen carbonate solution, dried over sodium sulfate, and concentrated. The residue is purified by silica gel column chromatography (60,2g, a saturated ammonia solution in chloroform) to give 3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethyl-amino]propyl)-1-methyl-1H-indole-7-carboxylic acid diethylamide (2 mg, yield: 71 %).

$^1\text{H-NMR}$ (CDCl_3 , 40°C) δ : 1.05 (3H, t, $J=7.1\text{Hz}$), 1.12 (3H, d, $J=5.5\text{Hz}$), 1.31 (3H, t, $J=7.1\text{Hz}$), 2.63 (1H, m), 2.79-2.87 (3H, m), 3.00 (1H, m), 3.23 (2H, q, $J=7.1\text{Hz}$), 3.55 (1H, m), 3.79 (1H, m), 4.50 (1H, dd, $J=3.5, 8.9\text{Hz}$), 6.82 (1H, s), 7.03 (1H, dd, $J=1.4, 7.2\text{Hz}$), 7.07 (1H, t, $J=7.6\text{Hz}$), 7.16-7.23 (3H, m), 7.35 (1H, s), 7.56 (1H, dd, $J=1.4, 7.6\text{Hz}$).

Reference Example 18

N-((7-Benzyloxy)-1H-indol-3-yl)methyl)-N,N-dimethylamine

To a solution of a 40 % aqueous dimethylamine solution (7.88 g, 69.9 mmol) and a 37 % aqueous formaldehyde solution (5.92 g, 72.9mmol) in acetic acid (70 mL) is added 7-(benzyloxy)-1H-indole (14.2 g, 63.6 mmol) at 0°C under nitrogen atmosphere, and the mixture is stirred at room temperature for 3.5 hours. Water is added to the reaction solution, and the mixture is washed with diethyl ether. The pH value of the aqueous layer is adjusted to pH 12 with a 3N aqueous sodium hydroxide solution, and the mixture is extracted with chloroform. The organic layer is washed with a saturated brine, and dried over anhydrous potassium carbonate. The solvent is evaporated, and the obtained crude product is dissolved in ethyl acetate (100 mL), crystallized from n-hexane (100 mL), and collected by filtration to give N-((7-benzyloxy)-1H-indol-3-yl)methyl)-N,N-dimethylamine (14.3 g, 50.9 mmol, yield: 80 %).

¹H-NMR (CDCl₃) δ: 8.35 (1H, brs), 7.47-7.49 (2H, m), 7.35-7.43 (3H, m), 7.32 (1H, d, J=8.0Hz), 7.10 (1H, d, J=2.3Hz), 7.03 (1H, dd, J=7.9, 7.7Hz), 6.73 (1H, d, J=7.7Hz), 5.20 (2H, s), 3.61 (2H, s), 2.27 (6H, s).

Reference Example 19

(7-(Benzyloxy)-1H-indol-3-yl)acetonitrile

To a solution of N-((7-benzyloxy)-1H-indol-3-yl)methyl)-N,N-dimethylamine (14.2 g, 50.6 mmol) in N,N-dimethylformamide (150 mL) is added a solution of potassium cyanide (13.2 g, 202.7 mmol) in water (25 ml) under nitrogen atmosphere, and after cooling thereto is added dropwise methyl iodide (34.5 g, 243.1 mmol), and the mixture is stirred

at room temperature for 14 hours. The reaction solution is poured into water, and the mixture is extracted with ethyl acetate. The organic layer is washed successively with water and a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated, and the obtained crude product is purified by silica gel column chromatography (n-hexane:ethyl acetate = 2:1) to give (7-(benzyloxy)-1H-indol-3-yl)acetonitrile (12.0 g, 45.7 mmol, yield: 90 %).

¹H-NMR (CDCl₃) δ: 8.42 (1H, brs), 7.46-7.48 (2H, m), 7.34-7.43 (3H, m), 7.19-7.21 (2H, m), 7.08 (1H, t, J=7.9Hz), 7.68 (1H, d, J=7.7Hz), 5.21 (2H, s), 3.83 (2H, d, J=0.84Hz).

Reference Example 20

(7-(Benzyloxy)-1H-indol-3-yl)acetic acid

Under nitrogen atmosphere, to a suspension of (7-(benzyloxy)-1H-indol-3-yl)acetonitrile (13.2 g, 50.3 mmol) in methanol (400 mL) is added 10N aqueous sodium hydroxide solution (130 mL), and the mixture is refluxed for 5 hours. The reaction solution is cooled to room temperature, and methanol alone is distilled off. The pH value of the resultant is adjusted to pH 1 with conc. hydrochloric acid while the mixture is cooled with ice. The precipitates are collected by filtration, and dissolved in chloroform, and dried over anhydrous sodium sulfate. The solvent is evaporated to give (7-(benzyloxy)-1H-indol-3-yl)acetic acid (12.2 g, 43.4 mmol, yield: 86 %).

¹H-NMR (DMSO-d₆) δ: 12.09 (1H, brs), 11.04 (1H, s), 7.55-7.57 (2H, m), 7.39-7.42 (2H, m), 7.31-7.35 (1H, m), 7.14 (1H, d, J=2.4Hz), 7.09 (1H, d, J=7.9Hz), 6.88 (1H, dd, J=7.9, 7.6Hz), 6.73 (1H, d, J=7.6Hz), 5.26 (2H, s), 3.61 (2H, s).

Reference Example 21

2-(7-(Benzyloxy)-1H-indol-3-yl)ethanol

To a solution of (7-(benzyloxy)-1H-indol-3-yl)acetic acid (7.21 g, 27.4 mmol) in tetrahydrofuran (150 mL) is added a 1M solution of borane · tetrahydrofuran complex in tetrahydrofuran (55 mL, 55 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for 17 hours. To the reaction solution is added methanol (100 mL), and the mixture is stirred at room temperature for one hour. The solvent is evaporated, and the resulting crude product is purified by silica gel column chromatography (n-hexane:ethyl acetate = 2:1 → 1:1) to give 2-(7-(benzyloxy)-1H-indol-3-yl)ethanol (6.37 g, 23.8 mmol, yield: 87 %).

$^1\text{H-NMR}$ (CDCl_3) δ : 8.31 (1H, s), 7.47-7.49 (2H, m), 7.34-7.43 (3H, m), 7.24 (1H, d, $J=8.1\text{Hz}$), 7.06 (1H, d, $J=3.2\text{Hz}$), 7.03 (1H, dd, $J=8.1, 7.7\text{Hz}$), 6.74 (1H, d, $J=7.7\text{Hz}$), 5.21 (2H, s), 3.90 (2H, td, $J=6.3, 5.3\text{Hz}$), 3.03 (2H, t, $J=6.3\text{Hz}$), 1.48 (1H, t, $J=5.3\text{Hz}$).

Reference Example 22

3-(2-Hydroxyethyl)-1H-indol-7-ol

To a solution of 2-(7-(benzyloxy)-1H-indol-3-yl)ethanol (6.31 g, 23.6 mmol) in ethanol (130 mL) are added ammonium formate (6.3 g, 99.9 mmol) and 10 % palladium on carbon (50 % wet, 555 mg), and the mixture is refluxed for one hour. The reaction solution is cooled to room temperature, and filtered through celite. The solvent is evaporated, and the obtained crude product is purified by silica gel column chromatography (n-hexane:ethyl acetate = 1:1) to give 3-(2-hydroxyethyl)-1H-indol-7-ol (3.19 g, 18.0 mmol, yield: 76 %).

$^1\text{H-NMR}$ (DMSO-d_6) δ : 10.54 (1H, s), 9.39 (1H, s), 7.00 (1H, d, $J=2.3\text{Hz}$), 6.94 (1H, d, $J=7.9\text{Hz}$), 6.75 (1H, t, $J=7.9\text{Hz}$), 6.46 (1H, d,

J=7.9Hz), 4.57 (1H, t, J=5.3Hz), 3.61 (2H, td, J=7.4, 5.3Hz), 2.79 (2H, t, J=7.4Hz).

Reference Example 23

Ethyl (((3-(2-hydroxyethyl)-1H-indol-7-yl)oxy)acetate

5 To a solution of 3-(2-hydroxyethyl)-1H-indol-7-ol (124 mg, 0.70 mmol) in acetone (5 mL) are added potassium carbonate (106 mg, 0.77 mmol), potassium iodide (11 mg, 0.066 mmol) and ethyl bromoacetate (172 μ L, 1.56 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for 13 hours. The insoluble materials are removed by filtration, and the solvent is removed by distillation. The resultant residue is purified by silica gel column chromatography (n-hexane:ethyl acetate = 5:1 \rightarrow 3:1 \rightarrow 1:2) to give ethyl (((3-(2-hydroxyethyl)-1H-indol-7-yl)oxy)acetate (122 mg, 0.46 mmol, yield: 66 %).

15 $^1\text{H-NMR}$ (CDCl_3) δ : 8.88 (1H, brs), 7.29 (1H, d, J=8.0Hz), 7.09 (1H, d, J=2.3Hz), 7.01 (1H, dd, J=8.0, 7.8Hz), 6.60 (1H, d, J=7.8Hz), 4.75 (2H, s), 4.28 (2H, q, J=7.1Hz), 3.90 (2H, td, J=6.3, 5.6Hz), 3.02 (2H, t, J=6.3Hz), 1.57 (1H, t, J=5.6Hz), 1.31 (3H, t, J=7.1Hz).

Reference Example 24

Ethyl (((3-(2-oxoethyl)-1H-indol-7-yl)oxy)acetate

20 To a solution of ethyl (((3-(2-hydroxyethyl)-1H-indol-7-yl)oxy)-acetate (1.2 g, 4.56 mmol) in dimethyl sulfide (56 mL) are added triethylamine (1.9 mL, 13.7 mmol) and sulfur trioxide \cdot pyridine complex (2.18 g, 13.7 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for one hour. The reaction solution is poured into water, and the mixture is extracted with ethyl acetate. The organic layer is washed successively with 1N aqueous hydrochloric acid solution, a saturated aqueous sodium hydrogen carbonate solution, and

a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated, and the obtained crude product is purified by silica gel column chromatography (n-hexane:ethyl acetate = 4:1) to give ethyl (((3-(2-oxoethyl)-1H-indol-7-yl)oxy)acetate (825 mg, 3.16 mmol, yield: 69 %).

¹H-NMR (CDCl₃) δ: 9.75 (1H, t, J=2.5Hz), 9.10 (1H, brs), 7.20 (1H, d, J=8.0Hz), 7.16 (1H, d, J=2.4Hz), 7.03 (1H, dd, J=8.0, 7.6Hz), 6.62 (1H, d, J=7.6Hz), 4.75 (2H, s), 4.28 (2H, q, J=7.2Hz), 3.78 (1H, dd, J=2.5, 0.7Hz), 1.30 (3H, t, J=7.2Hz).

Example 30

Ethyl (((3-(2-(((1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-methylethyl)-amino)ethyl)-1H-indol-7-yl)oxy)acetate

To a solution of ethyl (((3-(2-oxoethyl)-1H-indol-7-yl)oxy)acetate (103 mg, 0.394 mmol) in methylene chloride (10 mL) are added 4-((2S)-2-amino-(1R)-1-hydroxypropyl)phenol (J. Med. Chem., vol. 20, no. 7, p. 978 (1977)) (72 mg, 0.431 mmol), sodium triacetoxy borohydride (126 mg, 0.595 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for 5 hours. The reaction solution is poured into water, and the mixture is extracted with chloroform. The organic layer is washed with a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated, and the obtained crude product is purified by silica gel column chromatography (chloroform:methanol = 50:1 → 30:1 → 10:1) to give ethyl (((3-(2-(((1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-methylethyl)amino)ethyl)-1H-indol-7-yl)oxy)acetate (119 mg, 0.288 mmol, yield: 73 %).

¹H-NMR (CDCl₃) δ: 8.62 (1H, brs), 7.20 (1H, d, J=8.0Hz), 6.96 (1H, dd, J=8.0, 7.7Hz), 6.85 (2H, d, J=8.5Hz), 6.72 (1H, d, J=1.9Hz),

6.50 (1H, d, J=7.7Hz), 6.45 (2H, d, J=8.5Hz), 4.80 (1H, d, J=3.5Hz), 4.32-4.38 (3H, m), 3.01-3.50 (1H, m), 2.90-2.96 (1H, m), 2.75-2.85 (2H, m), 2.68 (1H, dq, J=7.0, 6.3Hz), 1.36 (1H, t, J=7.1Hz), 1.09 (3H, d, J=6.3Hz).

5 Example 31

((3-(2-(((1S,2R)-2-Hydroxy-2-(4-hydroxyphenyl)-1-methylethyl)amino)ethyl)-1H-indol-7-yl)oxy)acetic acid

To a solution of ethyl ((3-(2-(((1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-methylethyl)amino)ethyl)-1H-indol-7-yl)oxy)acetate (115 mg, 0.279 mmol) in tetrahydrofuran (5 mL) and water (0.5 mL) is added lithium hydroxide (10 mg, 0.418 mmol), and the mixture is stirred at room temperature for 11 hours. The pH value of the mixture is adjusted to pH 7 with 1N aqueous hydrochloric acid solution, and the solvent is distilled off. The obtained crude product is purified by octadecylsilyl (ODS) column chromatography (water → 10 % aqueous methanol) to give ((3-(2-(((1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-methylethyl)amino)ethyl)-1H-indol-7-yl)oxy)acetic acid (108 mg, 0.279 mmol, yield: 100 %).

¹H-NMR (CD₃OD) δ: 7.15 (1H, d, J=7.9Hz), 7.08 (1H, s), 7.03 (2H, d, J=8.5Hz), 6.93 (1H, dd, J=7.9, 7.6Hz), 6.70 (2H, d, J=8.5Hz), 6.57 (1H, d, J=7.6Hz), 4.83 (1H, d, J=3.7Hz), 4.47 (2H, s), 3.20-3.37 (3H, m), 3.11 (2H, t, J=7.4Hz), 0.90 (3H, d, J=6.8Hz).

Reference Example 25

3-((Dimethylamino)methyl)-N,N-diethyl-1H-indole-7-carboxamide

The title compound is obtained from 1H-indole-7-carboxylic acid diethylamide in a similar manner to Reference Example 18.

¹H-NMR (CDCl₃) δ: 8.96 (1H, brs), 7.77 (1H, d, J=7.9Hz), 7.22

(1H, dd, J=7.3, 0.9Hz), 7.17 (1H, d, J=2.3Hz), 7.08 (1H, dd, J=7.9, 7.3Hz), 3.63 (2H, s), 3.55 (4H, q, J=7.1Hz), 2.27 (6H, s), 1.27 (6H, t, J=7.1Hz).

Reference Example 26

5 3-(Cyanomethyl)-N,N-diethyl-1H-indole-7-carboxamide

The title compound is obtained from 3-((dimethylamino)methyl)-N,N-diethyl-1H-indole-7-carboxamide in a similar manner to Reference Example 19.

10 ¹H-NMR (CDCl₃) δ: 9.23 (1H, brs), 7.65 (1H, d, J=7.9Hz), 7.28 (1H, dd, J=7.3, 0.8Hz), 7.21 (1H, m), 7.16 (1H, dd, J=7.9, 7.3Hz), 3.84 (2H, d, J=0.9Hz), 3.55 (4H, q, J=7.1Hz), 1.27 (6H, t, J=7.1Hz).

Reference Example 27

(7-((Diethylamino)carbonyl)-1H-indol-3-yl)acetic acid

15 The title compound is obtained from 3-(cyanomethyl)-N,N-diethyl-1H-indole-7-carboxamide in a similar manner to Reference Example 20.

20 ¹H-NMR (CDCl₃) δ: 9.29 (1H, brs), 7.67 (1H, d, J=7.9Hz), 7.19 (1H, dd, J=7.3, 0.8Hz), 7.09 (1H, dd, J=7.9, 7.3Hz), 7.01 (1H, d, J=2.3Hz), 3.75 (2H, d, J=0.4Hz), 3.51 (4H, q, J=7.0Hz), 1.25 (6H, t, J=7.0Hz).

Reference Example 28

Methyl (7-((diethylamino)carbonyl)-1H-indol-3-yl)acetate

25 To a solution of (7-((diethylamino)carbonyl)-1H-indol-3-yl)acetic acid (2.38 g, 8.68 mmol) in methanol (10 mL) is added a 10 % hydrogen chloride solution in methanol (30 mL) under nitrogen atmosphere, and the mixture is refluxed for 30 minutes. The reaction solution is cooled to room temperature, and the solvent is distilled off. To the residue is

added chloroform, and the mixture is washed with a saturated aqueous sodium hydrogen carbonate solution and a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is removed by distillation, and the resultant crude product is purified by silica gel column chromatography (n-hexane:ethyl acetate = 1:1 → ethyl acetate) to give methyl (7-((diethylamino)carbonyl)-1H-indol-3-yl)acetate (1.93 g, yield: 77 %).

¹H-NMR (CDCl₃) δ: 8.97 (1H, brs), 7.67 (1H, d, J=7.9Hz), 7.22-7.25 (2H, m), 7.11 (1H, dd, J=7.9, 7.5Hz), 3.79 (2H, s), 3.70 (3H, s), 3.55 (4H, q, J=7.0Hz), 1.26 (6H, t, J=7.0Hz).

Reference Example 29

N,N-Diethyl-3-(2-hydroxyethyl)-1H-indole-7-carboxamide

To a solution of methyl (7-((diethylamino)carbonyl)-1H-indol-3-yl)acetate (1.87 g, 6.48 mmol) in tetrahydrofuran (30 mL) are added sodium borohydride (1.35 g, 35.7 mmol) and methanol (5 mL) under nitrogen atmosphere, and the mixture is stirred at room temperature for 24 hours, and then refluxed for 4 hours. The mixture is cooled to room temperature, and methanol is added to the reaction solution. The solvent is evaporated, and the residue is separated into water and chloroform. The organic layer is washed with a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated, and the obtained crude product is purified by silica gel column chromatography (n-hexane:ethyl acetate = 1:1 → ethyl acetate → ethyl acetate:methanol = 5:1) to give N,N-diethyl-3-(2-hydroxyethyl)-1H-indole-7-carboxamide (1.56 g, 5.99 mmol, yield: 92 %).

¹H-NMR (CDCl₃) δ: 8.94 (1H, brs), 7.67 (1H, d, J=7.9Hz), 7.23 (1H, dd, J=7.3, 0.8Hz), 7.13 (1H, d, J=2.2Hz), 7.10 (1H, dd, J=7.9,

7.3Hz), 3.90 (2H, td, J=6.0, 6.1Hz), 3.55 (4H, q, J=7.1Hz), 3.04 (2H, t, J=6.0Hz), 1.49 (1H, t, J=6.1Hz), 1.27 (6H, t, J=7.1Hz).

Reference Example 30

N,N-Diethyl-3-(2-oxoethyl)-1H-indole-7-carboxamide

5 The title compound is obtained from N,N-diethyl-3-(2-hydroxyethyl)-1H-indole-7-carboxamide in a similar manner to Reference Example 24.

10 ¹H-NMR (CDCl₃) δ: 9.75 (1H, t, J=2.5Hz), 9.10 (1H, brs), 7.59 (1H, d, J=7.9Hz), 7.26 (1H, dd, J=7.2, 0.7Hz), 7.21 (1H, d, J=2.3Hz), 7.12 (1H, dd, J=7.9, 7.2Hz), 3.82 (2H, dd, J=2.5, 0.7Hz), 3.56 (4H, q, J=7.1Hz), 1.27 (6H, t, J=7.1Hz).

Example 32

N,N-Diethyl-3-(2-(((1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-methylethyl)amino)ethyl)-1H-indole-7-carboxamide

15 The title compound is obtained from N,N-diethyl-3-(2-oxoethyl)-1H-indole-7-carboxamide in a similar manner to Example 30.

20 ¹H-NMR (CDCl₃) δ: 8.28 (1H, brs), 7.59 (1H, d, J=7.9Hz), 7.19 (1H, dd, J=7.3, 0.9Hz), 7.09 (1H, dd, J=7.9, 7.3Hz), 6.69 (2H, d, J=8.5Hz), 6.66 (1H, d, J=2.2Hz), 6.30 (2H, d, J=8.5Hz), 4.15 (1H, d, J=8.3Hz), 3.54 (4H, brs), 3.01-3.06 (1H, m), 2.92-2.97 (1H, m), 2.74-2.81 (1H, m), 2.64-2.70 (1H, m), 2.49-2.57 (1H, m), 1.26 (6H, brs), 1.24 (3H, d, J=6.1Hz).

Reference Example 31

1H-Indole-6-carboxylic acid diethylamide

25 The title compound is obtained from indole-6-carboxylic acid in a similar manner to Reference Example 14.

¹H-NMR (CDCl₃) δ: 1.20 (6H, brs), 3.3-3.5 (4H, brs), 6.54 (1H, s),

7.11 (1H, d, J=6.8Hz), 7.25 (1H, m), 7.47 (1H, s), 7.61 (1H, d, J=8.1Hz), 8.80 (1H, brs).

Reference Example 32

(R)-[2-(6-Diethylcarbamoyl-1H-indol-3-yl)-1-methyl-2-oxoethyl]carbamic acid 9H-fluoren-9-ylmethyl ester

The title compound is obtained from 1H-indole-6-carboxylic acid diethylamide in a similar manner to Reference Example 15.

¹H-NMR (CDCl₃) δ: 1.08 (3H, brs), 1.25 (3H, brs), 1.42 (3H, d, J=6.9Hz), 3.25 (2H, brs), 3.56 (2H, brs), 4.21 (1H, t, J=7.1Hz), 4.36 (2H, m), 4.96 (1H, dq, J=7.2, 7.2Hz), 6.06 (1H, d, J=7.6Hz), 7.21 (1H, m), 7.27 (2H, m), 7.37 (2H, dt, J=3.1, 7.4Hz), 7.59 (2H, dd, J=3.8, 7.3Hz), 7.67 (1H, d, J=3.0Hz), 7.74 (1H, d, J=7.5Hz), 8.31 (1H, d, J=8.2Hz).

Reference Example 33

(R)-3-(2-Aminopropyl)-1H-indole-6-carboxylic acid diethylamide

The title compound is obtained from (R)-[2-(6-diethylcarbamoyl-1H-indol-3-yl)-1-methyl-2-oxoethyl]carbamic acid 9H-fluoren-9-ylmethyl ester in a similar manner to Reference Example 16.

¹H-NMR (CDCl₃) δ: 1.14 (3H, d, J=6.3Hz), 1.20 (6H, brs), 2.60 (1H, d, J=8.3, 14.2Hz), 2.84 (1H, dd, J=4.7, 14.2Hz), 3.22 (1H, m), 3.3-3.5 (4H, brs), 7.00 (1H, s), 7.06 (1H, dd, J=1.3, 8.1Hz), 7.39 (1H, s), 7.54 (1H, d, J=8.1Hz).

Example 33

3-[(2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl]-1H-indole-6-carboxylic acid diethylamide · trifluoroacetate

The title compound is obtained from (R)-3-(2-aminopropyl)-1H-indole-6-carboxylic acid diethylamide in a similar manner to Example 9.

¹H-NMR (CD₃OD) δ: 1.14 (3H, brs), 1.26 (3H, brs), 1.31 (3H, d,

J=6.5Hz), 3.03 (1H, dd, J=9.3, 14.2Hz), 3.16-3.35 (5H, m), 3.56 (2H, brs), 3.66 (1H, m), 4.95 (1H, dd, J=3.2Hz, 10.0Hz), 7.07 (1H, dd, J=1.4, 8.2Hz), 7.3-7.4 (4H, m), 7.43 (1H, m), 7.47 (1H, m), 7.67 (1H, d, J=8.2Hz), 10.9 (1H, brs).

5 Reference Example 34

3-((2R)-2-[2-(4-Benzyloxy-3-methanesulfonylamino-phenylphenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid diethylamide

The title compound is obtained from (R)-3-(2-aminopropyl)-1H-indole-7-carboxylic acid diethylamide in a similar manner to Reference
10 Example 8.

¹H-NMR (CDCl₃) δ: 8.88 (1H, s), 7.62 (1H, d, J=7.8Hz), 7.31-7.43 (6H, m), 7.20 (1H, d, J=6.6Hz), 7.03-7.12 (3H, m), 6.92 (1H, d, J=8.5Hz), 5.07 (2H, s), 4.51 (1H, dd, J=8.5, 4.1Hz), 3.54 (4H, q, J=7.1Hz), 2.91-3.07 (1H, m), 2.85 (3H, s), 2.78-2.86 (3H, m), 2.71 (dd, J=11.8, 8.5Hz), 1.26 (6H, t, J=7.1Hz), 1.14 (3H, d, J=9.2Hz).

15 Example 34

3-((2R)-2-[(2R)-2-Hydroxy-2-(4-hydroxy-3-methanesulfonylamino-phenyl)ethylamino]propyl)-1H-indole-7-carboxylic acid diethylamide trifluoroacetate

20 To a solution of 3-((2R)-2-[2-(4-benzyloxy-3-methanesulfonylamino-phenylphenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid diethylamide (15.1 mg, 0.0255 mmol) in methanol (3 mL) is added 10 % palladium on carbon (50 % wet, 5 mg), and the mixture is stirred at room temperature under hydrogen atmosphere for
25 4 hours. The reaction solution is filtered through celite, and the filtrate is evaporated under reduced pressure to remove the solvent. The residue is purified by preparative TLC (a saturated ammonia solution in

chloroform:a saturated ammonia solution in methanol = 5:1) and preparative HPLC (trifluoroacetic acid-acetonitrile water) to give the desired 3-((2R)-2-((2R)-2-hydroxy-2-(4-hydroxy-3-methanesulfonylaminophenyl)ethylamino)propyl)-1H-indole-7-carboxylic acid diethylamide · trifluoroacetate (10.0 mg, 0.0162 mmol, yield: 64 %).

¹H-NMR (CD₃OD) δ: 10.7 (1H, s), 7.67-7.71 (1H, m), 7.40 (1H, d, J=2.1Hz), 7.26 (1H, s), 7.07-7.20 (3H, m), 6.90 (1H, d, J=8.3Hz), 4.80-4.86 (1H, m), 3.30-3.68 (5H, m), 3.18-3.28 (3H, m), 3.00-3.05 (1H, m), 2.92 (3H, s), 1.08-1.34 (9H, m).

Reference Example 35

tert-Butyl [2-(4-benzyloxy-3-methanesulfonylaminophenyl)-(2R)-2-hydroxyethyl]-[2-(7-diethylcarbamoyl-1H-indol-3-yl)-(1R)-1-methylethyl]carbamate

To a solution of 3-((2R)-2-[2-(4-benzyloxy-3-methanesulfonylaminophenylphenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid diethylamide (72.3 mg, 0.122 mmol) in tetrahydrofuran (3 mL) is added di-tert-butyl bicarbonate (40 mg, 0.183 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for 17 hours. The reaction solution is poured into water, and the mixture is extracted with chloroform. The organic layer is washed with a saturated brine, and dried over anhydrous sodium sulfate. The solvent is distilled off, and the obtained crude product is purified by preparative TLC (chloroform:methanol = 30:1) to give the desired tert-butyl [2-(4-benzyloxy-3-methanesulfonylaminophenyl)-(2R)-2-hydroxyethyl]-[2-(7-diethylcarbamoyl-1H-indol-3-yl)-(1R)-1-methylethyl]carbamate (79 mg, 0.114 mmol, yield: 93 %).

¹H-NMR (CDCl₃) δ: 8.88 (1H, s), 7.63 (1H, d, J=7.3Hz), 7.49 (1H,

brs), 7.35-7.43 (6H, m), 7.20 (1H, d, J=7.2Hz), 7.09 (1H, dd, J=8.8, 7.7Hz), 6.95-7.00 (2H, m), 6.82 (1H, s), 5.10 (2H, s), 4.71 (1H, d, J=8.3Hz), 4.28-4.34 (1H, m), 3.50-3.62 (5H, m), 3.11-3.15 (1H, m), 2.90 (3H, s), 2.79 (2H, m), 1.22-1.26 (18H, m).

5 Reference Example 36

tert-Butyl [2-(7-diethylcarbamoyl-1H-indol-3-yl)-(1R)-1-methylethyl]-[(2R)-2-hydroxy-2-(4-methanesulfonyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)ethyl]carbamate

To a solution of tert-butyl [2-(4-benzyloxy-3-methanesulfonyl-aminophenyl)-(2R)-2-hydroxyethyl]-[2-(7-diethylcarbamoyl-1H-indol-3-yl)-(1R)-1-methylethyl]carbamate (70.0 mg, 0.101 mmol) in methanol (3 mL) is added a 10 % palladium on carbon (50 % wet, 35 mg), and the mixture is stirred at room temperature under hydrogen atmosphere for one hour. The reaction solution is filtered through celite, and the solvent is evaporated to remove the solvent to give crude tert-butyl [2-(7-diethylcarbamoyl-1H-indol-3-yl)-(1R)-1-methylethyl]-[(2R)-2-hydroxy-2-(4-hydroxy-3-methanesulfonylaminophenyl)ethyl]carbamate (54.8 mg). To a solution of crude tert-butyl [2-(7-diethylcarbamoyl-1H-indol-3-yl)-(1R)-1-methylethyl]-[(2R)-2-hydroxy-2-(4-hydroxy-3-methanesulfonylaminophenyl)ethyl]carbamate (54.8 mg) in N,N-dimethylformamide (3 mL) are added 1,2-dibromoethane (18 μ L, 0.209 mmol) and potassium carbonate (27.9 mg, 0.202 mmol), and the mixture is stirred at 80°C for 2.5 hours. The reaction solution is cooled to room temperature, and the mixture is poured into a saturated aqueous sodium hydrogen carbonate solution. The mixture is extracted with ethyl acetate, and the organic layer is washed with a saturated brine, and dried over anhydrous sodium sulfate. The solvent is distilled off, and the

resulting crude product is purified by preparative TLC (ethyl acetate) to give the desired tert-butyl [2-(7-diethylcarbamoyl-1H-indol-3-yl)-(1R)-1-methylethyl]-[(2R)-2-hydroxy-2-(4-methanesulfonyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)ethyl]carbamate (54.2 mg, 0.0862 mmol, yield: 85 %).

¹H-NMR (CDCl₃) δ: 8.85 (1H, s), 7.62-7.66 (2H, m), 7.21 (1H, d, J=7.0 Hz), 7.06-7.16 (2H, m), 6.89-6.95 (2H, m), 4.70 (1H, d, J=8.3 Hz), 4.23-4.35 (3H, m), 3.81-3.95 (2H, m), 3.48-3.65 (5H, m), 3.11-3.15 (1H, m), 2.96 (3H, s), 2.80 (3H, m), 1.22-1.27 (18H, m).

Example 35

3-[(2R)-2-[(2R)-2-Hydroxy-2-(4-methanesulfonyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)ethylamino]propyl]-1H-indole-7-carboxylic acid diethylamide · trifluoroacetate

To a solution of tert-butyl [2-(7-diethylcarbamoyl-1H-indol-3-yl)-(1R)-1-methylethyl]-[(2R)-2-hydroxy-2-(4-methanesulfonyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)ethyl]carbamate (48.4 mg, 0.077 mmol) in methylene chloride (3 mL) is added trifluoroacetic acid (250 μL) under nitrogen atmosphere, and the mixture is stirred at room temperature for 18 hours, and to the mixture is added trifluoroacetic acid (1 mL). The mixture is stirred at room temperature for 3 hours, and thereto are added trifluoroacetic acid (1 mL) and thioanisole (50 μL). The mixture is stirred at room temperature for one hour. The pH value of the reaction solution is adjusted to pH 8 with a saturated aqueous sodium hydrogen carbonate solution, and the solvent is distilled off. The residue is dissolved in chloroform:methanol (2:1, 75 mL), and the insoluble materials are removed by filtration. The solvent is removed by distillation, and the crude product is purified by preparative TLC (a

saturated ammonia solution in chloroform) and preparative HPLC (trifluoroacetic acid- acetonitrile-water) to give the desired 3-((2R)-2-[(2R)-2-hydroxy-2-(4-methanesulfonyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)ethylamino]propyl)-1H-indole-7-carboxylic acid diethylamide trifluoroacetate (6.6 mg, 0.0103 mmol, yield: 13 %).

¹H-NMR (CD₃OD) δ: 10.6 (1H, s), 7.67 (1H, d, J=7.7Hz), 7.40 (1H, d, J=2.0Hz), 7.08-7.16 (3H, m), 6.92 (1H, d, J=8.5Hz), 6.87 (1H, dd, J=8.5, 2.0Hz), 4.78 (1H, dd, J=6.7, 4.1Hz), 4.25-4.32 (2H, m), 3.63-3.92 (5H, m), 3.32-3.59 (4H, m), 3.39 (1H, dd, J=16.7, 3Hz), 3.07 (1H, dd, J=16.7, 11.4Hz), 2.92 (3H, s), 1.51 (3H, d, J=6.6Hz), 0.97-1.30 (6H, m).

Reference Example 37

Methyl (R)-3-(2-aminopropyl)-1H-indole-7-carboxylate

(R)-3-(2-Aminopropyl)-1H-indole-7-carboxylic acid diethylamide (1.80 g, 6.59 mmol) is dissolved in a mixture of 6N hydrochloric acid (10 mL) and 1,4-dioxane (10 mL), and the mixture is stirred at 150°C for 14 hours in a sealed vessel. After cooling, the mixture is concentrated, and the residue is dissolved in 10 % HCl/methanol. The mixture is refluxed for 12 hours, cooled and concentrated. To the residue is added ethyl acetate, and the mixture is washed with a saturated aqueous sodium hydrogen carbonate solution and a saturated brine, dried over sodium sulfate, and concentrated to give the desired methyl (R)-3-(2-aminopropyl)-1H-indole-7-carboxylate (1.53 g, yield: 100 %).

¹H-NMR (CDCl₃) δ: 1.18 (3H, d, J=6.3Hz), 2.71 (1H, dd, J=8.1, 14.4Hz), 2.90 (1H, ddd, J=0.7, 5.2, 14.3Hz), 3.99 (3H, s), 7.15 (1H, dd, J=7.6, 7.6Hz), 7.17 (1H, d, J=1.2Hz), 7.84 (1H, J=7.8Hz), 7.89 (1H, dd, J=0.9, 7.5Hz), 9.69 (1H, brs).

Reference Example 38

Methyl 3-[(2R)-2-(2-pyridin-3-yl-(2R)-2-(triethylsilyloxy)ethylamino)-propyl]-1H-indole-7-carboxylate

A solution of methyl (R)-(pyridin-3-yl)oxirane (1.17 g, 9.7 mmol) and methyl (R)-3-(2-aminopropyl)-1H-indole-7-carboxylate (1.50 g, 6.5 mmol) in methanol (10 mL) is stirred at 100°C for 2 hours under sealed conditions, and the mixture is cooled and concentrated. The residue is dissolved in N,N-dimethylformamide (20 mL), and thereto are added imidazole (1.70 g, 25 mmol) and 4-dimethylaminopyridine (61 mg, 0.5 mmol) and triethylsilyl chloride (3.25 mL, 19.4 mmol). The mixture is stirred at room temperature for 3 hours, and thereto are added ethyl acetate. The mixture is washed with a saturated aqueous sodium hydrogen carbonate solution, water, and a saturated brine, dried over sodium sulfate, and concentrated. The residue is purified by silica gel column chromatography (methanol/chloroform = 1/50) to give the desired methyl 3-[(2R)-2-(2-pyridin-3-yl-(2R)-2-(triethylsilyloxy)ethylamino)propyl]-1H-indole-7-carboxylate (0.74 g, yield: 24 %).

¹H-NMR (CDCl₃) δ: 0.43 (6H, dq, J=2.5, 8.3Hz), 0.80 (9H, t, J=8.0Hz), 1.09 (3H, d, J=6.2Hz), 2.69-2.77 (2H, m), 2.84 (1H, dd, J=6.7, 14.4Hz), 2.91-2.99 (2H, m), 3.99 (3H, s), 4.76 (1H, dd, J=5.0, 7.1Hz), 7.05 (1H, d, J=2.1Hz), 7.11-7.15 (2H, m), 7.53-7.56 (1H, m), 7.78 (1H, d, J=7.8Hz), 7.88 (1H, dd, J=1.1, 7.6Hz), 8.45 (1H, dd, J=1.7, 4.8Hz), 8.52 (1H, d, J=2.1Hz), 9.60 (1H, brs).

Example 36

Methyl 3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indole-7-carboxylate

methyl 3-[(2R)-2-(2-pyridin-3-yl-(2R)-2-(triethylsilyloxy)ethylamino)propyl]-1H-indole-7-carboxylate (0.72 g, 1.54 mmol) is dissolved

in trifluoroacetic acid (20 mL), and the mixture is stirred at room temperature for one day. After concentrated, to the residue is added ethyl acetate, and the mixture is washed with a saturated aqueous sodium hydrogen carbonate solution and a saturated brine, dried over sodium sulfate, and concentrated. The residue is purified by silica gel column chromatography (methanol/chloroform = 1/100 → 1/3) to give the desired methyl 3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indole-7-carboxylate (440 mg, yield: 81 %).

¹H-NMR (CDCl₃) δ: 1.16 (3H, d, J=6.2Hz), 2.70 (1H, dd, J=9.3, 12.2Hz), 2.88-2.95 (3H, m), 3.06-3.11 (1H, m), 3.99 (3H, s), 4.61 (1H, dd, J=3.6, 9.2), 7.14-7.18 (2H, m), 7.22-7.26 (1H, m), 7.65-7.68 (1H, m), 7.81 (1H, d, J=7.8Hz), 7.90 (1H, dd, J=0.8, 7.5Hz), 8.49 (1H, dd, J=1.7, 4.8Hz), 8.54 (1H, d, J=1.2Hz), 9.71 (1H, brs).

Example 37

3-[(2R)-2-((2R)-2-Hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indole-7-carboxylic acid · 2 trifluoroacetate

A solution of methyl 3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indole-7-carboxylate (440 mg, 1.24 mmol) and potassium hydroxide (1.14 g, 20 mmol) in water (2 mL)/methanol (10 mL) is stirred at room temperature for 19 hours. After concentrated, the resultant is purified by reversed phase column chromatography (octadecylsilyl, trifluoroacetic acid/water/methanol) to give the desired 3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indole-7-carboxylic acid · 2 trifluoroacetate (560 mg, yield: 80 %).

¹H-NMR (CD₃OD) δ: 1.32 (3H, d, J=6.5Hz), 3.05 (1H, dd, J=9.5, 14.2Hz), 3.30-3.41 (3H, m), 3.65-3.73 (1H, m), 5.13 (1H, dd, J=3.3, 9.9Hz), 7.16 (1H, dd, J=7.7, 7.7Hz), 7.33 (1H, s), 7.67 (1H, dd, J=5.2,

8.0Hz), 7.88 (1H, d, J=7.4Hz), 7.89 (1H, d, J=7.8Hz), 8.13-8.17 (1H, m), 8.62 (1H, dd, J=1.2, 5.1Hz), 8.72 (1H, d, J=1.8Hz), 10.67 (1H, brs).

Examples 38 to 41

The compounds of the following Examples are obtained from 3-
5 [(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indole-7-carboxylic acid · 2 trifluoroacetate in a similar manner to Example 11.

Example 38

(3-((2R)-2-[(2R)-2-Hydroxy 2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yl)-piperidin-1-ylmethanone · 2 trifluoroacetate

10 ¹H-NMR (CD₃OD) δ: 1.32 (3H, d, J=6.6Hz), 1.53-1.71 (6H, m), 3.04 (1H, dd, J=9.3, 14.2Hz), 3.32-3.47 (4H, m), 3.44 (1H, dd, J=3.2, 12.8Hz), 3.67-3.97 (3H, m), 5.26 (1H, dd, J=3.0, 9.9Hz), 7.12-7.19 (2H, m), 7.28 (1H, s), 7.72 (1H, dd, J=1.8, 7.2Hz), 8.03 (1H, dd, J=7.7, 8.1Hz), 8.57 (1H, d, J=8.1Hz), 8.81 (1H, d, J=5.5Hz), 8.88 (1H, d, J=1.3Hz).

15 Example 39

(3-((2R)-2-[(2R)-2-Hydroxy-2-pyridin-3-ylethylamino]propyl)-1H-indol-7-yl)-morpholin-4-ylmethanone · 2 trifluoroacetate

20 ¹H-NMR (CD₃OD) δ: 1.33 (3H, d, J=6.5Hz), 2.85-3.12 (2H, m), 3.32-3.48 (2H, m), 3.68 (9H, m), 5.17 (1H, dd, J=3.1, 10.0Hz), 7.10-7.21 (2H, m), 7.30 (1H, s), 7.73 (1H, dd, J=1.1, 7.8Hz), 7.84 (1H, dd, J=5.2, 7.8Hz), 8.34 (1H, d, J=8.1Hz), 8.71 (1H, d, J=4.8Hz), 8.79 (1H, s), 10.78 (1H, brs).

Example 40

25 3-((2R)-2-[(2R)-2-Hydroxy-2-pyridin-3-ylethylamino]propyl)-1H-indole-7-carboxylic acid dimethylamide · 2 trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.33 (3H, d, J=6.6Hz), 2.87-3.33 (9H, m), 3.39 (1H, dd, J=3.3, 12.9Hz), 3.68-3.72 (1H, m), 5.16 (1H, dd, J=3.2,

9.9Hz), 7.14 (1H, dd, J=7.4, 7.7Hz), 7.19 (1H, dd, J=0.9, 7.2Hz), 7.29 (1H, s), 7.72 (1H, dd, J=1.1, 7.8Hz), 7.83 (1H, dd, J=5.4, 8.1Hz), 8.32 (1H, d, J=8.1Hz), 8.70 (1H, d, J=4.8Hz), 8.77 (1H, s), 10.72 (1H, brs).

Example 41

5 3-((2R)-2-[(2R)-2-Hydroxy-2-pyridin-3-ylethylamino]propyl)-1H-indole-7-carboxylic acid ethylamide · 2 trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.26 (3H, t, J=7.2Hz), 1.33 (3H, d, J=6.6Hz), 2.85-3.12 (2H, m), 3.31-3.49 (2H, m), 3.46 (2H, q, J=7.3Hz), 3.68-3.72 (1H, m), 5.21 (1H, dd, J=3.1, 10.0Hz), 7.13 (1H, dd, J=7.7, 7.7Hz), 7.32 (1H, s), 7.60 (1H, d, J=7.0Hz), 7.80 (1H, dd, J=0.8, 7.9Hz), 7.92 (1H, dd, J=5.4, 8.0Hz), 8.45 (1H, d, J=8.1Hz), 8.75 (1H, d, J=5.4Hz), 8.85 (1H, s).

Example 42

(1R)-1-(3-Chlorophenyl)-2-[2-(7-diethylaminomethyl-1H-indol-3-yl)-(1R)-1-(methyl)ethylamino]ethanol

15 A solution of 3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid diethylamide (30 mg, 0.07 mmol) in tetrahydrofuran (3 mL) is cooled to 0°C, and thereto is added LiAlH₄ (9 mg, 0.21 mmol), and the mixture is stirred at 0°C for 4 hours. To the mixture are added water (10 μL), a 15 % aqueous NaOH solution (10 μL) and water (30 μL), and the insoluble materials are removed by filtration, and the filtrate is concentrated. The residue is purified by preparative reversed phase HPLC (trifluoroacetic acid/water/acetonitrile) to give the desired (1R)-1-(3-chlorophenyl)-2-[2-(7-diethylaminomethyl-1H-indol-3-yl)-(1R)-1-(methyl)ethylamino]ethanol (16 mg, yield: 36 %).

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.6Hz), 1.33 (3H, t, J=7.3Hz), 3.03 (1H, dd, J=9.6, 14.2Hz), 3.15-3.34 (3H, m), 3.26 (2H, q, J=7.3Hz),

3.65-3.69 (1H, m), 4.61 (2H, s), 4.98 (1H, dd, $J=3.1, 10.1\text{Hz}$), 7.20 (1H, dd, $J=7.5, 7.7\text{Hz}$), 7.30-7.39 (5H, m), 7.47 (1H, s), 7.76 (1H, dd, $J=0.7, 7.9$).

Example 43

{3-[(2R)-2-((2R)-2-Hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yloxy}acetic acid 1-(cyclohexyloxycarbonyloxy)ethyl ester

{3-[(2R)-2-((2R)-2-Hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yloxy}acetic acid (4 mg, 0.011 mmol) and potassium carbonate (3 mg, 0.022 mmol) are added to N,N-dimethylformamide (2 mL), and to the mixture is added at 0°C 1-iodoethylcyclohexyl carbonate (J.

Antibiot., vol. 40, no. 1, p. 81-90 (1987)) (7.8 mg, 0.026 mmol). The mixture is stirred at 0°C for 2 hours, and thereto is added a saturated brine. The mixture is extracted with ethyl acetate, and the organic layer is washed with a saturated brine, dried over sodium sulfate, and concentrated. The residue is purified by silica gel column chromatography (methanol/chloroform = 1/20 to 1/1) to give the desired {3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yloxy}-acetic acid 1-(cyclohexyloxycarbonyloxy)ethyl ester (1 mg, yield: 17 %).

¹H-NMR (CDCl₃) δ : 1.23 (3H, d, $J=6.2\text{Hz}$), 1.28-1.53 (6H, m), 1.56 (1H, d, $J=5.4\text{Hz}$), 1.73 (2H, m), 1.91 (2H, m), 2.77-2.82 (1H, m), 2.89-3.01 (3H, m), 3.19-3.26 (1H, m), 4.60-4.66 (1H, m), 4.77 (2H, s), 4.81 (1H, m), 6.58 (1H, d, $J=7.6\text{Hz}$), 6.87 (1H, q, $J=5.3\text{Hz}$), 7.00 (1H, dd, $J=7.8, 7.9\text{Hz}$), 7.09 (1H, s), 7.22-7.24 (2H, m), 7.68 (1H, d, $J=7.8\text{Hz}$), 8.50 (1H, d, $J=3.5\text{Hz}$), 8.55 (1H, s), 8.76 (1H, s).

Example 44

Ethyl {3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yloxy}acetate · 2 hydrochloride

{3-[(2R)-2-((2R)-2-Hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yloxy}acetic acid (65 mg, 0.176 mmol) is dissolved in ethanol (5 mL), and thereto are added conc. sulfuric acid (0.1 mL) and Molecular Sieves 4A powder (trade name, 0.3 g), and the mixture is refluxed for 20 hours. After cooling, the insoluble materials are removed by filtration, and a saturated aqueous sodium hydrogen carbonate solution is added to the filtrate. The mixture is evaporated under reduced pressure to remove ethanol, and extracted with ethyl acetate. The organic layer is washed with a saturated brine, dried over sodium sulfate, and concentrated. The residue is purified by silica gel column chromatography (ethanol/chloroform = 1/5 → 1/1). The obtained purified product is dissolved in ethanol (3 mL), and thereto is added 1N HCl/diethyl ether (0.5 mL). The mixture is concentrated under reduced pressure to give the desired ethyl {3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-yloxy}acetate · 2 hydrochloride (63 mg, yield: 76 %).

¹H-NMR (DMSO-d₆) δ: 1.17 (3H, d, J=6.5Hz), 1.22 (3H, t, J=7.1Hz), 2.85 (1H, dd, J=10.2, 13.9Hz), 3.25-3.39 (3H, m), 3.50 (1H, m), 4.18 (2H, q, J=7.1Hz), 4.91 (2H, s), 5.11-5.16 (1H, m), 6.54 (1H, d, J=7.7Hz), 6.89 (1H, dd, J=7.9, 7.9Hz), 7.17 (1H, d, J=2.4Hz), 7.24 (1H, d, J=7.9Hz), 7.72 (1H, brs), 8.17 (1H, brs), 8.70 (1H, d, J=4.5Hz), 8.78 (1H, s), 8.96 (1H, brs), 11.15 (1H, d, J=2.2Hz).

Example 45

3-[(2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl]-1H-indole-6-carboxylic acid diethylamide

To a solution of (R)-3-(2-aminopropyl)-1H-indole-6-carboxylic acid diethylamide (960 mg, 3.51 mmol) in acetonitrile (10 ml) is added

(R)-(+)-3-chlorostyrene oxide (1.09 g, 7.02 mmol), and the mixture is refluxed for 5 hours. The reaction solution is concentrated under reduced pressure, and the residue is purified by silica gel column chromatography (chloroform/methanol = 20/1 → (a saturated ammonia solution in chloroform)/methanol = 20/1) to give 3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid diethylamide (950 mg, 2.22 mmol, 63 %).

¹H-NMR (CDCl₃) δ: 1.12 (3H, d, J=6.2Hz), 1.20 (6H, br), 2.64 (1H, dd, J=9.0, 12.1Hz), 2.83 (2H, d, J=6.6Hz), 2.88 (1H, dd, J=3.7, 12.1Hz), 3.04 (1H, m), 3.30 (4H, br), 4.50 (1H, dd, J=3.5, 8.9Hz), 7.08 (1H, d, J=2.2Hz), 7.12 (1H, dd, J=2.2, 8.1Hz), 7.21 (3H, m), 7.34 (1H, s), 7.43 (1H, s), 7.57 (1H, d, J=8.1Hz), 8.16 (1H, br).

Example 46

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid

To a solution of 3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid diethylamide (800 mg, 1.87 mmol) in 1,4-dioxane (10 ml) is added 6N-HCl (10 ml), and the mixture is refluxed for 24 hours. After allowed to cool, the reaction solution is concentrated under reduced pressure, and the residue is purified by octadecylsilyl column chromatography (trifluoroacetic acid/methanol/water), and further crystallized from aqueous methanol (pH 7) to give 3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid (304 mg, 0.815 mmol, 44 %) as white crystals.

¹H-NMR (CD₃OD) δ: 1.28 (3H, d, J=6.5Hz), 2.99 (1H, dd, J=8.7, 14.2Hz), 3.18 (3H, m), 3.56 (1H, m), 4.92 (1H, m), 7.31 (4H, m), 7.43

(1H, s), 7.55 (1H, d, J=8.4Hz), 7.71 (1H, dd, J=1.2, 8.3Hz), 8.06 (1H, d, J=8.2Hz).

Example 47

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid · trifluoroacetate

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid (9.8 mg, 0.0263 mmol) is purified by preparative reversed phase HPLC (octadecylsilyl, trifluoroacetic acid/ acetonitrile/water) to give 3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid · trifluoroacetate (3.9 mg, 0.0080 mmol, 30 %).

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.5Hz), 3.03 (1H, dd, J=9.3, 14.2Hz), 3.18 (1H, dd, J=10.0, 12.5Hz), 3.20 (2H, m), 3.66 (1H, m), 4.95 (1H, dd, J=9.2, 10.0Hz), 7.31 (3H, m), 7.40 (1H, s), 7.45 (1H, s), 7.65 (1H, d, J=8.4Hz), 7.75 (1H, dd, J=1.4, 8.4Hz), 8.13 (1H, m).

Examples 48 to 59

The compounds of the following Examples are obtained from 3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid in a similar manner to Example 11.

Example 48

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid dibutylamide · trifluoroacetate

¹H-NMR (CD₃OD) δ: 0.74 (3H, brs), 1.01 (3H, brs), 1.10 (2H, brs), 1.30 (3H, d, J=6.5Hz), 1.42 (2H, brs), 1.54 (2H, brs), 1.68 (2H, brs), 3.02 (1H, dd, J=9.3, 14.2Hz), 3.15-3.31 (4H, m), 3.52 (2H, brs), 3.66 (1H, m), 4.95 (1H, dd, J=3.1, 10.0Hz), 7.06 (1H, dd, J=1.3, 8.2Hz), 7.36 (4H, m), 7.41 (1H, m), 7.47 (1H, m), 7.67 (1H, d, J=8.0Hz).

Example 49

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid (3-methylbutyl)amide · trifluoroacetate

¹H-NMR (CD₃OD) δ: 0.97 (6H, d, J=6.6Hz), 1.29 (3H, d, J=6.5Hz), 1.53 (2H, dt, J=7.2, 8.5Hz), 1.69 (1H, m), 3.02 (2H, dd, J=9.3, 14.2Hz), 3.27 (2H, m), 3.45 (2H, t, J=7.4Hz), 3.66 (1H, m), 4.95 (1H, dd, J=3.2, 10.0Hz), 7.34 (4H, m), 7.45 (1H, m), 7.53 (1H, dd, J=1.6, 8.4Hz), 7.66 (1H, d, J=8.7Hz), 7.91 (1H, m).

Example 50

(3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indol-6-yl)-piperidin-1-ylmethanone · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.5Hz), 1.41-1.65 (6H, m), 3.02 (1H, dd, J=9.3, 14.2Hz), 3.17 (1H, m), 3.27 (2H, m), 3.30-3.50 (2H, m), 3.66-3.88 (2H, m), 4.95 (1H, dd, J=3.2, 10.0Hz), 7.10 (2H, dd, J=1.4, 8.2Hz), 7.34 (4H, m), 7.45 (1H, m), 7.47 (1H, m), 7.66 (1H, d, J=8.2Hz).

Example 51

(3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indol-6-yl)-morpholin-4-ylmethanone · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.5Hz), 3.02 (1H, dd, J=9.4, 14.2Hz), 3.17 (1H, dd, J=10.1, 12.5Hz), 3.25-3.76 (6H, m), 4.95 (1H, dd, J=3.2, 10.0Hz), 7.13 (1H, dd, J=1.4, 8.2Hz), 7.27 (4H, m), 7.47 (1H, m), 7.50 (1H, m), 7.67 (1H, d, J=8.3Hz).

Example 52

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid ((1S)-1-hydroxymethyl-3-methylbutyl)amide · trifluoroacetate

¹H-NMR (CD₃OD) δ: 0.97 (6H, d, J=6.6Hz), 1.31 (3H, d,

J=6.5Hz), 1.49-1.57 (4H, m), 3.04 (2H, dd, J=9.1, 14.3Hz), 3.27 (2H, m), 3.61 (3H, d, J=5.6Hz), 3.66 (1H, m), 4.30 (1H, tt, J=4.7Hz), 4.93 (1H, dd, J=3.2, 10.2Hz), 7.34 (4H, m), 7.45 (1H, s), 7.54 (1H, m), 7.66 (1H, m), 7.91 (1H, d, J=16.2Hz).

5 Example 53

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxamide · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 3.04 (1H, dd, J=9.1, 14.3Hz), 3.17 (2H, m), 3.25-3.35 (3H, m), 3.66 (1H, m), 4.94 (1H, dd, J=3.2, 9.9Hz), 7.35 (4H, m), 7.45 (1H, m), 7.59 (1H, dd, J=1.5, 8.4Hz), 7.98 (1H, m).

Example 54

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid methylamide · trifluoroacetate

15 ¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.5Hz), 2.94 (3H, s), 3.03 (2H, dd, J=9.3, 14.2Hz), 3.27-3.35 (3H, m), 3.65 (3H, m), 4.94 (1H, dd, J=3.2, 10.0Hz), 7.34 (4H, m), 7.46 (1H, m), 7.53 (1H, d, J=1.5, 8.3Hz), 7.65 (1H, d, J=8.3Hz), 7.91 (1H, m).

Example 55

20 3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid dimethylamide · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.5Hz), 3.10-3.35 (6H, m), 3.04 (4H, m), 3.66 (1H, m), 4.93 (1H, dd, J=3.2, 10.2Hz), 7.13 (1H, dd, J=1.3, 8.2Hz), 7.27 (3H, m), 7.47 (2H, m), 7.66 (1H, d, J=8.2Hz).

25 Example 56

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carboxylic acid ethylamide · trifluoroacetate

$^1\text{H-NMR}$ (CD_3OD) δ : 1.24 (3H, t, $J=7.2\text{Hz}$), 1.31 (3H, d, $J=6.5\text{Hz}$), 3.02 (1H, dd, $J=9.3, 14.2\text{Hz}$), 3.16-3.35 (3H, m), 3.46 (2H, q, $J=7.2\text{Hz}$), 3.65 (1H, m), 4.94 (1H, dd, $J=3.2, 10.0\text{Hz}$), 7.35 (4H, m), 7.46 (1H, m), 7.54 (1H, dd, $J=1.5, 8.3\text{Hz}$), 7.65 (1H, d, $J=8.3\text{Hz}$), 7.91 (1H, m).

5 Example 57

Ethyl [(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carbonyl)amino]acetate · trifluoroacetate

10 $^1\text{H-NMR}$ (CD_3OD) δ : 1.31 (6H, m), 3.04 (1H, dd, $J=9.3, 14.2\text{Hz}$), 3.17 (1H, dd, $J=10.0, 12.5\text{Hz}$), 3.30 (2H, m), 3.66 (1H, m), 4.12 (2H, s), 4.20 (2H, q, $J=7.2\text{Hz}$), 4.95 (1H, d, $J=3.1\text{Hz}, 9.9\text{Hz}$), 7.37 (4H, m), 7.46 (1H, m), 7.58 (1H, dd, $J=1.5, 8.4\text{Hz}$), 7.67 (1H, d, $J=8.4\text{Hz}$), 7.97 (1H, m).

Example 58

Ethyl (2S)-2-[(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carbonyl)amino]propionate · trifluoroacetate

15 $^1\text{H-NMR}$ (CD_3OD) δ : 1.27 (3H, t, $J=7.1\text{Hz}$), 1.30 (3H, d, $J=6.5\text{Hz}$), 1.51 (3H, d, $J=7.3\text{Hz}$), 3.03 (1H, dd, $J=9.3, 14.2\text{Hz}$), 3.17 (1H, dd, $J=12.5\text{Hz}$), 3.20-3.35 (2H, m), 3.66 (1H, m), 4.19 (2H, q, $J=7.1\text{Hz}$), 4.60 (1H, q, $J=7.3\text{Hz}$), 4.93 (1H, d, $J=3.2\text{Hz}, 10.0\text{Hz}$), 7.37 (4H, m), 7.46 (1H, m), 7.59 (1H, dd, $J=1.5, 8.4\text{Hz}$), 7.67 (1H, d, $J=8.4\text{Hz}$), 7.96 (1H, m).

20 Example 59

Ethyl 3-[(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carbonyl)amino]propionate · trifluoroacetate

25 $^1\text{H-NMR}$ (CD_3OD) δ : 1.24 (3H, t, $J=7.1\text{Hz}$), 1.30 (3H, d, $J=6.5\text{Hz}$), 2.66 (3H, d, $J=6.8\text{Hz}$), 3.03 (1H, dd, $J=9.3, 14.2\text{Hz}$), 3.16-3.35 (3H, m), 3.66 (1H, m), 3.66 (2H, t, $J=6.8\text{Hz}$), 4.14 (2H, q, $J=7.1\text{Hz}$), 4.95 (1H, d, $J=1.8\text{Hz}, 9.9\text{Hz}$), 7.27 (4H, m), 7.45 (1H, m), 7.52 (1H, dd, $J=1.5, 8.4\text{Hz}$), 7.65 (1H, d, $J=8.4\text{Hz}$), 7.90 (1H, m).

Examples 60 to 62

The compounds of the following Examples are obtained in a similar manner to Example 24.

Example 60

5 [(3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carbonyl)amino]acetic acid · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 3.04 (1H, dd, J=9.3, 14.2Hz), 3.17 (1H, dd, J=10.0, 12.6Hz), 3.20-3.35 (2H, m), 3.66 (1H, m), 4.11 (2H, s), 4.95 (1H, d, J=3.1Hz, 10.0Hz), 7.35 (4H, m), 7.47 (1H, m),
10 7.59 (1H, dd, J=1.5, 8.4Hz), 7.67 (1H, d, J=8.4Hz), 7.97 (1H, m).

Example 61

Ethyl (2S)-2-[(3-((2R)-2-[2-(3-chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carbonyl)amino]propionate · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 1.52 (3H, d, J=7.3Hz), 3.03 (1H, dd, J=9.3, 14.2Hz), 3.18 (1H, dd, J=10.0, 12.6Hz), 3.20-3.35 (2H, m), 3.66 (1H, m), 4.62 (2H, q, J=7.3Hz), 4.95 (1H, d, J=3.2Hz, 10.0Hz), 7.35 (4H, m), 7.46 (1H, m), 7.60 (1H, dd, J=1.5, 8.4Hz), 7.66 (1H, d, J=8.4Hz), 7.97 (1H, m).
15

Example 62

20 3-[(3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-6-carbonyl)amino]propionic acid · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.5Hz), 2.65 (3H, d, J=6.8Hz), 3.02 (1H, dd, J=9.3, 14.1Hz), 3.16-3.35 (3H, m), 3.66 (1H, m), 3.69 (2H, t, J=6.8Hz), 4.94 (1H, d, J=3.1Hz, 10.0Hz), 7.35 (4H, m), 7.45 (1H, m), 7.53 (1H, d, J=8.4Hz), 7.65 (1H, d, J=8.4Hz), 7.92 (1H, s)
25

Reference Example 39

3-((2R)-2-[2-(4-Benzyloxy-3-methanesulfonylaminophenyl)-(2R)-2-

(triethylsilyloxy)ethylamino]propyl}-1H-indole-6-carboxylic acid diethylamide

To a solution of (R)-3-(2-aminopropyl)-1H-indole-6-carboxylic acid diethylamide (20.0 mg, 0.0742 mmol) in acetonitrile (1.0 ml) is added (R)-N-[2-benzyloxy-5-(2-iodo-1-triethylsilyloxyethyl)phenyl]-methanesulfonamide (41.1 mg, 0.0742 mmol), and the mixture is stirred at 110°C for 21 hours in a sealed vessel. The reaction solution is concentrated under reduced pressure, and the residue is purified by preparative TLC (a saturated ammonia solution in chloroform/methanol = 50/1) to give 3-((2R)-2-[2-(4-benzyloxy-3-methanesulfonylamino-phenyl)-(2R)-2-(triethylsilyloxy)ethylamino]propyl)-1H-indole-6-carboxylic acid diethylamide (19.2 mg, 0.0272 mmol, 37 %).

¹H-NMR (CDCl₃) δ: 0.44 (6H, m), 0.81 (9H, t, J=8.0Hz), 1.09 (3H, d, J=6.2Hz), 1.20 (6H, br), 2.63 (1H, dd, J=5.6, 11.3Hz), 2.76 (2H, m), 2.85 (3H, s), 2.94 (2H, m), 3.30 (4H, br), 4.66 (1H, t, J=6.1Hz), 5.07 (2H, s), 6.80 (1H, d, J=8.4Hz), 6.93 (1H, dd, J=2.1, 8.4Hz), 6.97 (1H, d, J=2.1Hz), 7.05 (1H, dd, J=1.3, 8.1Hz), 7.35 (1H, s), 7.40 (7H, m), 7.50 (1H, d, J=8.2Hz), 8.41 (1H, br).

Example 63

3-((2R)-2-[(2R)-2-Hydroxy-2-(4-hydroxy-3-methanesulfonylamino-phenyl)ethylamino]propyl)-1H-indole-6-carboxylic acid diethylamide trifluoroacetate

To a solution of 3-((2R)-2-[2-(4-benzyloxy-3-methanesulfonylamino-phenyl)-(2R)-2-(triethylsilyloxy)ethylamino]propyl)-1H-indole-6-carboxylic acid diethylamide (18.0 mg, 0.0254 mmol) in methylene chloride (1.0 ml) is added trifluoroacetic acid (100 μl), and the mixture is stirred at room temperature for one hour. To the reaction solution is

added a 5 % aqueous potassium carbonate solution, and the mixture is extracted with chloroform. The organic layers are combined and washed with a saturated brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the residue (19 mg).

5 To a solution of the residue in methanol (1.0 ml) is added 10 % palladium on carbon (50 % wet, 5.0 mg), and the mixture is stirred under hydrogen atmosphere for 4 hours. The reaction solution is filtered through celite and the filtrate is concentrated under reduced pressure. The resulting residue is purified by preparative reversed
10 phase HPLC (octadecylsilyl, trifluoroacetic acid/acetonitrile/water) to give 3-((2R)-2-[(2R)-2-hydroxy-2-(4-hydroxy-3-methanesulfonylamino-phenyl)ethylamino]propyl)-1H-indole-6-carboxylic acid diethylamide · trifluoroacetate (950 mg, 2.22 mmol, 63 %).

¹H-NMR (CD₃OD) δ: 1.14-1.27 (6H, br), 1.30 (3H, d, J=6.6Hz),
15 2.93 (3H, s), 3.01 (1H, dd, J=9.2, 14.2Hz), 3.18 (2H, m), 3.27 (1H, m), 3.04 (1H, m), 3.30 (2H, br), 3.56 (2H, br), 3.66 (1H, m), 4.85 (1H, m), 6.89 (1H, d, J=8.3Hz), 7.07 (1H, dd, J=1.4, 8.2Hz), 7.11 (1H, dd, J=2.2, 8.5Hz), 7.31 (1H, s), 7.39(1H, d, J=2.2Hz), 7.43 (1H, m), 7.66 (1H, d, J=8.2Hz).

20 Reference Example 40

3-[(2R)-2-(2-Pyridin-3-yl-(2R)-2-(triethylsilyloxy)ethylamino)propyl]-1H-indole-6-carboxylic acid diethylamide

The title compound is obtained from (R)-3-(2-aminopropyl)-1H-indole-6-carboxylic acid diethylamide in a similar manner to Reference
25 Example 38.

¹H-NMR (CDCl₃) δ: 0.44 (6H, m), 0.81 (9H, t, J=7.8Hz), 1.08 (3H, d, J=6.2Hz), 1.20 (6H, br), 2.70 (2H, m), 2.80 (1H, dd, J=6.8, 14.2Hz),

2.96 (2H, m), 3.30 (4H, br), 4.76 (1H, m), 6.99 (1H, d, J=1.3Hz), 7.07 (1H, dd, J=1.3, 8.1Hz), 7.15 (1H, dd, J=4.9, 8.1Hz), 7.41 (1H, s), 7.52 (1H, d, J=8.1Hz), 7.56 (1H, m), 8.44 (1H, br), 8.46 (1H, dd, J=1.7, 4.8Hz), 8.50 (1H, d, J=2.0Hz).

5 Example 64

3-[(2R)-2-((2R)-2-Hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indole-7-carboxylic acid diethylamide

10 The title compound is obtained from 3-[(2R)-2-(2-pyridin-3-yl-(2R)-2-(triethylsilyloxy)ethylamino)propyl]-1H-indole-6-carboxylic acid diethylamide in a similar manner to Example 36.

15 ¹H-NMR (CDCl₃) δ: 1.13 (3H, d, J=6.2Hz), 1.20 (6H, br), 2.66 (2H, dd, J=9.2, 12.2Hz), 2.80 (2H, m), 2.87 (2H, d, J=3.7, 12.2Hz), 3.05 (1H, m), 3.40 (4H, m), 4.54 (1H, dd, J=3.6, 9.1Hz), 7.06 (1H, d, J=2.2Hz), 7.10 (1H, dd, J=1.4, 8.2Hz), 7.22 (1H, m), 7.43 (1H, s), 7.56 (1H, d, J=8.1Hz), 7.64 (1H, dt, 7.9, 1.8Hz), 8.47 (1H, dd, J=1.6, 4.8Hz), 8.51 (1H, d, J=2.1Hz), 8.68 (1H, brs).

Reference Example 41

3-((Dimethylamino)methyl)-N,N-diethyl-1H-indole-6-carboxamide

20 The title compound is obtained from 1H-indole-6-carboxylic acid diethylamide in a similar manner to Reference Example 18.

¹H-NMR (CDCl₃) δ: 1.18 (6H, brs), 2.27 (6H, s), 3.34-3.57 (4H, m), 3.60 (2H, s), 7.11 (1H, dd, J=8.1, 1.4Hz), 7.15 (1H, d, J=1.4Hz), 7.42 (1H, brs), 7.67 (1H, d, J=8.1Hz), 8.80 (1H, brs).

Reference Example 42

25 3-(Cyanomethyl)-N,N-diethyl-1H-indole-6-carboxamide

The title compound is obtained from 3-((dimethylamino)methyl)-N,N-diethyl-1H-indole-6-carboxamide in a similar manner to Reference

Example 19.

^1H -NMR (CDCl_3) δ : 1.14-1.25 (6H, m), 3.32-3.58 (4H, m), 3.83 (1H, d, $J=0.9\text{Hz}$), 7.15 (1H, dd, $J=8.1, 1.3\text{Hz}$), 7.23-7.24 (1H, m), 7.47 (1H, brs), 7.56 (1H, d, $J=8.1\text{Hz}$), 8.97 (1H, brs).

5 Reference Example 43

(6-((Diethylamino)carbonyl)-1H-indol-3-yl)acetic acid

The title compound is obtained from 3-(cyanomethyl)-N,N-diethyl-1H-indole-6-carboxamide in a similar manner to Reference Example 20.

10 ^1H -NMR (CDCl_3) δ : 1.08-1.29 (6H, m), 3.29-3.60 (4H, m), 3.58 (2H, s), 6.22 (1H, d, $J=1.3\text{Hz}$), 7.01 (1H, dd, $J=8.1, 1.3\text{Hz}$), 7.36 (1H, s), 7.45 (1H, d, $J=8.1\text{Hz}$), 9.08 (1H, s).

Reference Example 44

Methyl (6-((diethylamino)carbonyl)-1H-indol-3-yl)acetate

15 The title compound is obtained from (6-((diethylamino)-carbonyl)-1H-indol-3-yl)acetic acid in a similar manner to Reference Example 28.

20 ^1H -NMR (CDCl_3) δ : 1.19 (6H, brs), 3.35-3.54 (4H, m), 3.71 (3H, s), 3.78 (2H, s), 7.13 (1H, dd, $J=8.2, 1.3\text{Hz}$), 7.22 (1H, m), 7.44 (1H, brs), 7.59 (1H, d, $J=8.2\text{Hz}$), 8.67 (1H, brs).

Reference Example 45

N,N-Diethyl-3-(2-hydroxyethyl)-1H-indole-6-carboxamide

25 The title compound is obtained from methyl (6-((diethylamino)-carbonyl)-1H-indol-3-yl)acetic acid in a similar manner to Reference Example 29.

^1H -NMR (CDCl_3) δ : 1.19 (6H, brs), 1.54 (1H, t, $J=6.1\text{Hz}$), 3.02 (2H, t, $J=6.3\text{Hz}$), 3.25-3.53 (4H, m), 3.90 (2H, td, $J=6.3, 6.1\text{Hz}$), 7.11-

7.13 (2H, m), 7.43 (1H, s), 7.60 (1H, d, J=8.1Hz), 8.37 (1H, brs).

Reference Example 46

N,N-Diethyl-3-(2-oxoethyl)-1H-indole-6-carboxamide

The title compound is obtained from N,N-diethyl-3-(2-hydroxyethyl)-1H-indole-6-carboxamide in a similar manner to Reference Example 24.

¹H-NMR (CDCl₃) δ: 1.11-1.25 (6H, m), 3.31-3.59 (4H, m), 3.76 (2H, dd, J=2.4, 0.5Hz), 7.07-7.10 (2H, m), 7.42 (1H, m), 7.47 (1H, d, J=8.2Hz), 9.66 (1H, brs), 9.70 (1H, t, J=2.4Hz).

Example 65

N,N-Diethyl-3-(2-(((1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-methylethyl)amino)ethyl)-1H-indole-6-carboxamide

The title compound is obtained from N,N-diethyl-3-(2-oxoethyl)-1H-indole-6-carboxamide in a similar manner to Example 30.

¹H-NMR (CDCl₃) δ: 1.15-1.32 (6H, m), 1.23 (3H, d, J=6.1Hz), 2.50-2.65 (2H, m), 2.80-2.87 (1H, m), 2.91-2.96 (1H, m), 3.0-3.09 (1H, m), 3.38-3.64 (4H, m), 4.03 (1H, d, J=8.5Hz), 6.21 (2H, d, J=8.5Hz), 6.65 (2H, d, J=8.5Hz), 6.91 (1H, d, J=2.1Hz), 6.97 (1H, dd, J=8.1, 2.1Hz), 7.35 (1H, s), 7.39 (1H, d, J=8.1Hz), 8.30 (1H, brs).

Example 66

(R)-N,N-Diethyl-3-{2-[2-(3-chlorophenyl)-2-hydroxyethylamino]ethyl}-1H-indole-7-carboxamide

The title compound is obtained from N,N-diethyl-3-(2-oxoethyl)-1H-indole-7-carboxamide and (R)-2-amino-1-(3-chlorophenyl)ethanol in a similar manner to Example 30.

¹H-NMR (CDCl₃) δ: 1.26 (6H, t, J=7.1Hz), 2.65 (1H, dd, J=12.3, 9.2Hz), 2.90 (1H, dd, J=12.3, 3.6Hz), 2.92-3.06 (4H, m), 3.54 (4H, q,

J=7.1Hz), 4.65 (1H, dd, J=9.2, 3.6Hz), 7.03 (1H, d, J=1.5Hz), 7.08 (1H, dd, J=7.7, 7.5Hz), 7.17-7.23 (4H, m), 7.35 (1H, s), 7.64 (1H, d, J=7.7Hz), 9.02 (1H, s).

Example 67

5 (3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indol-7-yl)-pyrrolidin-1-ylmethanone · trifluoroacetate

The title compound is obtained in a similar manner to Example 11.

10 ¹H-NMR (CD₃OD) δ:1.31 (3H, d, J=6.5Hz), 1.85-2.03 (4H, m), 3.04 (1H, dd, J=9.2, 14.2Hz), 3.17 (1H, dd, J=10.2, 12.6Hz), 3.24-3.47 (3H, m), 3.68 (4H, m), 4.93 (1H, dd, J=6.7, 10.0Hz), 7.14 (1H, dd, J=7.4, 7.4Hz), 7.27-7.39 (5H, m), 7.47 (1H, s), 7.72 (1H, d, J=7.9Hz).

Example 68

15 Ethyl 3-{2-[(1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethyl-amino]ethyl}-1H-indole-7-carboxylate

To a solution of N,N-diethyl-3-(2-(((1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-methylethyl)amino)ethyl)-1H-indole-7-carboxamide (215.5 mg, 0.526 mmol) in acetonitrile (18 mL) is added disodium hydrogen phosphate (112 mg, 0.789 mmol) under nitrogen atmosphere, and thereto is further added a 1N solution of triethyloxonium tetra-
20 fluoroborate in methylene chloride (1.58 mL, 1.58 mmol), and the mixture is stirred at room temperature for 15 hours. To the reaction solution are added a saturated aqueous sodium hydrogen carbonate solution (18 mL) and sodium hydrogen carbonate (200 mg), and the
25 mixture is further stirred for 2 hours. The mixture is separated into water and chloroform, and the organic layer is washed with a saturated aqueous sodium hydrogen carbonate solution and dried over anhydrous

sodium sulfate. The solvent is evaporated, and the obtained crude product is purified by silica gel column chromatography (chloroform: methanol = 50:1 → 5:1) to give ethyl 3-{2-[(1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]ethyl}-1H-indole-7-carboxylate (128.4 mg, 0.336 mmol, 64 %).

¹H-NMR (CDCl₃) δ: 0.89 (3H, d, J=6.4Hz), 1.45 (3H, t, J=7.1Hz), 2.80-2.86 (1H, m), 2.89-2.97 (3H, m), 3.04-3.08 (1H, m), 4.45 (2H, q, J=7.1Hz), 4.57 (1H, d, J=4.9Hz), 6.68 (2H, d, J=8.6Hz), 7.02 (1H, d, J=2.1Hz), 7.06 (2H, d, J=8.6Hz), 7.14 (1H, dd, J=7.8, 7.5Hz), 7.80 (1H, d, J=7.8Hz), 7.90 (1H, d, J=7.5Hz), 9.60 (1H, brs).

Example 69

3-{2-[(1S,2R)-2-Hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]ethyl}-1H-indole-7-carboxylic acid

To a solution of ethyl 3-{2-[(1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]ethyl}-1H-indole-7-carboxylate (115 mg, 0.301 mmol) in tetrahydrofuran (10 mL) and water (5 mL) is added lithium hydroxide (36 mg, 1.51 mmol), and the mixture is stirred at room temperature for 15 hours. The pH value of the reaction solution is adjusted to pH 7 with 1N aqueous hydrochloric acid solution, and the solvent is distilled off. The residue is isolated by reversed phase column (octadecylsilyl, water → water-methanol = 10:1 → 9:1 → 6:1) to give 3-{2-[(1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]ethyl}-1H-indole-7-carboxylic acid (69.3 mg, 0.195 mmol, 65 %).

¹H-NMR (CD₃OD) δ: 1.08 (3H, d, J=6.7Hz), 3.20-3.26 (2H, m), 3.33-3.43 (3H, m), 5.01 (1H, d, J=3.1Hz), 6.75 (2H, d, J=8.6Hz), 7.03-7.15 (3H, m), 7.30 (1H, s), 7.74 (1H, d, J=7.8Hz), 7.79 (1H, d, J=7.3Hz).

Example 70

3-{2-[(1S,2R)-2-Hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]ethyl}-1H-indole-7-carboxylic acid ethylamide

To a solution of 3-{2-[(1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]ethyl}-1H-indole-7-carboxylic acid (29.1 mg, 0.0821 mmol) in N,N-dimethylformamide (3 mL) are added ethylamine hydrochloride (67 mg, 0.822 mmol), triethylamine (171 μ L, 1.23 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (WSCl) hydrochloride (78.7 mg, 0.411 mmol) and 1-hydroxybenzotriazole (HOBT, 55.5 mg, 0.411 mmol), and the mixture is stirred at room temperature for 17 hours. The reaction solution is poured into a saturated aqueous sodium hydrogen carbonate solution, and the mixture is separated into chloroform. The organic layer is washed with a saturated brine and dried over anhydrous sodium sulfate. The solvent is removed by distillation, and the resulting crude product is purified by preparative TLC (a saturated ammonia solution in chloroform:methanol = 20:1) to give 3-{2-[(1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]ethyl}-1H-indole-7-carboxylic acid ethylamide (7.5mg, 0.0197 mmol, 24 %).

$^1\text{H-NMR}$ (CDCl_3) δ : 1.02 (3H, d, $J=6.3\text{Hz}$), 1.31 (3H, t, $J=7.3\text{Hz}$), 2.68 (1H, qd, $J=6.3, 6.3\text{Hz}$), 2.76-2.90 (2H, m), 2.97 (1H, dt, $J=14.3, 5.3\text{Hz}$), 3.02-3.08 (1H, m), 3.57 (2H, qd, $J=7.3, 5.1\text{Hz}$), 4.39 (1H, d, $J=6.3\text{Hz}$), 6.38 (1H, brt, $J=5.1\text{Hz}$), 6.56 (2H, d, $J=8.5\text{Hz}$), 6.85 (1H, d, $J=2.1\text{Hz}$), 6.91 (2H, d, $J=8.5\text{Hz}$), 7.10 (1H, dd, $J=7.8, 7.2\text{Hz}$), 7.36 (1H, d, $J=7.2\text{Hz}$), 7.72 (1H, d, $J=7.8\text{Hz}$), 9.70 (1H, s).

Examples 71 to 74

The compounds of the following Examples are obtained from 3-{2-[(1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]ethyl}-

1H-indole-7-carboxylic acid in a similar manner to Example 11.

Example 71

(3-{2-[(1S,2R)-2-Hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]-ethyl}-1H-indol-7-yl)-piperidin-1-ylmethanone · trifluoroacetate

5 $^1\text{H-NMR}$ (CD_3OD) δ : 1.08 (3H, d, $J=6.5\text{Hz}$), 1.50 (2H, brs), 1.70 (4H, m), 3.22 (2H, m), 3.41 (3H, m), 3.75 (4H, brs), 4.99 (1H, d, $J=3.1\text{Hz}$), 6.76 (2H, m), 7.14 (4H, m), 7.27 (1H, s), 7.70 (1H, dd, $J=2.4, 6.9\text{Hz}$).

Example 72

10 (3-{2-[(1S,2R)-2-Hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]-ethyl}-1H-indol-7-yl)morpholin-4-ylmethanone · trifluoroacetate

$^1\text{H-NMR}$ (CD_3OD) δ : 1.08 (3H, d, $J=6.5\text{Hz}$), 3.22 (2H, m), 3.41 (3H, m), 3.70 (8H, brs), 4.99 (1H, d, $J=3.1\text{Hz}$), 6.76 (2H, m), 7.15 (2H, m), 7.18 (2H, m), 7.29 (1H, s), 7.73 (1H, dd, $J=1.3, 7.7\text{Hz}$).

15 Example 73

(3-{2-[(1S,2R)-2-Hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]-ethyl}-1H-indol-7-yl)-(4-methylpiperazin-1-yl)methanone · 2 trifluoroacetate

20 $^1\text{H-NMR}$ (CD_3OD) δ : 1.08 (3H, d, $J=6.5\text{Hz}$), 2.93 (3H, s), 3.29 (4H, m), 3.39 (3H, m), 3.50 (4H, brs), 4.43 (2H, brs), 5.00 (1H, d, $J=3.1\text{Hz}$), 6.76 (2H, m), 7.15 (2H, d, $J=8.4\text{Hz}$), 7.18 (1H, t, $J=7.8\text{Hz}$), 7.28 (1H, d, $J=6.9\text{Hz}$), 7.31 (1H, s), 7.78 (1H, d, $J=8.0\text{Hz}$).

Example 74

25 3-{2-[(1S,2R)-2-Hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]-ethyl}-1H-indole-7-carboxylic acid dimethylamide · trifluoroacetate

$^1\text{H-NMR}$ (CD_3OD) δ : 1.08 (3H, d, $J=6.5\text{Hz}$), 2.98 (3H, brs), 3.16 (3H, brs), 3.22 (2H, m), 3.39 (3H, m), 5.00 (1H, d, $J=3.0\text{Hz}$), 6.76 (2H,

m), 7.16 (4H, m), 7.28 (1H, s), 7.88 (1H, dd, J=1.3, 7.7Hz).

Examples 75 to 76

The compounds of the following Examples are obtained from 3-
 [(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indole-7-
 5 carboxylic acid · 2 trifluoroacetate in a similar manner to Example 11.

Example 75

{3-[(2R)-2-((2R)-2-Hydroxy-2-pyridin-3-ylethylamino)propyl]-1H-indol-7-
 yl}-pyrrolidin-1-ylmethanone · 2 trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.32 (3H, d, J=6.5Hz), 1.90 (2H, m), 2.01
 10 (2H, m), 3.30 (1H, dd, J=9.3, 14.2Hz), 3.34 (2H, m), 3.43 (3H, m), 3.69
 (3H, m), 5.25 (1H, dd, J=3.0, 9.9Hz), 7.13 (1H, t, J=7.8Hz), 7.30 (2H, m),
 7.71 (1H, m), 8.03 (1H, dd, J=5.7, 8.0Hz), 8.58 (1H, m), 8.80 (1H, d,
 J=5.5Hz), 8.89 (1H, s).

Example 76

15 Azepan-1-yl-{3-[(2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethylamino)propyl]-
 1H-indol-7-yl}methanone · 2 trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.32 (3H, d, J=6.5Hz), 1.57 (4H, m), 1.68
 (2H, m), 1.90 (2H, m), 3.03 (1H, dd, J=9.3, 14.2Hz), 3.34 (2H, m), 3.43
 (3H, m), 3.69 (3H, m), 3.76 (2H, m), 5.27 (1H, dd, J=3.0, 9.9Hz), 7.13
 20 (2H, m), 7.28 (1H, s), 7.71 (1H, dd, J=2.3, 6.7Hz), 8.05 (1H, dd, J=5.7,
 8.0Hz), 8.58 (1H, m), 8.81 (1H, d, J=5.5Hz), 8.89 (1H, s).

Examples 77 to 89

The compounds of the following Examples 77 to 89 are obtained
 in a similar manner to Example 11.

25 Example 77

N-Benzyl-3-((2R)-2-(((2R)-2-(3-chlorophenyl)-2-hydroxyethyl)amino)-
 propyl)-1H-indole-7-carboxamide · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 3.05 (1H, dd, J=9.0, 14.3Hz), 3.16-3.31 (3H, m), 3.64-3.69 (1H, m), 4.63 (2H, s), 4.93 (1H, dd, J=3.3, 9.9Hz), 7.15 (1H, dd, J=7.7, 7.7Hz), 7.21-7.39 (9H, m), 7.67 (1H, d, J=7.0Hz), 7.81 (1H, dd, J=0.8, 7.9Hz).

5 Example 78

3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-N-phenyl-1H-indole-7-carboxamide · trifluoroacetate

10 ¹H-NMR (CD₃OD) δ: 1.33 (3H, d, J=6.5Hz), 3.07 (1H, dd, J=9.1, 14.3Hz), 3.15-3.33 (3H, m), 3.65-3.71 (1H, m), 4.95 (1H, dd, J=3.6, 10.1Hz), 7.12-7.17 (1H, m), 7.21 (1H, dd, J=7.7, 7.7Hz), 7.31-7.39 (6H, m), 7.45-7.46 (1H, m), 7.72-7.75 (2H, m), 7.85 (2H, dd, J=7.6, 9.1Hz).

Example 79

(1R)-2-(((1R)-2-{7-[(4-Benzylpiperazin-1-yl)carbonyl]-1H-indol-3-yl}-1-methylethyl)amino)-1-(3-chlorophenyl)ethanol · trifluoroacetate

15 ¹H-NMR (CD₃OD) δ: 1.29 (3H, d, J=6.6Hz), 3.03 (1H, dd, J=9.5, 14.2Hz), 3.18 (1H, dd, J=10.2, 12.6Hz), 3.25-3.35 (6H, m), 3.65-3.69 (1H, m), 3.90 (4H, m), 4.37 (2H, s), 4.96 (1H, dd, J=3.1, 10.1Hz), 7.17 (1H, t, J=7.4Hz), 7.27 (1H, dd, J=0.8, 7.2Hz), 7.31-7.39 (5H, m), 7.46-7.53 (6H, m), 7.78 (1H, dd, J=0.9, 7.9Hz).

20 Example 80

(1R)-1-(3-Chlorophenyl)-2-(((1R)-1-methyl-2-{7-[(4-phenylpiperazin-1-yl)-carbonyl]-1H-indol-3-yl}ethyl)amino)ethanol · 2 trifluoroacetate

25 ¹H-NMR (CD₃OD) δ: 1.32 (3H, d, J=6.5Hz), 3.04 (1H, dd, J=9.2, 14.3Hz), 3.15-3.32 (7H, m), 3.65-3.70 (1H, m), 3.80 (4H, m), 4.94 (1H, dd, J=2.8, 9.7Hz), 6.88-6.92 (1H, m), 7.01 (2H, dd, J=0.9, 8.7Hz), 7.15 (1H, t, J=7.3Hz), 7.21-7.42 (8H, m), 7.48 (1H, m), 7.74 (1H, dd, J=1.0, 7.9Hz).

Example 81

3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-N-cyclopentyl-1H-indole-7-carboxamide · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 1.61-1.68 (4H, m),
 5 1.78-1.80 (2H, m), 2.02-2.07 (2H, m), 3.04 (1H, dd, J=9.1, 14.4Hz),
 3.14-3.29 (3H, m), 3.63-3.69 (1H, m), 4.93 (1H, dd, J=3.1, 9.8Hz), 7.13
 (1H, t, J=7.7Hz), 7.31-7.38 (4H, m), 7.44 (1H, m), 7.63 (1H, dd, J=0.6,
 7.5Hz), 7.79 (1H, dd, J=0.7, 7.9Hz).

Example 82

10 3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-N-cyclohexyl-1H-indole-7-carboxamide · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.24-1.45 (6H, m), 1.31 (3H, d, J=6.6Hz),
 1.81-1.85 (2H, m), 1.97-2.02 (2H, m), 3.04 (1H, dd, J=9.1, 14.3Hz),
 3.14-3.27 (3H, m), 3.65-3.67 (1H, m), 3.93 (1H, m), 4.93 (1H, dd, J=3.4,
 15 9.9Hz), 7.13 (1H, dd, J=7.7, 7.7Hz), 7.31-7.38 (4H, m), 7.44 (1H, s),
 7.63 (1H, d, J=7.1Hz), 7.78 (1H, d, J=7.9Hz).

Example 83

(1R)-1-(3-Chlorophenyl)-2-(((1R)-1-methyl-2-{7-[(4-methylpiperidin-1-yl)-carbonyl]-1H-indol-3-yl}ethyl)amino)ethanol · trifluoroacetate

20 ¹H-NMR (CD₃OD) δ: 0.97 (3H, d, J=6.4Hz), 1.31 (3H, d,
 J=6.5Hz), 1.03-1.93 (5H, m), 2.81-3.20 (4H, m), 3.03 (1H, dd, J=9.1,
 14.3Hz), 3.17 (1H, dd, J=10.1, 12.6Hz), 3.24-3.28 (2H, m), 3.64-3.69
 (1H, m), 4.94 (1H, d, J=3.2, 10.1Hz), 7.12-7.18 (2H, m), 7.27 (1H, m),
 7.31-7.39 (3H, m), 7.47 (1H, m), 7.68-7.73 (1H, m).

25 Example 84

(1R)-2-(((1R)-2-[7-(Azepan-1-ylcarbonyl)-1H-indol-3-yl]-1-methylethyl)-amino)-1-(3-chlorophenyl)ethanol · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 1.57-1.69 (6H, m),
 1.90 (2H, m), 3.03 (1H, dd, J=9.2, 14.3Hz), 3.18 (1H, dd, J=10.2,
 12.6Hz), 3.25-3.28 (2H, m), 3.42 (2H, m), 3.63-3.69 (1H, m), 3.74-3.77
 (2H, m), 4.94 (1H, dd, J=2.8, 9.7Hz), 7.12-7.16 (2H, m), 7.27 (1H, s),
 5 7.31-7.42 (3H, m), 7.47 (1H, s), 7.67-7.72 (1H, m).

Example 85

3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-N-
 tetrahydro-2H-pyran-4-yl-1H-indole-7-carboxamide · trifluoroacetate

10 ¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 1.67-1.77 (2H, m),
 1.91-1.95 (2H, m), 3.04 (1H, dd, J=9.1, 14.3Hz), 3.17-3.28 (3H, m),
 3.51-3.57 (2H, m), 3.65 (1H, m), 3.98-4.02 (2H, m), 4.16-4.21 (1H, m),
 4.94 (1H, dd, J=3.1, 9.9Hz), 7.14 (1H, dd, J=7.7, 7.7Hz), 7.31-7.36 (4H,
 m), 7.44 (1H, s), 7.65 (1H, d, J=7.5Hz), 7.80 (1H, dd, J=0.8, 8.0Hz).

Example 86

15 3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-N-
 cyclohexyl-N-methyl-1H-indole-7-carboxamide · trifluoroacetate

20 ¹H-NMR (CD₃OD) δ: 0.87-1.15 (2H, m), 1.30 (3H, d, J=6.5Hz),
 1.46-1.93 (8H, m), 2.90 (3H, m), 3.03 (1H, dd, J=9.2, 14.2Hz), 3.18 (1H,
 dd, J=10.2, 12.6Hz), 3.25-3.28 (2H, m), 3.47 (1H, m), 3.64-3.69 (1H, m),
 4.95 (1H, dd, J=3.1, 10.1Hz), 7.12-7.16 (2H, m), 7.26 (1H, s), 7.32-7.40
 (3H, m), 7.48 (1H, s), 7.70 (1H, d, J=8.9Hz).

Example 87

25 3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-N-
 methyl-N-(1-methylpiperidin-4-yl)-1H-indole-7-carboxamide · 2
 trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.6Hz), 2.13-2.23 (4H, m),
 2.85 (3H, s), 2.96 (3H, s), 3.03 (1H, dd, J=9.4, 14.2Hz), 3.18 (1H, dd,

J=10.2, 12.6Hz), 3.25-3.31 (2H, m), 3.31-3.68 (3H, m), 4.96 (1H, dd, J=3.1, 10.0Hz), 7.15 (1H, dd, J=7.3, 7.3Hz), 7.18-7.20 (1H, m), 7.29 (1H, s), 7.32-7.38 (3H, m), 7.48 (1H, s), 7.74 (1H, dd, J=1.4, 7.5Hz).

Example 88

5 (1R)-2-(((1R)-2-{7-[(4-Benzylpiperidin-1-yl)carbonyl]-1H-indol-3-yl}-1-methylethyl)amino)-1-(3-chlorophenyl)ethanol · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 1.15-1.89 (5H, m), 2.57 (2H, d, J=6.9Hz), 3.00 (2H, m), 3.03 (1H, dd, J=9.1, 14.3Hz), 3.17 (1H, dd, J=10.2, 12.5Hz), 3.24-3.29 (2H, m), 3.64-3.69 (3H, m), 4.93 (1H, dd, J=3.1, 10.0Hz), 7.13-7.16 (5H, m), 7.22-7.26 (3H, m), 7.31-7.37 (3H, m), 7.47 (1H, s), 7.70 (1H, dd, J=2.1, 6.7Hz).

Example 89

(1R)-1-(3-Chlorophenyl)-2-(((1R)-2-{7-[(3,5-dimethylpiperidin-1-yl)-carbonyl]-1H-indol-3-yl}-1-methylethyl)amino)ethanol · trifluoroacetate

15 ¹H-NMR (CD₃OD) δ: 0.73-0.99 (8H, m), 1.31 (3H, d, J=6.5Hz), 1.51-1.89 (2H, m), 2.36-2.60 (2H, m), 3.04 (1H, dd, J=9.2, 14.3Hz), 3.18 (1H, dd, J=10.3, 12.6Hz), 3.25-3.28 (2H, m), 3.53-3.97 (2H, m), 3.64-3.70 (1H, m), 4.94 (1H, dd, J=3.0, 9.9Hz), 7.12-7.18 (2H, m), 7.27 (1H, s), 7.32-7.40 (3H, m), 7.48 (1H, s), 7.70-7.74 (1H, m).

20 Example 90

Ethyl 3-((2R)-2-(((2R)-2-(3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1H-indole-7-carboxylate

3-((2R)-2-[2-(3-Chlorophenyl)-(2R)-2-hydroxyethylamino]propyl)-1H-indole-7-carboxylic acid (50 mg, 0.134 mmol) is dissolved in ethanol (5 mL), and thereto are added conc. sulfuric acid (0.05 mL) and Molecular Sieves 4A powder (trade name, 0.3 g), and the mixture is heated at 150°C for 16 hours. After cooling, the insoluble materials are

removed by filtration, and a saturated aqueous sodium hydrogen carbonate solution is added to the filtrate. The mixture is evaporated under reduced pressure to remove ethanol, and the resultant is extracted with ethyl acetate. The organic layer is washed with a saturated brine, dried over sodium sulfate, and concentrated. The residue is purified by silica gel column chromatography (chloroform/ethanol = 4 / 1) to give the title compound (50 mg, yield: 93 %).

¹H-NMR (CDCl₃) δ: 1.13 (3H, d, J=6.2Hz), 1.45 (3H, t, J=7.1Hz), 3.04 (1H, dd, J=6.4, 12.7Hz), 2.85-2.89 (3H, m), 3.02-3.06 (1H, m), 4.45 (2H, q, J=7.1Hz), 4.53 (1H, dd, J=3.7, 8.9Hz), 7.12-7.17 (3H, m), 7.20-7.22 (2H, m), 7.32 (1H, s), 7.80 (1H, d, J=7.8Hz), 7.91 (1H, d, J=7.4Hz), 9.70 (1H, brs).

Reference Example 47

Ethyl 3-((2R)-2-[(5R)-5-(3-chlorophenyl)-2-oxo-1,3-oxazolidin-3-yl]-propyl)-1H-indole-7-carboxylate

The title compound is obtained from ethyl 3-((2R)-2-[(2R)-2-(3-chlorophenyl)-2-hydroxyethyl]amino)propyl)-1H-indole-7-carboxylate in a similar manner to Reference Example 17.

¹H-NMR (CDCl₃) δ: 1.32 (3H, d, J=6.8Hz), 1.46 (3H, t, J=7.1Hz), 2.96-2.98 (2H, m), 3.22 (1H, dd, J=6.5, 8.5Hz), 3.83 (1H, dd, J=8.7, 8.8Hz), 4.42-4.49 (3H, m), 5.34 (1H, dd, J=6.6, 8.9Hz), 6.86 (1H, d, J=7.7Hz), 7.06-7.14 (4H, m), 7.20-7.23 (1H, m), 7.80 (1H, d, J=7.8Hz), 7.89 (1H, dd, J=0.8, 7.5Hz), 9.65 (1H, brs).

Reference Example 48

3-((2R)-2-[(5R)-5-(3-Chlorophenyl)-2-oxo-1,3-oxazolidin-3-yl]propyl)-1H-indole-7-carboxylic acid

A solution of ethyl 3-((2R)-2-[(5R)-5-(3-chlorophenyl)-2-oxo-1,3-

oxazolidin-3-yl]propyl}-1H-indole-7-carboxylate (55 mg, 0.129 mmol) and potassium hydroxide (70 mg, 1.3 mmol) in water (1 mL) and ethanol (2 mL) is stirred at room temperature for 3 days. The reaction mixture is neutralized with 1N hydrochloric acid, and the precipitated crystals are collected by filtration to give the title compound (44 mg, yield: 86 %).

¹H-NMR (CD₃OD) δ: 1.34 (3H, d, J=6.8Hz), 2.98-3.00 (2H, m), 3.24-3.28 (1H, m), 3.95 (1H, dd, J=9.1, 9.1Hz), 4.37-4.46 (1H, m), 5.42 (1H, dd, J=6.2, 9.1Hz), 6.77 (1H, d, J=7.8Hz), 7.02-7.13 (4H, m), 7.20-7.23 (1H, m), 7.79-7.82 (2H, m).

Example 91

(1R)-1-(3-Chlorophenyl)-2-[(1R)-2-{7-[(2,6-dimethylpiperidin-1-yl)-carbonyl]-1H-indol-3-yl}-1-methylethyl)amino]ethanol · trifluoroacetate

To a solution of 3-[(2R)-2-[(5R)-5-(3-chlorophenyl)-2-oxo-1,3-oxazolidin-3-yl]propyl]-1H-indole-7-carboxylic acid (30 mg, 0.075 mmol) in tetrahydrofuran (5 mL) is added thionyl chloride (0.011 mL, 0.15 mmol), and the mixture is stirred at room temperature for 4 hours. The reaction solution is concentrated, and thereto are added tetrahydrofuran (3 mL) and 2,6-dimethylpiperidine (85 mg, 0.75 mmol). The mixture is stirred at room temperature for one day. The reaction solution is concentrated, and thereto are added ethanol (1 mL)/water (1 mL) and potassium hydroxide (0.60 g). The mixture is stirred at 70°C for 6.5 hours. The reaction solution is acidified with hydrochloric acid, and purified by preparative HPLC (trifluoroacetic acid-acetonitrile-water) to give the title compound (8 mg, yield: 18 %).

¹H-NMR (CD₃OD) δ: 1.31 (9H, m), 1.46-1.61 (3H, m), 1.78 (2H, m), 1.88-1.98 (1H, m), 3.03 (1H, dd, J=9.2, 14.3Hz), 3.18 (1H, dd,

J=10.2, 12.5Hz), 3.25-3.27 (2H, m), 3.34 (2H, m), 3.64-3.69 (1H, m), 4.94 (1H, dd, J=3.1, 10.1Hz), 7.09-7.16 (2H, m), 7.26 (1H, m), 7.31-7.39 (3H, m), 7.47 (1H, m), 7.68 (1H, m).

Examples 92 to 95

5 The compounds of Examples 92 to 95 are obtained in a similar manner to Example 91.

Example 92

(1R)-1-(3-Chlorophenyl)-2-(((1R)-2-(7-(((2R,6S)-2,6-dimethylpiperidin-1-yl)carbonyl)-1H-indol-3-yl)-1-methylethyl)amino)ethanol ·

10 trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.31 (9H, m), 1.46-1.61 (3H, m), 1.78 (2H, m), 1.88-1.98 (1H, m), 3.03 (1H, dd, J=9.2, 14.3Hz), 3.18 (1H, dd, J=10.3, 12.6Hz), 3.25-3.27 (2H, m), 3.34 (2H, m), 3.64-3.69 (1H, m), 4.94 (1H, dd, J=3.1, 10.1Hz), 7.09-7.18 (2H, m), 7.26 (1H, s), 7.31-7.39 (3H, m), 7.47 (1H, s), 7.68 (1H, dd, J=1.1, 7.7Hz).

Example 93

3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-N,N-diisopropyl-1H-indole-7-carboxamide · trifluoroacetate

20 ¹H-NMR (CDCl₃) δ: 1.12 (3H, d, J=6.2Hz), 1.40 (9H, m), 2.62 (1H, dd, J=9.2, 12.1Hz), 2.83 (2H, m), 2.87 (1H, dd, J=3.6, 12.1Hz), 3.00-3.06 (1H, m), 3.97 (2H, m), 4.50 (1H, dd, J=3.5, 9.1Hz), 7.05-7.24 (6H, m), 7.35 (1H, s), 7.61 (1H, d, J=7.7Hz), 8.78 (1H, brs).

Example 94

(1R)-1-(3-Chlorophenyl)-2-(((1R)-2-(7-(((2R,5R)-2,5-dimethylpyrrolidin-1-yl)carbonyl)-1H-indol-3-yl)-1-methylethyl)amino)ethanol ·

25 trifluoroacetate

¹H-NMR (CD₃OD) δ: 0.68 (3H, d, J=6.2Hz), 1.31 (3H, d,

J=6.5Hz), 1.36 (3H, d, J=6.2Hz), 1.57-1.69 (2H, m), 2.17-2.39 (2H, m), 3.04 (1H, dd, J=9.0, 14.3Hz), 3.17 (1H, dd, J=10.2, 12.6Hz), 3.24-3.34 (2H, m), 3.64-3.69 (1H, m), 4.24-4.28 (1H, m), 4.49-4.52 (1H, m), 4.93 (1H, dd, J=3.2, 10.1Hz), 7.13 (1H, dd, J=7.8, 8.4Hz), 7.24-7.27 (2H, m), 7.31-7.40 (3H, m), 7.46-7.47 (1H, m), 7.71 (1H, dd, J=0.8, 7.9Hz).

Example 95

(1R)-1-(3-Chlorophenyl)-2-[[[(1R)-2-{7-[(2,5-dimethylpyrrolidin-1-yl)-carbonyl]-1H-indol-3-yl}-1-methylethyl)amino]ethanol · trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.21 (6H, m), 1.30 (3H, d, J=6.6Hz), 1.73 (2H, m), 2.11 (2H, m), 3.02 (1H, dd, J=9.2, 14.3Hz), 3.17 (1H, dd, J=10.2, 12.6Hz), 3.25-3.34 (2H, m), 3.63-3.68 (1H, m), 4.10 (2H, m), 4.95 (1H, dd, J=3.1, 10.1Hz), 7.13 (1H, dd, J=7.3, 7.6Hz), 7.18 (1H, dd, J=1.2, 7.2Hz), 7.26 (1H, s), 7.31-7.39 (3H, m), 7.47-7.48 (1H, m), 7.70 (1H, dd, J=1.2, 7.7Hz).

Reference Example 49

tert-Butyl 3-((2R)-2-((5R)-5-(3-chlorophenyl)-2-oxo-1,3-oxazolidin-3-yl)-propyl)-1H-indol-7-ylcarbamate

To a solution of 3-((2R)-2-((5R)-5-(3-chlorophenyl)-2-oxo-1,3-oxazolidin-3-yl)propyl)-1H-indole-7-carboxylic acid (504 mg, 1.26 mmol) in toluene (25 mL) and t-butyl alcohol (12.5 mL) are added diphenylphosphoryl azide (379 μL, 1.76 mmol) and triethylamine (245 μL, 1.76 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for 30 minutes, and further stirred at 100°C for 5 hours. The reaction solution is poured into a saturated aqueous ammonium chloride solution, and the mixture is extracted with ethyl acetate. The organic layer is washed with a saturated brine, and dried over anhydrous sodium sulfate. The solvent is evaporated and the obtained

crude product is purified by silica gel column chromatography (n-hexane:ethyl acetate = 2:1 → 1:1) to give the title compound (427 mg, 72 %).

¹H-NMR (CDCl₃) δ: 1.30 (3H, d, J=6.8Hz), 1.56 (9H, s), 2.96 (2H, d, J=7.0Hz), 3.22 (1H, dd, J=8.6, 6.8Hz), 3.80 (1H, t, J=8.8Hz), 4.37-4.36 (1H, m), 5.33 (1H, dd, J=8.9, 6.8Hz), 6.68 (1H, brd, J=6.8Hz), 6.74 (1H, brs), 6.87 (1H, d, J=7.6Hz), 6.96 (1H, dd, J=7.9, 7.6Hz), 7.02 (1H, d, J=2.3Hz), 7.10-7.17 (2H, m), 7.22-7.24 (1H, m), 7.35 (1H, d, J=7.9Hz), 9.78 (1H, brs).

Reference Example 50

(5R)-3-((1R)-2-(7-Amino-1H-indol-3-yl)-1-methylethyl)-5-(3-chlorophenyl)-1,3-oxazolidin-2-one

To a solution of tert-butyl 3-((2R)-2-((5R)-5-(3-chlorophenyl)-2-oxo-1,3-oxazolidin-3-yl)propyl)-1H-indol-7-ylcarbamate (419.2 mg, 0.894 mmol) in methanol (2 mL) is added a 10 % solution of hydrochloric acid in methanol (10 mL) under nitrogen atmosphere, and the mixture is stirred at room temperature for 21 hours. The solvent is evaporated, and the residue is separated into a saturated ammonia solution in chloroform and water, and the organic layer is washed with a saturated brine, and dried over anhydrous sodium sulfate. The solvent is evaporated, and the obtained crude product is purified by silica gel column chromatography (a saturated ammonia solution in chloroform-methanol = 1:0 → 20:1) to give the title compound (299 mg, 90 %).

¹H-NMR (CDCl₃) δ: 1.33 (3H, d, J=6.8Hz), 2.93 (2H, d, J=7.6Hz), 3.24 (1H, dd, J=8.6, 6.8Hz), 3.81 (1H, t, J=8.8Hz), 4.42-4.51 (1H, m), 5.31 (1H, dd, J=9.0, 6.8Hz), 6.55 (1H, dd, J=7.3, 0.6Hz), 6.71 (1H, d,

J=7.8Hz), 6.91 (1H, t, J=7.9, 7.5Hz), 7.00 (1H, d, J=2.4Hz), 7.02-7.06 (2H, m), 7.16-7.19 (1H, m), 8.32 (1H, s).

Reference Example 51

N-(3-((2R)-2-((5R)-5-(3-Chlorophenyl)-2-oxo-1,3-oxazolidin-3-yl)propyl)-1H-indol-7-yl)propanamide

To a solution of (5R)-3-((1R)-2-(7-amino-1H-indol-3-yl)-1-methylethyl)-5-(3-chlorophenyl)-1,3-oxazolidin-2-one (28.4 mg, 0.0768 mmol) in tetrahydrofuran (1 mL) are added triethylamine (13.5 μ L, 0.0969 mmol) and propionyl chloride (8 μ L, 0.0921 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for 1 hour. The reaction solution is poured into water, and the mixture is extracted with ethyl acetate. The organic layer is washed with a saturated brine, and dried over anhydrous sodium sulfate. The solvent is evaporated, and the obtained crude product is purified by preparative thin layer chromatography (n-hexane:ethyl acetate = 1:1) to give the title compound (34.3 mg, yield: 100 %).

$^1\text{H-NMR}$ (CDCl_3) δ : 1.21 (3H, t, J=7.5Hz), 1.23 (3H, d, J=6.8Hz), 2.40 (2H, q, J=7.5Hz), 2.86 (2H, d, J=7.4Hz), 3.19 (1H, dd, J=8.5, 7.0Hz), 3.74 (1H, t, J=8.8Hz), 4.29-4.38 (1H, m), 5.26 (1H, dd, J=8.8, 7.0Hz), 6.78-6.82 (2H, m), 6.86-6.93 (2H, m), 7.01-7.02 (1H, m), 7.08 (1H, dd, J=7.9, 7.7Hz), 7.14-7.17 (1H, m), 7.31 (1H, d, J=7.7Hz), 7.72 (1H, s), 9.74 (1H, s).

Example 96

N-(3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-1H-indol-7-yl)propanamide

To a solution of N-(3-((2R)-2-((5R)-5-(3-chlorophenyl)-2-oxo-1,3-oxazolidin-3-yl)propyl)-1H-indol-7-yl)propanamide (28.4 mg, 0.0666

mmol) in ethanol (1 mL) is added a 5N aqueous sodium hydroxide solution (0.2 mL), and the mixture is stirred at room temperature for 15 hours, and further stirred at 60°C for 11 hours. The ethanol is evaporated, and the resulting residue is separated into a saturated

5 aqueous sodium hydrogen carbonate solution and chloroform. The organic layer is washed with a saturated brine, and dried over anhydrous sodium sulfate. The solvent is evaporated, and the obtained crude product is purified by preparative HPLC (water-acetonitrile-trifluoroacetic acid), and preparative thin layer chromatography (1st; a saturated ammonia solution in chloroform-methanol = 10:1, 2nd; chloroform:methanol = 10:1) to give the title compound (3.3 mg, 12.5 %).

¹H-NMR (CDCl₃) δ: 1.09 (3H, d, J=6.2Hz), 1.24 (3H, t, J=7.6Hz), 2.45 (2H, q, J=7.6Hz), 2.58 (1H, dd, J=12.2, 9.3Hz), 2.79-2.83 (3H, m), 2.98-3.04 (1H, m), 4.48 (1H, dd, J=9.3, 3.4Hz), 6.70 (1H, d, J=7.4Hz), 6.94 (1H, t, J=7.7Hz), 6.99 (1H, d, J=2.1Hz), 7.07-7.16 (3H, m), 7.24 (1H, s), 7.35 (1H, d, J=7.8Hz), 7.48 (1H, brs), 9.78 (1H, brs).

Reference Example 52

Tetrahydro-4H-pyran-4,4-dicarboxylic acid

Diethyl tetrahydro-2H-pyran-4,4-dicarboxylate (4.04 g, 20 mmol) is suspended in a 30 % aqueous sodium hydroxide solution (10 mL), and the mixture is stirred at room temperature for 28 hours. The pH value of the reaction solution is adjusted to pH 1 with conc. hydrochloric acid, and the mixture is separated into water and ethyl acetate. The organic layer is dried over anhydrous sodium sulfate, and the solvent is evaporated. Ethyl acetate (30mL) is added to the obtained crude product, and the mixture is washed by repulping, and the resulting solid is collected by filtration, and dried to give tetrahydro-

4H-pyran-4,4-dicarboxylic acid (3.19 g, 18.3 mmol, 92 %).

¹H-NMR (DMSO-d₆) δ: 1.90 (4H, t, J=5.3Hz), 3.55 (4H, t, J=5.3Hz), 12.93 (2H, s).

Reference Example 53

5 Tetrahydro-2H-pyran-4-carboxylic acid

Tetrahydro-4H-pyran-4,4-dicarboxylic acid (1.01 g, 5.80 mmol) is stirred at 185°C for one hour. The resulting residue is crystallized from toluene (5 mL) to give tetrahydro-2H-pyran-4-carboxylic acid (480.6 mg, 3.69 mmol, 64 %).

10 ¹H-NMR (DMSO-d₆) δ: 1.48-1.58 (2H, m), 1.70-1.74 (2H, m), 2.42-2.48 (1H, m), 3.29-3.36 (2H, m), 3.78-3.83 (2H, m), 12.22 (1H, s).

Example 97

N-(3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-1H-indol-7-yl)cyclohexanecarboxamide · trifluoroacetate

15 A solution of (5R)-3-((1R)-2-(7-amino-1H-indol-3-yl)-1-methylethyl)-5-(3-chlorophenyl)-1,3-oxazolidin-2-one (30 mg, 0.08 mmol), cyclohexanecarboxylic acid (15 mg, 0.12 mmol), bis(2-oxo-3-oxazolidinyl)phosphinic chloride (31 mg, 0.12 mmol) and triethylamine (0.033 mL, 0.24 mmol) in tetrahydrofuran (2 mL) is stirred at room temperature for
20 2.5 hours. The reaction solution is poured into a saturated aqueous sodium hydrogen carbonate solution, and the mixture is extracted with ethyl acetate. The organic layer is dried over anhydrous sodium sulfate and concentrated. The residue is purified by silica gel column chromatography (chloroform:methanol = 20:1). A solution of the
25 purified residue and potassium hydroxide (0.5 g) in ethanol (2 mL)/water (1 mL) is stirred at 80°C for 3.5 hours. The reaction solution is acidified with hydrochloric acid, and purified by preparative

HPLC (trifluoroacetic acid-acetonitrile-water) to give the title compound (29 mg, yield: 65 %).

¹H-NMR (CD₃OD) δ: 1.28-1.43 (3H, m), 1.31 (3H, d, J=6.5Hz), 1.57-1.61 (2H, m), 1.73-1.76 (1H, m), 1.85-1.89 (2H, m), 1.96-2.02 (2H, m), 2.47-2.51 (1H, m), 3.02 (1H, dd, J=9.0, 14.3Hz), 3.16 (1H, dd, J=10.1, 12.6Hz), 3.12-3.27 (2H, m), 3.62-3.68 (1H, m), 4.92 (1H, dd, J=3.2, 9.9Hz), 7.04 (1H, dd, J=7.7, 7.7Hz), 7.15 (1H, dd, J=0.6, 7.5Hz), 7.21 (1H, s), 7.30-7.38 (3H, m), 7.44-7.46 (2H, m).

Example 98

N-(3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-1H-indol-7-yl)tetrahydro-2H-pyran-4-carboxamide · trifluoroacetate

The title compound is obtained in a similar manner to Example 97.

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 1.85-1.98 (4H, m), 2.74-2.80 (1H, m), 3.02 (1H, dd, J=8.9, 14.3Hz), 3.16-3.28 (3H, m), 3.49-3.55 (2H, m), 3.63-3.68 (1H, m), 4.01-4.06 (2H, m), 4.93 (1H, m), 7.05 (1H, dd, J=7.7, 7.7Hz), 7.14 (1H, d, J=0.8, 7.5Hz), 7.30-7.38 (3H, m), 7.45 (1H, m), 7.47 (1H, dd, J=0.8, 7.9Hz).

Example 99

N-(3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-1H-indol-7-yl)-1-methylpiperazine-4-carboxamide · trifluoroacetate

A solution of (5R)-3-((1R)-2-(7-amino-1H-indol-3-yl)-1-methylethyl)-5-(3-chlorophenyl)-1,3-oxazolidin-2-one (30 mg, 0.08 mmol), N-methylpiperidine-4-carboxylic acid (34 mg, 0.24 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (46 mg, 0.24 mmol), 1-hydroxybenzotriazole (32 mg, 0.24 mmol) and triethylamine (67 μL, 0.48 mmol) in tetrahydrofuran (2 mL) is stirred at room temperature for

16 hours. The mixture is diluted with ethyl acetate, washed with a saturated aqueous sodium hydrogen carbonate solution, dried over sodium sulfate, and concentrated. The residue is purified by silica gel column chromatography (a saturated ammonia solution in chloroform: methanol = 100:1 → 20:1). A solution of the purified residue and potassium hydroxide (0.5 g) in ethanol (2 mL)/water (1 mL) is stirred at 80°C for 3.5 hours. The reaction solution is acidified with hydrochloric acid, and purified by preparative HPLC (trifluoroacetic acid-acetonitrile-water) and preparative thin layer chromatography (silica gel, a saturated ammonia solution in chloroform:methanol = 10:1) to give the title compound (5 mg, yield: 13 %).

¹H-NMR (CDCl₃) δ: 1.12 (3H, d, J=6.2Hz), 1.90-2.07 (6H, m), 2.32 (3H, s), 2.36 (1H, m), 2.61 (1H, dd, J=9.2, 12.1Hz), 2.81-2.88 (3H, m), 2.96-3.05 (3H, m), 4.49 (1H, dd, J=3.6, 9.2Hz), 6.75 (1H, d, J=7.4Hz), 7.01 (1H, dd, J=7.8, 7.8Hz), 7.05 (1H, d, J=2.1Hz), 7.15-7.18 (1H, m), 7.19-7.23 (2H, m), 7.33 (1H, s), 7.42 (1H, d, J=7.9Hz), 7.57 (1H, s), 9.88 (1H, brs).

Reference Example 54

Methyl 3-[(2R)-2-[(2R)-2-{4-(benzyloxy)-3-[(methylsulfonyl)amino]phenyl}-2-[(triethylsilyl)oxy]ethyl]amino)propyl]-1H-indole-7-carboxylate

A solution of (R)-N-[2-benzyloxy-5-(2-iodo-1-triethylsilyloxy-ethyl)phenyl]methanesulfonamide (2.09 g, 3.72 mmol), methyl (R)-3-(2-aminopropyl)-1H-indole-7-carboxylate (1.00 g, 3.72 mmol), diisopropylethylamine (1.9 mL, 11.2 mmol) in tetrahydrofuran (30 mL)-acetonitrile (20 mL) is stirred at 110°C for 18 hours in a sealed vessel. After cooling, to the reaction solution are added a saturated aqueous sodium hydrogen carbonate solution and a saturated brine, and the mixture is

extracted with ethyl acetate. The organic layer is dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue is purified by silica gel column chromatography (chloroform/methanol = 100/1 → 100/4) to give the title compound (1.05 g, yield: 42 %).

¹H-NMR (CDCl₃) δ: 0.42 (6H, m), 0.73 (12H, m), 2.83 (3H, s), 3.00 (5H, m), 3.93 (3H, s), 5.05 (2H, s), 5.06 (1H, m), 6.7-7.5 (11H, m), 7.77 (1H, d, J=7.7Hz), 7.86 (1H, d, J=7.4Hz), 9.62 (1H, br).

Reference Example 55

3-[(2R)-2-[(2R)-2-{4-(Benzyloxy)-3-[(methylsulfonyl)amino]phenyl}-2-hydroxyethyl)amino]propyl]-1H-indole-7-carboxylic acid

To a solution of methyl 3-[(2R)-2-[(2R)-2-{4-(benzyloxy)-3-[(methylsulfonyl)amino]phenyl}-2-[(triethylsilyl)oxy]ethyl)amino]propyl]-1H-indole-7-carboxylate (400 mg, 0.601 mmol) in methanol (20 ml) is added a solution of potassium hydroxide (336 mg, 6.01 mmol) in water (2.0 ml), and the mixture is stirred at 40°C for 9 hours. The reaction solution is neutralized with 1N hydrochloric acid at 0°C, and thereto is added a pH standard solution (pH 6.8, 5.0 ml). The resultant is purified by reversed phase column (octadecylsilyl, water/methanol = 95/5 → 40/60) to give the title compound (296 mg, yield: 92 %).

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 1.51 (2H, br), 1.71 (4H, brs), 3.03 (1H, dd, J=9.1, 14.2Hz), 3.17 (1H, dd, J=10.1, 12.5Hz), 3.25 (2H, m), 3.65 (1H, m), 3.70 (2H, br), 4.95 (1H, dd, J=3.2, 10.1Hz), 7.15 (2H, m), 7.27 (1H, s), 7.34 (3H, m), 7.48 (1H, s), 7.71 (1H, dd, J=2.2, 6.9Hz).

Example 100

N-{2-Hydroxy-5-[(1R)-1-hydroxy-2-[(1R)-1-methyl-2-[7-(pyrrolidin-1-yl-

carbonyl)-1H-indol-3-yl]ethyl)amino)ethyl]phenyl]methanesulfonamide trifluoroacetate

To a solution of 3-((2R)-2-(((2R)-2-(4-(benzyloxy)-3-((methylsulfonyl)amino]phenyl)-2-hydroxyethyl)amino]propyl)-1H-indole-7-carboxylic acid (40.0 mg, 0.0744 mmol) in dimethylformamide (1.5 ml) are added triethylamine (124 μ l, 1.22 mmol), pyrrolidine (62.1 μ l, 0.744 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (143 mg, 0.744 mmol) and 1-hydroxybenzotriazole (101 mg, 0.744 mmol), and the mixture is stirred at room temperature for 16 hours. To the reaction solution are added a saturated aqueous sodium hydrogen carbonate solution and a saturated brine, and the mixture is extracted with chloroform. The organic layers are combined, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue is purified by preparative thin layer chromatography (a 15 % methanol solution in chloroform). The obtained compound is dissolved in methanol (2.0 ml), and thereto is added a 10 % palladium on carbon (50 % wet, 20 mg), and the mixture is stirred under hydrogen atmosphere for 30 minutes. The mixture is filtered through celite, and the filtrate is concentrated under reduced pressure. To a solution of the residue in acetonitrile is added trifluoroacetic acid (15 ml), and the mixture is purified by preparative reversed phase HPLC (trifluoroacetic acid/water/acetonitrile) to give the title compound (25.0 mg, yield: 55 %).

$^1\text{H-NMR}$ (CD_3OD) δ : 1.29 (3H, d, $J=6.6\text{Hz}$), 1.88 (2H, m), 2.01 (2H, m), 2.93 (3H, s), 3.02 (1H, dd, $J=9.0, 14.3\text{Hz}$), 3.20 (2H, m), 3.27 (1H, m), 3.43 (2H, m), 3.64 (3H, m), 4.83 (1H, m), 6.89 (1H, d, $J=8.3\text{Hz}$), 7.1-7.2 (2H, m), 7.25 (2H, m), 7.39 (1H, d, $J=2.1\text{Hz}$), 7.71 (1H, dd, $J=0.9,$

7.9Hz).

Examples 101 to 104

The compounds of Examples 101 to 127 are obtained in a similar manner to Example 100.

5 Example 101

N-{2-Hydroxy-5-[(1R)-1-hydroxy-2-(((1R)-1-methyl-2-[7-(piperidin-1-yl-carbonyl)-1H-indol-3-yl]ethyl)amino)ethyl]phenyl}methanesulfonamide · trifluoroacetate

10 ¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.6Hz), 1.5-1.7 (6H, m), 2.92 (3H, s), 3.02 (1H, dd, J=9.0, 14.3Hz), 3.20 (2H, m), 3.27 (1H, m), 3.40 (2H, brs), 3.65 (1H, m), 3.70 (2H, brs), 4.83 (1H, m), 6.89 (1H, d, J=8.3Hz), 7.1-7.2 (3H, m), 7.26 (1H, s), 7.39 (1H, d, J=2.1Hz), 7.70 (1H, dd, J=2.2, 6.8Hz).

Example 102

15 N-{2-Hydroxy-5-[(1R)-1-hydroxy-2-(((1R)-1-methyl-2-[7-(morpholin-4-yl-carbonyl)-1H-indol-3-yl]ethyl)amino)ethyl]phenyl}methanesulfonamide · trifluoroacetate

20 ¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.6Hz), 2.93 (3H, s), 3.02 (1H, dd, J=9.0, 14.3Hz), 3.20 (2H, m), 3.27 (1H, m), 3.65 (9H, m), 4.83 (1H, m), 6.89 (1H, d, J=6.9Hz), 7.1-7.2 (3H, m), 7.28 (1H, s), 7.39 (1H, d, J=2.1Hz), 7.71 (1H, dd, J=1.2, 7.7Hz).

Example 103

25 N-(2-Hydroxy-5-[(1R)-1-hydroxy-2-(((1R)-1-methyl-2-{7-[(4-methyl-piperazin-1-yl)carbonyl]-1H-indol-3-yl]ethyl)amino)ethyl]phenyl)-methanesulfonamide · 2 trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.5Hz), 2.93 (3H, s), 2.93 (3H, s), 3.02 (1H, dd, J=9.1, 14.3Hz), 3.20 (2H, m), 3.27 (1H, m), 3.45

(4H, m), 3.65 (2H, m), 4.83 (1H, m), 6.89 (1H, d, J=8.3Hz), 7.11 (1H, dd, J=2.2, 8.4Hz), 7.18 (1H, t, J=7.8Hz), 7.27 (1H, m), 7.31 (1H, s), 7.37 (1H, d, J=2.1Hz), 7.77 (1H, dd, J=1.0, 8.8Hz).

Example 104

5 3-((2R)-2-(((2R)-2-Hydroxy-2-{4-hydroxy-3-[(methylsulfonyl)amino]-phenyl}ethyl)amino)propyl)-N,N-dimethyl-1H-indole-7-carboxamide trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.6Hz), 2.92 (3H, s), 3.02 (4H, m), 3.20 (5H, m), 3.25 (1H, m), 3.40 (2H, brs), 3.65 (1H, m), 4.84 (1H, dd, J=4.8, 8.3Hz), 6.89 (1H, d, J=8.3Hz), 7.1-7.2 (3H, m), 7.27 (1H, s), 7.38 (1H, d, J=2.1Hz), 7.70 (1H, dd, J=1.2, 7.8Hz).

Reference Example 56

N-(3-((1R)-2-Iodo-1-[(triethylsilyl)oxy]ethyl)phenyl)methanesulfonamide

15 The title compound is obtained from N-[3-(2-bromoacetyl)-phenyl]methanesulfonamide in a similar manner to Reference Example 5, Reference Example 6 and Reference Example 7.

¹H-NMR (CDCl₃) δ: 0.54-0.61 (6H, m), 0.91 (9H, t, J=7.7Hz), 3.01 (3H, s), 3.33 (2H, d, J=6.1Hz), 4.75 (1H, t, J=5.8Hz), 7.15-7.18 (2H, m), 7.22-7.23 (1H, m), 7.33 (1H, dd, J=7.9, 7.9Hz).

20 Example 105

N,N-Diethyl-3-((2R)-2-(((2R)-2-hydroxy-2-{3-[(methylsulfonyl)amino]-phenyl}ethyl)amino)propyl)-1H-indole-7-carboxamide

25 The title compound is obtained from N-(3-((1R)-2-iodo-1-[(triethylsilyl)oxy]ethyl)phenyl)methanesulfonamide and (R)-3-(2-amino-propyl)-1H-indole-7-carboxylic acid diethylamide in a similar manner to Reference Example 8.

¹H-NMR (CDCl₃) δ: 1.15 (3H, d, J=6.1Hz), 1.26 (6H, t, J=7.0Hz),

2.59 (1H, dd, J=5.0, 11.8Hz), 2.69-2.76 (2H, m), 2.82-2.87 (1H, m), 2.86 (3H, s), 2.95-3.03 (1H, m), 3.53 (4H, m), 4.45 (1H, dd, J=5.0, 7.9Hz), 6.85 (1H, m), 6.99 (1H, m), 7.03-7.07 (3H, m), 7.14-7.19 (2H, m), 7.58 (1H, dd, J=0.6, 7.9Hz), 9.00 (1H, brs).

5 Example 106

4-[(1R,2S)-1-Hydroxy-2-({2-[7-(pyrrolidin-1-ylcarbonyl)-1H-indol-3-yl]-ethyl}amino)propyl]phenol · trifluoroacetate

10 The title compound is obtained from 3-{2-[(1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-(methyl)ethylamino]ethyl}-1H-indole-7-carboxylic acid in a similar manner to Example 11.

¹H-NMR (CD₃OD) δ: 1.08 (3H, d, J=6.5Hz), 1.88 (2H, m), 2.01 (2H, m), 3.22 (2H, m), 3.41 (3H, m), 3.68 (2H, t, J=6.9Hz), 4.99 (1H, d, J=3.1Hz), 6.76 (2H, m), 7.14 (3H, m), 7.30 (2H, m), 7.72 (1H, dd, J=0.9, 7.9Hz).

15 Example 107

3-((2R)-2-{[(2R)-2-(4-Chlorophenyl)-2-hydroxyethyl]amino}propyl)-N,N-diethyl-1H-indole-7-carboxamide

The title compound is obtained from (R)-4-chlorostyrene oxide in a similar manner to Example 9.

20 ¹H-NMR (CDCl₃) δ: 1.12 (3H, d, J=6.2Hz), 1.27 (6H, t, J=7.1Hz), 2.61 (1H, dd, J=9.2, 12.1Hz), 2.83-2.87 (m, 3H), 3.00-3.05 (1H, m), 3.55 (4H, q, J=6.4Hz), 4.51 (1H, dd, J=3.6, 9.1Hz), 7.04-7.10 (2H, m), 7.21-7.31 (5H, m), 7.63 (1H, d, J=7.9Hz), 8.91 (1H, brs).

Reference Example 57

25 N,N-Diethyl-2-(2-nitrophenyl)acetamide

The title compound is obtained from 2-nitrophenylacetic acid in a similar manner to Reference Example 14.

¹H-NMR (CDCl₃) δ: 1.14 (3H, t, J=7.1Hz), 1.30 (3H, t, J=7.2Hz), 3.38-3.46 (4H, m), 4.05 (2H, s), 7.34 (1H, dd, J=1.1, 7.6Hz), 7.42-7.46 (1H, m), 7.55-7.59 (1H, m), 8.09 (1H, dd, J=1.3, 8.2Hz).

Reference Example 58

5 N,N-Diethyl-2-(1H-indol-7-yl)acetamide

N,N-Diethyl-2-(2-nitrophenyl)acetamide (87.2 g, 0.369 mol) is dissolved in tetrahydrofuran (1.5L), and the mixture is cooled to -70°C. To the mixture is added dropwise a 1M solution of vinylmagnesium bromide in tetrahydrofuran (1.11 L, 1.11 mol). After the addition, the mixture is stirred at -50°C for additional 5 hours. The reaction solution is poured into an aqueous ammonium chloride solution, and the mixture is extracted with ethyl acetate. The organic layer is washed with a saturated brine, dried over anhydrous magnesium sulfate, and concentrated. The resulting residue is purified by silica gel column chromatography (chloroform, hexane/ethyl acetate = 4/1) to give the title compound (4.02 g, yield: 5 %).

¹H-NMR (CDCl₃) δ: 1.05 (3H, t, J=7.1Hz), 1.13 (3H, t, J=7.2Hz), 3.33 (2H, q, J=7.1Hz), 3.42 (2H, q, J=7.2Hz), 3.94 (2H, s), 6.52-6.53 (1H, m), 6.96 (1H, d, J=7.0Hz), 7.04 (1H, dd, J=7.4, 7.6Hz), 7.23-7.26 (2H, m), 7.57 (1H, d, J=7.8Hz), 10.05 (1H, brs).

Reference Example 59

2-{3-[(2R)-2-Aminopropyl]-1H-indol-7-yl}-N,N-diethylacetamide

The title compound is obtained from N,N-diethyl-2-(1H-indol-7-yl)acetamide in a similar manner to Reference Example 4.

¹H-NMR (CDCl₃) δ: 1.06 (3H, t, J=7.1Hz), 1.15 (3H, t, J=7.2Hz), 1.16 (3H, d, J=6.3Hz), 2.64 (1H, dd, J=8.2, 14.0Hz), 2.87 (1H, dd, J=4.4, 14.2Hz), 3.24-3.29 (1H, m), 3.34 (2H, q, J=7.1Hz), 3.43 (2H, q, J=7.1Hz),

3.92 (2H, s), 6.97 (1H, d, J=7.2Hz), 7.04 (1H, dd, J=7.1, 7.8Hz), 7.07 (1H, d, J=2.3Hz), 7.53 (1H, d, J=7.9Hz), 9.88 (1H, brs).

Example 108

2-[3-((2R)-2-(((2R)-2-(3-Chlorophenyl)-2-hydroxyethyl)amino)propyl)-1H-indol-7-yl]-N,N-diethylacetamide

To a solution of 2-{3-[(2R)-2-aminopropyl]-1H-indol-7-yl}-N,N-diethylacetamide (1.0 g, 3.48 mmol) in dimethylsulfoxide (5 mL) is added N-trimethylsilylacetamide (0.50 g, 3.83 mmol), and the mixture is stirred at room temperature for 1 hour. To the reaction solution is added (R)-(+)-3-chlorostyrene oxide (0.89 mL, 6.96 mmol), and the mixture is stirred at 110°C for 2 hours. After cooling, a 2 N hydrochloric acid (5 mL) is added to the mixture, and the mixture is stirred at room temperature for 0.5 hour. To the mixture are added a saturated aqueous sodium hydrogen carbonate solution and a saturated brine, and the mixture is extracted with chloroform. The organic layer is dried over anhydrous sodium sulfate and concentrated. The residue is purified by silica gel column chromatography (a saturated ammonia solution in chloroform) to give the title compound (0.90 g, yield: 59 %).

¹H-NMR (CDCl₃) δ: 1.06 (3H, t, J=7.1Hz), 1.12 (3H, d, J=6.2Hz), 1.14 (3H, t, J=7.2Hz), 2.62 (1H, dd, J=9.2, 12.1Hz), 2.82-2.90 (3H, m), 3.01-3.06 (1H, m), 3.34 (2H, q, J=7.1Hz), 3.43 (2H, q, J=7.2Hz), 3.92 (2H, s), 4.50 (1H, dd, J=3.6, 9.1Hz), 6.97 (1H, d, J=7.0Hz), 7.03-7.07 (2H, m), 7.16-7.25 (3H, m), 7.33-7.34 (1H, m), 7.51 (1H, d, J=7.9Hz), 9.90 (1H, brs).

Reference Example 60

(1R)-2-(7-(Benzyloxy)-1H-indol-3-yl)-1-methylethylamine

The title compound is obtained in a similar manner to

Reference Example 4.

¹H-NMR (CDCl₃) δ: 1.16 (3H, d, J=6.3Hz), 2.64 (1H, dd, J=14.1, 8.2Hz), 2.86 (1H, d, J=14.1, 4.9Hz), 3.24-3.32 (1H, m), 6.73 (1H, d, J=7.7Hz), 7.01-7.04 (2H, m), 7.24 (1H, d, J=8Hz), 7.34-7.43 (3H, m), 7.47-7.49 (2H, m), 8.30 (1H, s).

Reference Example 61

(2R)-N-((1R)-2-(7-(Benzyloxy)-1H-indol-3-yl)-1-methylethyl)-2-hydroxy-2-pyridin-3-ylacetamide

To a solution of (1R)-2-(7-(benzyloxy)-1H-indol-3-yl)-1-methylethylamine (5.0 g, 17.8 mmol) in tetrahydrofuran (100 mL) are added successively with stirring (R)-2-hydroxy-2-(3-pyridyl)acetic acid · sulfate (4.68 g, 11.6 mmol), triethylamine (3.22 mL, 23.1 mmol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide · hydrochloride (4.78 g, 24.9 mmol), and 1-hydroxybenzotriazole (3.36 g, 24.9 mmol) under ice-cooling, and the mixture is stirred at room temperature for 2.5 hours. The reaction mixture is diluted with a mixture of a saturated aqueous sodium hydrogen carbonate solution and water (1:1, 100 mL), and extracted with ethyl acetate (2 x 100 mL). The organic layers are combined and washed with a mixture of a saturated brine and water (1:1, 100 mL), dried over magnesium sulfate, and separated by filtration. The filtrate is concentrated under reduced pressure, and the obtained pale brown solid is dispersed into chloroform (80 mL). The insoluble white solid is collected by filtration, washed with chloroform (40 mL), and dried under reduced pressure to give the title compound (6.38 g, yield: 86 %).

¹H-NMR (CDCl₃) δ: 1.20 (3H, d, J=6.6Hz), 2.85 (1H, dd, J=14.6, 5.5Hz), 2.91 (1H, dd, J=14.6, 6.1Hz), 4.32-4.39 (1H, m), 4.96 (1H, s), 5.20 (2H, s), 5.97 (1H, d, J=8.2Hz), 6.72 (1H, d, 7.8Hz), 6.73 (1H, s),

7.00 (1H, dd, J=8.0, 7.8Hz), 7.13 (1H, d, J=8.0Hz), 7.20 (1H, d, J=7.9, 4.8Hz), 7.38-7.44 (3H, m), 7.48-7.50 (2H, m), 7.59 (1H, ddd, J=7.9, 2.2, 1.6Hz), 8.34 (1H, brs), 8.53 (1H, dd, J=4.8, 1.6Hz), 8.57 (1H, d, J=2.2Hz).

5 Reference Example 62

(1R)-2-(((1R)-2-(7-(Benzyloxy)-1H-indol-3-yl)-1-methylethyl)amino)-1-pyridin-3-ylethanol

10 To a solution of (2R)-N-((1R)-2-(7-(benzyloxy)-1H-indol-3-yl)-1-methylethyl)-2-hydroxy-2-pyridin-3-ylacetamide (283 mg, 0.681 mmol) in tetrahydrofuran (15 mL) is added a 2M solution of boranedimethyl sulfide complex in tetrahydrofuran (2.04 mL, 4.09 mmol), and the mixture is refluxed for 3 hours. To the reaction solution is added a 10 % hydrogen chloride solution in methanol (30 mL), and the mixture is stirred for 30 minutes. The solvent is evaporated, and the residue is
15 separated into a saturated aqueous sodium hydrogen carbonate solution and chloroform. The organic layer is washed with a saturated brine, and dried over sodium sulfate. The solvent is distilled off, and the obtained crude product is purified by silica gel column chromatography (a saturated ammonia solution in chloroform - methanol =
20 100:1) to give the title compound (184.8 mg, yield: 68 %).

¹H-NMR (CDCl₃) δ: 1.13 (3H, d, J=6.2Hz), 2.65 (1H, dd, J=12.1, 9.2Hz), 2.77-2.88 (3H, m), 3.01-3.08 (1H, m), 4.54 (1H, dd, J=9.2, 3.6Hz), 5.19 (2H, s), 6.72 (1H, d, J=7.7Hz), 6.97 (1H, d, J=2.1Hz), 7.02 (1H, t, J=7.8Hz), 7.20-7.22 (2H, m), 7.33-7.41 (3H, m), 7.46-7.48 (2H,
25 m), 7.63-7.66 (1H, m), 8.45-8.50 (3H, m).

Reference Example 63

tert-Butyl (1R)-2-(7-(benzyloxy)-1H-indol-3-yl)-1-methylethyl-((2R)-2-

hydroxy-2-pyridin-3-ylethyl)carbamate

To a solution of (1R)-2-(((1R)-2-(7-(benzyloxy)-1H-indol-3-yl)-1-methylethyl)amino)-1-pyridin-3-ylethanol (182 mg, 0.453 mmol) in tetrahydrofuran (5 mL) is added di-tert-butyl bicarbonate (148 mg, 0.677 mmol), and the mixture is stirred at room temperature for one day. Without concentration, the reaction solution is purified by silica gel column chromatography (n-hexane → n-hexane:ethyl acetate = → 2:1 → 1:1 → 1:2) to give the title compound (205.4 mg, yield: 91 %).

¹H-NMR (CDCl₃) δ: 1.27 (3H, d, J=6.9Hz), 1.29 (9H, s), 2.79-2.85 (2H, m), 3.10 (1H, m), 3.59-3.65 (1H, m), 4.26-4.31 (1H, m), 4.69 (1H, m), 5.20 (2H, s), 5.55 (1H, s), 6.72 (1H, d, J=7.6Hz), 6.88 (1H, brs), 7.02 (1H, dd, J=8.0, 7.5Hz), 7.20 (1H, d, J=7.5Hz), 7.26-7.32 (1H, m), 7.32-7.41 (3H, m), 7.44-7.52 (2H, m), 7.74 (1H, m), 8.27 (1H, s), 8.51-8.53 (3H, m).

Reference Example 64

tert-Butyl (1R)-2-(7-(benzyloxy)-1H-indol-3-yl)-1-methylethyl((2R)-2-pyridin-3-yl-2-((triethylsilyl)oxy)ethyl)carbamate

To a solution of tert-butyl (1R)-2-(7-(benzyloxy)-1H-indol-3-yl)-1-methylethyl((2R)-2-hydroxy-2-pyridin-3-ylethyl)carbamate (8.05 g, 16.1 mmol) in N,N-dimethylformamide (60 mL) are added imidazole (4.37 g, 64.2 mmol) and triethylsilane chloride (5.44 mL, 32.1 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for 3 hours. The reaction solution is poured into water, and the mixture is extracted with ethyl acetate. The organic layer is washed with water and a saturated brine, and dried over anhydrous sodium sulfate. The solvent is evaporated, and the obtained crude product is purified by silica gel column chromatography (n-hexane:ethyl

acetate = 5:1 → 3:1) to give the title compound (7.94 g, yield: 80 %).

IR (ATR/FT-IR, cm^{-1}): 1682, 1578, 1500, 1454, 1404, 1365, 1330, 1257, 1238, 1164, 1083, 1025, 1003.

Reference Example 65

5 tert-Butyl (1R)-2-(7-hydroxy-1H-indol-3-yl)-1-methylethyl((2R)-2-pyridin-3-yl-2-((triethylsilyl)oxy)ethyl)carbamate

To a solution of tert-butyl (1R)-2-(7-(benzyloxy)-1H-indol-3-yl)-1-methylethyl((2R)-2-pyridin-3-yl-2-((triethylsilyl)oxy)ethyl)carbamate (2.5 g, 4.06 mmol) in methanol (50 mL) are added a phosphate standard
10 buffer (pH 6.86, 2.5 mL) and a 10 % palladium on carbon (50 % wet, 2.5g), and the mixture is stirred at room temperature for 1.5 hour under hydrogen atmosphere. The reaction solution is filtered through celite, and the filtrate is evaporated under reduced pressure to remove the solvent. To the residue are added water and ethyl acetate, and the
15 organic layer is washed with a saturated brine, and dried over anhydrous sodium sulfate. The solvent is evaporated, and the obtained crude product is purified by silica gel column chromatography (n-hexane:ethyl acetate = 3:1 → 3:2) to give the title compound (1.84 g, yield: 87 %).

20 IR (ATR/FT-IR, cm^{-1}): 3336, 1681, 1577, 1454, 1430, 1407, 1365, 1331, 1238, 1164, 1083, 1003.

Example 109

2-((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)propanoic acid

25 To a solution of tert-butyl (1R)-2-(7-hydroxy-1H-indol-3-yl)-1-methylethyl((2R)-2-pyridin-3-yl-2-((triethylsilyl)oxy)ethyl)carbamate (100 mg, 0.190 mmol) in acetone (10 mL) is added ethyl α -bromopropanoate

(51.6 mg, 0.285 mmol), potassium carbonate (31.5 mg, 0.228 mmol) and potassium iodide (5 mg), and the mixture is refluxed for 16 hours. The reaction solution is allowed to cool, filtered, and concentrated. The residue is purified by silica gel column chromatography (n-hexane/ethyl acetate = 10/1 → 1/1) to give ethyl 2-((3-((2R)-((tert-butoxycarbonyl)-((2R)-2-pyridin-3-yl-2-((triethyl)oxy)ethyl)amino)propyl)-1H-indol-7-yl)-oxy)propanoate (81.3 mg, yield: 68 %).

Further, to a solution of ethyl 2-((3-((2R)-((tert-butoxycarbonyl)-((2R)-2-pyridin-3-yl-2-((triethyl)oxy)ethyl)amino)propyl)-1H-indol-7-yl)-oxy)propanoate (81.3 mg, 0.130 mmol) in ethanol (1.0 mL) is added a 4N hydrochloric acid in dioxane (1.0 mL), and the mixture is stirred at room temperature for 4 hours. Water (1.0 mL) is added to the mixture, and the mixture is further stirred for 1 hour. The mixture is concentrated under reduced pressure, and the residue is dissolved in a mixture of ethanol (1.25 mL) and water (0.25 mL), and thereto is added KOH (72.8 mg, 1.30 mmol). The mixture is stirred at room temperature overnight, and neutralized with 1N hydrochloric acid and a pH buffer (pH 6.8). The mixture is evaporated under reduced pressure to remove ethanol, and thereto is added a 0.1 % aqueous trifluoroacetic acid solution (2.0 mL). The mixture is purified by reversed phase column (trade name; COSMOSIL 75C18-OPN (Nacalai Tesque), 0.1 % aqueous trifluoroacetic acid solution/methanol = 100/0 → 50/50) to give the title compound (80.1 mg, yield: 100 %).

¹H-NMR (CD₃OD) δ: 1.31 and 1.32 (total 3H, d, J=6.5Hz), 1.66 (3H, d, J=6.6Hz), 3.03 (1H, m), 3.31 (2H, m), 3.40 (1H, m), 3.65 (1H, m), 3.70 (2H, br), 4.96 (1H, m), 5.26 (1H, m), 6.56 (1H, d, J=7.7Hz), 6.94 (1H, t, J=7.9Hz), 7.17 (1H, s), 7.21 (1H, dd, J=1.4, 8.0Hz), 8.01 (1H, dd,

$J=5.7, 8.0\text{Hz}$), 8.53 (1H, d, $J=8.1\text{Hz}$), 8.78 (1H, d, $J=5.2\text{H}$), 8.88 (1H, m).

Example 110

N,N-Diethyl-2-((3-((2R)-2-(((2R)-2-hydroxy-2-pyridin-3-ylethyl)amino)-propyl)-1H-indol-7-yl)oxy)propanamide · 2 trifluoroacetate

5 To a solution of 2-((3-((2R)-2-(((2R)-2-hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)propanoic acid (18.8 mg, 0.0490 mmol) in N,N-dimethylformamide (1.5 mL) are added triethylamine (170 μL , 1.22 mmol), diethylamine (50.7 μL , 0.490 mmol), N,N-bis(2-oxo-3-oxazolidinyl)phosphinic chloride (125 mg, 0.490 mmol), and the mixture
10 is stirred at 0°C for 2 hours. To the reaction solution are added a saturated aqueous sodium hydrogen carbonate solution and a saturated brine, and the mixture is extracted with chloroform. The organic layers are combined, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The obtained residue is
15 dissolved in acetonitrile (2.0 mL). Trifluoroacetic acid (30 μL) is added to the mixture, and the mixture is purified by preparative reversed phase HPLC (octadecylsilyl, trifluoroacetic acid/water/acetonitrile) to give the title compound (10.2 mg, yield: 31 %).

20 $^1\text{H-NMR}$ (CD_3OD) δ : 1.11 (3H, t, $J=7.0\text{Hz}$), 1.17 (3H, m), 1.32 (3H, m), 1.60 (3H, m), 3.03 (1H, m), 3.29 (2H, m), 3.36 (2H, m), 3.39 (1H, m), 3.53 (2H, q, $J=6.4\text{Hz}$), 3.67 (1H, m), 5.19 (1H, m), 5.30 (1H, m), 6.59 (1H, dd, $J=2.6, 7.7\text{Hz}$), 6.94 (1H, dd, $J=7.8, 7.8\text{Hz}$), 7.18 (1H, s), 7.22 (1H, dd, $J=1.4, 7.9\text{Hz}$), 7.91 (1H, dd, $J=5.6, 8.0\text{Hz}$), 8.41 (1H, d, $J=8.1\text{Hz}$), 8.74 (1H, d, $J=5.1\text{Hz}$), 8.78 (1H, m).

25 Example 111

2-((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)-N,N-dimethylpropanamide · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 110.

¹H-NMR (CD₃OD) δ: 1.32 (3H, m), 1.60 (3H, m), 2.97 (3H, s), 3.03 (1H, m), 3.19 (3H, s), 3.27 (2H, m), 3.35 (1H, m), 3.69 (1H, m), 3.52 (1H, m), 5.38 (1H, m), 6.53 (1H, dd, J=2.5, 7.7Hz), 6.94 (1H, dd, J=7.8, 7.8Hz), 7.18 (1H, s), 7.21 (1H, dd, J=2.5, 8.0Hz), 7.98 (1H, m), 8.48 (1H, m), 8.78 (2H, m).

Example 112

(1R)-2-(((1R)-1-Methyl-2-(7-(1-methyl-2-oxo-2-piperidin-1-ylethoxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 11.

¹H-NMR (CD₃OD) δ: 1.31 and 1.32 (total 3H, d, J=6.5Hz), 1.50 (4H, m), 1.60 (3H, d, J=6.6Hz), 1.65 (2H, m), 3.03 (1H, m), 3.31 (2H, m), 3.38 (1H, m), 3.56 (2H, t, J=5.3Hz), 3.65 (2H, m), 3.70 (1H, m), 5.23 (1H, m), 5.36 (1H, m), 6.57 (1H, dd, J=2.6, 7.7Hz), 6.95 (1H, t, J=7.8Hz), 7.18 (1H, s), 7.22 (1H, dd, J=2.2, 8.0Hz), 8.01 (1H, dd, J=5.7, 8.0Hz), 8.53 (1H, d, J=8.2Hz), 8.80 (2H, m).

Example 113

(1R)-2-(((1R)-1-Methyl-2-(7-(1-methyl-2-morpholin-4-yl-2-oxoethoxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 11.

¹H-NMR (CD₃OD) δ: 1.31 and 1.32 (total 3H, d, J=6.5Hz), 1.62 (3H, d, J=6.6Hz), 3.03 (1H, m), 3.31 (1H, m), 3.40 (2H, m), 3.58 (6H, m), 3.70 (3H, m), 5.25 (1H, m), 5.34 (1H, m), 6.60 (1H, dd, J=1.7, 7.7Hz), 6.96 (1H, t, J=7.9Hz), 7.19 (1H, s), 7.23 (1H, dd, J=0.9, 7.9Hz), 8.01 (1H,

dd, J=5.7, 8.0Hz), 8.55 (1H, d, J=8.1Hz), 8.80 (1H, d, J=5.3Hz), 8.86 (1H, m).

Example 114

2-((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)-2-methylpropanoic acid

The title compound is obtained in a similar manner to Example 109.

¹H-NMR (CD₃OD) δ: 1.32 (3H, d, J=6.5Hz), 1.63 (6H, s), 3.00 (1H, dd, J=9.1, 14.2Hz), 3.31 (2H, m), 3.41 (1H, dd, J=4.3, 12.8Hz), 3.65 (1H, m), 5.25 (1H, dd, J=3.0, 9.7Hz), 6.63 (1H, d, J=7.7Hz), 6.92 (1H, t, J=7.9Hz), 7.17 (1H, s), 7.26 (1H, d, J=7.9Hz), 8.01 (1H, dd, J=5.7, 8.1Hz), 8.53 (1H, d, J=8.1Hz), 8.80 (1H, d, J=5.4Hz), 8.89 (1H, brs).

Example 115

2-((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)-N,N,2-trimethylpropanamide · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 110.

¹H-NMR (CD₃OD) δ: 1.32 (3H, d, J=6.5Hz), 1.68 (6H, s), 2.95 (3H, s), 2.99 (1H, dd, J=9.4, 14.2Hz), 3.21 (3H, s), 3.25-3.35 (2H, m), 3.38 (1H, dd, J=3.4, 12.8Hz), 3.65-3.69 (1H, m), 5.15 (1H, dd, J=3.2, 10.0Hz), 6.43 (1H, d, J=7.6Hz), 6.89 (1H, t, J=7.9Hz), 7.17 (1H, s), 7.23 (1H, d, J=7.7Hz), 7.78 (1H, dd, J=5.3, 8.1Hz), 8.28 (1H, d, J=8.0Hz), 8.68 (1H, dd, J=1.3, 5.3Hz), 8.77 (1H, d, J=1.8Hz).

Example 116

N,N-Diethyl-2-((3-((2R)-2-(((2R)-2-hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)-2-methylpropanamide · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example

110.

¹H-NMR (CDCl₃) δ: 0.88 (3H, t, J=7.1Hz), 1.11-1.15 (6H, m), 1.71 (6H, s), 2.65 (1H, dd, J=9.3, 12.2Hz), 2.80-2.83 (2H, m), 2.92 (1H, dd, J=3.6, 12.2Hz), 3.04-3.09 (1H, m), 3.39 (2H, q, J=7.0Hz), 3.69 (2H, d, J=7.1Hz), 4.54 (1H, dd, J=3.6, 9.2Hz), 6.59 (1H, d, J=7.7Hz), 6.92 (1H, t, J=7.9Hz), 7.01 (1H, d, J=2.2Hz), 7.19 (1H, d, J=8.0Hz), 7.22-7.24 (1H, m), 7.65-7.68 (1H, m), 8.24 (1H, brs), 8.50 (1H, dd, J=1.6, 4.8Hz), 8.54 (1H, d, J=2.1Hz).

Example 117

(1R)-2-(((1R)-2-(7-(1,1-Dimethyl-2-oxo-2-piperidin-1-ylethoxy)-1H-indol-3-yl)-1-methylethyl)amino)-1-pyridin-3-ylethanol · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 11.

¹H-NMR (CD₃OD) δ: 1.10 (2H, brs), 1.30 (3H, d, J=6.5Hz), 1.47 (4H, brs), 1.69 (6H, s), 2.97 (1H, dd, J=9.6, 14.1Hz), 3.31 (2H, m), 3.45 (1H, dd, J=3.1, 12.7Hz), 3.56 (2H, brs), 3.65 (1H, m), 3.84 (2H, brs), 5.26 (1H, d, J=3.0, 9.9Hz), 6.50 (1H, d, J=7.7Hz), 6.88 (1H, t, J=7.9Hz), 7.17 (1H, s), 7.24 (1H, d, J=14.3Hz), 8.01 (1H, m), 8.56 (1H, d, J=12.9Hz), 8.80 (2H, d, J=5.3Hz), 8.91 (1H, s).

Example 118

(1R)-2-(((1R)-2-(7-(1,1-Dimethyl-2-morpholin-4-yl-2-oxoethoxy)-1H-indol-3-yl)-1-methylethyl)amino)-1-pyridin-3-ylethanol · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 11.

¹H-NMR (CD₃OD) δ: 1.30 (3H, d, J=6.5Hz), 1.71 (6H, s), 3.00 (1H, dd, J=6.7, 16.4Hz), 3.31 (2H, m), 3.45 (1H, dd, J=4.6, 12.8Hz),

3.50 (2H, brs), 3.60 (2H, brs), 3.67 (1H, m), 3.89 (2H, brs), 5.26 (1H, dd, $J=3.0, 10.0\text{Hz}$), 6.53 (1H, d, $J=7.7\text{Hz}$), 6.92 (1H, t, $J=7.9\text{Hz}$), 7.18 (1H, s), 7.25 (1H, d, $J=7.9\text{Hz}$), 8.00 (1H, dd, $J=5.6, 8.1\text{Hz}$), 8.56 (1H, d, $J=8.1\text{Hz}$), 8.80 (1H, d, $J=5.3\text{Hz}$), 8.91 (1H, s).

5 Example 119

(1R)-2-(((1R)-1-Methyl-2-(7-(((1R)-1-methyl-2-morpholin-4-yl-2-oxoethyl)oxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol

To a solution of tert-butyl (1R)-2-(7-hydroxy-1H-indol-3-yl)-1-methylethyl((2R)-2-pyridin-3-yl-2-((triethylsilyl)oxy)ethyl)carbamate (300 mg, 0.570 mmol) in N,N-dimethylformamide (1.0 mL) are added
 10 potassium carbonate (98.5 mg, 0.714 mmol), (1S)-1-methyl-2-morpholin-4-yl-2-oxoethyl 4-methylbenzenesulfonate (JP-A-2000-273085) (196 mg, 0.627 mmol), and the mixture is stirred at 50°C for one hour. To the reaction solution are added potassium carbonate (45
 15 mg), (1S)-1-methyl-2-morpholin-4-yl-2-oxoethyl 4-methylbenzenesulfonate (120 mg), and the mixture is stirred at 60°C for 10 hours. After allowed to cool, to the reaction solution is added ethyl acetate (50 mL), and the mixture is filtered, washed with water and a saturated brine, and dried over anhydrous sodium sulfate, and concentrated
 20 under reduced pressure. The obtained residue is purified by silica gel column chromatography (chloroform/methanol = 100/1 → 20/1) to give tert-butyl (1R)-1-methyl-2-(7-(((1R)-1-methyl-2-morpholin-4-yl-2-oxoethyl)oxy)-1H-indol-3-yl)ethyl((2R)-2-pyridin-3-yl-2-((triethylsilyl)oxy)-ethyl)carbamate (209 mg, yield: 55 %).

25 Further, to a solution of tert-butyl (1R)-1-methyl-2-(7-(((1R)-1-methyl-2-morpholin-4-yl-2-oxoethyl)oxy)-1H-indol-3-yl)ethyl((2R)-2-pyridin-3-yl-2-((triethylsilyl)oxy)ethyl)carbamate (190 mg, 0.285 mmol)

in ethanol (2.0 mL) is added a 4N hydrochloric acid/dioxane (2.0 mL, 8.0 mmol), and the mixture is stirred at room temperature for 14 hours. To the reaction solution are added a saturated aqueous sodium hydrogen carbonate solution and a saturated brine, and the mixture is extracted with chloroform, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue is purified by silica gel column chromatography (a saturated ammonia solution in chloroform/methanol = 100/1 → 100/4) to give the title compound (107 mg, yield: 83 %).

¹H-NMR (CDCl₃) δ: 1.14 (3H, d, J=6.2Hz), 1.65 (3H, d, J=6.6Hz), 2.66 (1H, dd, J=9.3, 12.2Hz), 2.83 (2H, m), 2.89 (1H, dd, J=3.6, 12.3Hz), 3.05 (1H, m), 3.50 (2H, m), 3.60 (6H, m), 4.54 (1H, dd, J=3.4, 9.3Hz), 5.14 (1H, q, J=6.7Hz), 6.62 (1H, d, J=7.7Hz), 6.99 (1H, t, J=7.9Hz), 7.02 (1H, m), 7.24 (1H, m), 7.25 (1H, m), 7.66 (1H, dt, J=1.7, 7.9Hz), 8.49 (1H, dd, J=1.6, 4.8Hz), 8.52 (1H, d, J=2.2Hz), 8.70 (1H, brs).

Example 120

(2R)-2-((3-(((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)propanoic acid · 2 trifluoroacetate

To a solution of (1R)-2-(((1R)-1-methyl-2-(7-(((1R)-1-methyl-2-morpholin-4-yl-2-oxoethoxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol (96.3 mg, 0.213 mmol) in methanol (850 μL) are added tetrahydrofuran (850 μL) and a 2N lithium hydroxide (850 μL), and the mixture is stirred at 50°C for one hour. The mixture is neutralized with 1N hydrochloric acid at 0°C, and thereto is added a pH standard solution (pH 6.8, 2.0 mL). The mixture is concentrated under reduced pressure, and purified by octadecylsilyl (ODS) column chromatography (Cosmosil, a 0.1% aqueous trifluoroacetic acid solution/methanol =

100/1 → 70/30) to give the title compound (99.7 mg, yield: 77 %).

¹H-NMR (CD₃OD) δ: 1.33 (3H, d, J=6.5Hz), 1.67 (3H, d, J=6.6Hz), 3.01 (1H, dd, J=8.8, 14.3Hz), 3.30 (2H, m), 3.40 (1H, dd, J=3.3, 12.8Hz), 3.70 (1H, m), 4.99 (1H, m), 5.23 (1H, dd, J=3.1, 9.6Hz), 6.57 (1H, d, J=7.7Hz), 6.95 (1H, t, J=7.9Hz), 7.18 (1H, s), 7.21 (1H, d, J=7.7Hz), 8.01 (1H, dd, J=5.7, 8.0Hz), 8.51 (1H, m), 8.78 (1H, d, J=5.2Hz), 8.87 (1H, m).

Example 121

(2R)-2-(((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)-N,N-dimethylpropanamide · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 110.

¹H-NMR (CD₃OD) δ: 1.33 (3H, d, J=6.5Hz), 1.60 (3H, d, J=6.6Hz), 2.97 (3H, s), 3.02 (1H, dd, J=8.7, 14.3Hz), 3.19 (3H, s), 3.30 (2H, m), 3.35 (1H, dd, J=3.4, 12.8Hz), 3.68 (1H, m), 5.16 (1H, dd, J=3.2, 9.7Hz), 5.36 (1H, q, J=6.6Hz), 6.53 (1H, d, J=7.7Hz), 6.94 (1H, t, J=7.9Hz), 7.18 (1H, s), 7.21 (1H, d, J=7.7Hz), 7.91 (1H, m), 8.51 (1H, m), 8.75 (2H, brs).

Example 122

(1R)-2-(((1R)-1-Methyl-2-(7-(((1R)-1-methyl-2-oxo-2-piperidin-1-ylethyl)-oxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 110.

¹H-NMR (CD₃OD) δ: 1.32 (3H, d, J=6.5Hz), 1.5-1.7 (6H, m), 1.61 (3H, d, J=6.6Hz), 3.00 (1H, dd, J=8.9, 14.3Hz), 3.30 (2H, m), 3.39 (1H, dd, J=3.3, 12.8Hz), 3.56 (2H, t, J=5.5Hz), 3.66 (3H, m), 5.23 (1H, dd, J=3.0, 9.6Hz), 5.35 (1H, q, J=6.6Hz), 6.56 (1H, d, J=7.7Hz), 6.95 (1H, t,

$J=7.9\text{Hz}$), 7.18 (1H, s), 7.21 (1H, d, $J=7.8\text{Hz}$), 8.04 (1H, dd, $J=5.7$, 8.1Hz), 8.57 (1H, m), 8.81 (2H, m).

Reference Example 66

(1R)-1-Methyl-2-morpholin-4-yl-2-oxoethyl 4-methylbenzenesulfonate

5 Under nitrogen atmosphere, to methyl D-(+)-lactate (20.8 g, 200 mmol) and morpholine (19.1 mL, 220 mmol) is added in portions a 60 % sodium hydride (800 mg, 20.0mmol) with stirring under ice-cooling, and the mixture is heated at 50°C with stirring for 3 hours. The mixture is cooled to room temperature, and subjected to azeotropic distillation
10 with toluene to remove excess morpholine. The resultant is dried under reduced pressure to give lactic morpholinamide (32.1 g).

Subsequently, to a suspension of 60 % sodium hydride (8.41 g, 210 mmol) in tetrahydrofuran (120 mL) is added dropwise a solution of the above lactic morpholinamide (32.1 g) in tetrahydrofuran (150 mL)
15 under nitrogen atmosphere with stirring under ice-cooling, and the mixture is heated with stirring at 50°C for 30 minutes. After cooling with ice, to the mixture is added dropwise a solution of p-toluene-sulfonyl chloride (45.8 g, 234 mmol) in tetrahydrofuran (180 mL), and the mixture is stirred at room temperature for 4 hours. To the mixture
20 is added 1N aqueous hydrochloric acid solution, and the mixture is extracted with ethyl acetate. The organic layer is washed with water and a saturated brine, dried over anhydrous magnesium sulfate, and the solvent is evaporated under reduced pressure. To the resulting oily residue is added diethyl ether, and the precipitated crystals are
25 collected by filtration, washed with diethyl ether, and dried under reduced pressure to give the title compound (36.1 g, yield: 58 %).

$^1\text{H-NMR}$ (CDCl_3) δ : 7.81 (2H, d, $J=8.0\text{Hz}$), 7.35 (2H, d, $J=8.0\text{Hz}$),

5.27 (1H, q, J=6.8Hz), 3.64-3.46 (8H, m), 2.46 (3H, s), 1.47 (3H, d, J=6.8Hz).

Example 123

(1R)-2-(((1R)-1-Methyl-2-(7-(((1S)-1-methyl-2-morpholin-4-yl-2-oxoethyl-
oxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.15 (3H, d, J=6.2Hz), 1.66 (3H, d, J=6.7Hz), 2.67 (1H, dd, J=9.4, 12.2Hz), 2.83 (2H, m), 2.91 (1H, dd, J=3.6, 12.2Hz), 3.07 (1H, m), 3.50 (2H, m), 3.61 (6H, m), 4.59 (1H, dd, J=3.5, 9.3Hz), 5.13 (1H, q, J=6.7Hz), 6.63 (1H, d, J=7.7Hz), 7.00 (1H, t, J=7.8Hz), 7.03 (1H, m), 7.23 (1H, m), 7.25 (1H, m), 7.66 (1H, dt, J=1.7, 7.9Hz), 8.49 (1H, dd, J=1.7, 4.8Hz), 8.53 (1H, d, J=2.1Hz), 8.60 (1H, brs).

Example 124

(2S)-2-(((3-(((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)propanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CD₃OD) δ: 1.32 (3H, d, J=6.5Hz), 1.67 (3H, d, J=6.8Hz), 3.00 (1H, dd, J=9.0, 14.2Hz), 3.30 (2H, m), 3.40 (1H, dd, J=3.2, 12.6Hz), 3.67 (1H, m), 4.97 (1H, q, J=6.8Hz), 5.23 (1H, dd, J=3.0, 10.5Hz), 6.57 (1H, d, J=7.7Hz), 6.95 (1H, t, J=7.9Hz), 7.18 (1H, s), 7.22 (1H, d, J=7.7Hz), 7.98 (1H, dd, J=5.7, 8.0Hz), 8.47 (1H, m), 8.78 (1H, d, J=5.2Hz), 8.86 (1H, m).

Example 125

Ethyl (2S)-2-(((3-(((2R)-2-(((2R)-2-hydroxy-2-pyridin-3-ylethyl)amino)-propyl)-1H-indol-7-yl)oxy)propanoate

To a suspension of (2S)-2-((3-(((2R)-2-(((2R)-2-hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)propanoic acid (6.72 g, 17.53 mmol) in ethanol (135 mL) is added a 4N solution of hydrogen chloride in dioxane (55 mL), and the mixture is stirred at room temperature for 2 hours. The reaction solution is evaporated to remove the solvent, and the residue is separated into a saturated aqueous sodium hydrogen carbonate solution and chloroform. The organic layer is washed with a saturated brine, and dried over anhydrous sodium sulfate. The mixture is evaporated to remove the solvent, and to the obtained crude product are added ethyl acetate (50 mL) and diisopropyl ether (50 mL). The product is crystallized and collected by filtration. The product on the filter is washed with diisopropyl ether, and dried to give the title compound (6.26 g, yield: 87 %).

Example 126

(2S)-6-(((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-2-methyl [1,4]oxazino[2,3,4-hi]indol-3(2H)-one

To a solution of ethyl (2S)-2-((3-(((2R)-2-(((2R)-2-hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)propanoate · 2 hydrochloride (205 mg, 0.423 mmol) in acetonitrile (8.7 ml) is added potassium carbonate (294 mg, 2.13 mmol), and the mixture is stirred at 60°C for 4 hours. After the reaction is completed, the mixture is cooled to room temperature, and filtered. The filtrate is evaporated under reduced pressure to remove the solvent, and the obtained residue is purified by preparative thin layer chromatography (chloroform/methanol = 10/1) to give the title compound (131 mg, yield: 85 %).

¹H-NMR (CDCl₃) δ: 1.16 (3H, d, J=6.3Hz), 1.74 (3H, d, J=6.9Hz), 2.74 (1H, dd, J=12, 9.3Hz), 2.77 (1H, dd, J=14.3, 6.5Hz), 2.88 (1H, dd,

J=14.3, 6.6Hz), 2.96 (1H, dd, J=12, 3.5Hz), 3.10-3.15 (1H, m), 4.65 (1H, dd, J=9.3, 3.5Hz), 5.10 (1H, q, J=6.9Hz), 6.83 (1H, dd, J=6.6, 1.7Hz), 7.13-7.18 (2H, m), 7.25-7.28 (1H, m), 7.41 (1H, s), 7.70 (1H, d, J=7.9Hz), 8.50 (1H, dd, J=4.8, 1.4Hz), 8.55 (1H, d, J=1.7Hz).

5 Example 127

(2S)-2-((3-(((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)-N,N-dimethylpropanamide · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 110.

10 ¹H-NMR (CD₃OD) δ: 1.32 (3H, d, J=6.5Hz), 1.60 (3H, d, J=6.6Hz), 2.97 (3H, s), 3.02 (1H, m), 3.19 (3H, s), 3.30 (2H, m), 3.39 (1H, dd, J=3.2, 12.8Hz), 3.68 (1H, m), 5.22 (1H, dd, J=3.0, 9.7Hz), 5.36 (1H, q, J=6.6Hz), 6.54 (1H, d, J=7.7Hz), 6.95 (1H, t, J=7.9Hz), 7.18 (1H, s), 7.22 (1H, d, J=7.7Hz), 8.00 (1H, dd, J=5.9, 8.0Hz), 8.51 (1H, m), 8.80
15 (2H, m).

Example 128

(1R)-2-(((1R)-1-Methyl-2-(7-(((1S)-1-methyl-2-oxo-2-piperidin-1-ylethyl)-oxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol · 2 trifluoroacetate

20 The title compound is obtained in a similar manner to Example 110.

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 1.5-1.7 (6H, m), 1.61 (3H, d, J=6.6Hz), 3.00 (1H, dd, J=9.1, 14.2Hz), 3.30 (2H, m), 3.42 (1H, dd, J=3.2, 12.8Hz), 3.56 (2H, t, J=5.5Hz), 3.66 (3H, m), 5.25 (1H, dd, J=3.0, 9.7Hz), 5.34 (1H, q, J=6.6Hz), 6.57 (1H, d, J=7.7Hz), 6.95 (1H, t, J=7.9Hz), 7.18 (1H, s), 7.22 (1H, d, J=7.9Hz), 8.03 (1H, dd, J=5.7, 8.1Hz), 8.56 (1H, m), 8.81 (2H, m).

Reference Example 67

4-(2-Bromobutanoyl)morpholine

To a solution of 2-bromobutanoic acid (0.3 mL, 2.81 mmol) in N,N-dimethylformamide (14 mL) are added morpholine (0.3 mL, 3.44 mmol), 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (650 mg, 3.39 mmol) and 7-hydroxybenzotriazole (456 mg, 3.38 mmol), and the mixture is stirred at room temperature for 4.5 hours. After the reaction is completed, water is added to the mixture, and the mixture is extracted with ethyl acetate. The organic layer is washed with 1N-hydrochloric acid, a saturated aqueous sodium hydrogen carbonate solution, and a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated, and the obtained residue is purified by silica gel column chromatography (n-hexane/ethyl acetate = 3/1) to give the title compound (323 mg, yield: 49 %).

$^1\text{H-NMR}$ (CDCl_3) δ : 1.04 (3H, t, $J=7.3\text{Hz}$), 1.96-2.16 (2H, m), 3.46-3.54 (2H, m), 3.62-3.74 (5H, m), 3.79-3.84 (1H, m), 4.28 (1H, dd, $J=7.8, 6.2\text{Hz}$).

Example 129

(1R)-2-(((1R)-1-Methyl-2-(7-(1-(Morpholin-4-ylcarbonyl)propoxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 119.

$^1\text{H-NMR}$ (CD_3OD) δ : 1.14 and 1.15 (total 3H, J, t=7.3Hz), 1.33 and 1.34 (total 3H, d, $J=6.5\text{Hz}$), 2.00-2.08 (2H, m), 3.02-3.05 (1H, m), 3.27-3.42 (3H, m), 3.52-3.82 (9H, m), 5.10-5.15 (1H, m), 5.21-5.25 (1H, m), 6.57 and 6.58 (total 1H, d, $J=7.7\text{Hz}$), 6.96 (1H, t, $J=7.8\text{Hz}$), 7.20 (s, 1H), 7.24 and 7.25 (total 1H, d, $J=8.0\text{Hz}$), 8.00 (1H, m), 8.52 (1H, d, $J=8.2\text{Hz}$), 8.80 (1H, d, $J=4.2\text{Hz}$), 8.84 (s, 1H).

Example 130

2-((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)butanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CD₃OD) δ: 1.14 (3H, t, J=7.4Hz), 1.35 and 1.34 (total 3H, d, J=6.5Hz), 1.99-2.14 (2H, m), 3.03-3.06 (1H, m), 3.25-3.36 (3H, m), 3.66-3.70 (1H, m), 4.81-4.88 (1H, m), 5.11-5.17 (m, 1H), 6.57 (1H, d, J=7.7Hz), 6.95 (1H, t, J=7.9Hz), 7.19 (1H, s), 7.22 (1H, d, J=8.1Hz), 7.81 (dd, 1H, J=7.8, 5.6Hz), 8.26 (1H, d, J=8.2Hz), 8.69 (1H, s), 8.76 (1H, brs).

Example 131

2-((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)pentanoic acid · 2 trifluoroacetate

6-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-2-propyl-[1,4]oxadino[2,3,4-hi]indol-3(2H)-one · 2 trifluoroacetate

To a solution of tert-butyl (1R)-2-(7-hydroxy-1H-indol-3-yl)-1-methylethyl ((2R)-2-pyridin-3-yl-2-((triethylsilyl)oxy)ethyl)carbamate (49.7 mg, 0.0945 mmol) in acetonitrile (2 ml) are added potassium carbonate (20.1 mg, 0.176 mmol), ethyl 2-bromo-n-valerate (0.03 ml, 0.176 mmol) and a catalytic amount of potassium iodide, and the mixture is stirred at 70°C for 2 days. After the reaction is completed, the mixture is cooled to room temperature, filtered, and the solvent is evaporated under reduced pressure. The resulting residue is purified by preparative thin layer chromatography (n-hexane/ethyl acetate = 2/1) to give ethyl 2-((3-((2R)-2-((tert-butoxycarbonyl)((2R)-2-pyridin-3-yl-2-((triethylsilyl)oxy)ethyl)amino)propyl)-1H-indol-7-yl)oxy)pentanoate

(3.6 mg, yield: 5.7 %) and tert-butyl (1R)-1-methyl-2-(3-oxo-2-propyl-2,3-dihydro[1,4]oxazino[2,3,4-hi]indol-6-yl)ethyl((2R)-2-pyridin-3-yl-2-(triethylsilyl)oxy)ethyl)carbamate (9.5 mg, yield: 16 %).

To a solution of the obtained ethyl 2-((3-((2R)-2-((tert-butoxycarbonyl)((2R)-2-pyridin-3-yl-2-((triethylsilyl)oxy)ethyl)amino)propyl)-1H-indol-7-yl)oxy)pentanoate in ethanol (0.2 ml) is added a solution of 4N-hydrogen chloride in dioxane (0.2 ml), and the mixture is stirred at room temperature for 19 hours. After the reaction is completed, water (0.2 ml) is added to the mixture, and the solvent is evaporated under reduced pressure. To a solution of the obtained residue in ethanol (0.35 ml) is added a 5N aqueous potassium hydroxide solution (0.09 ml), and the mixture is stirred at room temperature for 4 hours. After the reaction is completed, the mixture is neutralized with 1N hydrochloric acid, and the solvent is evaporated under reduced pressure. The residue is purified by preparative high performance liquid chromatography (0.035 % trifluoroacetic acid/acetonitrile-0.05% trifluoroacetic acid/water) to give 2-((3-((2R)-2-((2R)-2-hydroxy-2-pyridin-3-ylethyl)-amino)propyl)-1H-indol-7-yl)oxy)pentanoic acid · 2 trifluoroacetate (13.2 mg).

Further, to a solution of tert-butyl (1R)-1-methyl-2-(3-oxo-2-propyl-2,3-dihydro[1,4]oxazino[2,3,4-hi]indol-6-yl)ethyl((2R)-2-pyridin-3-yl-2-(triethylsilyl)oxy)ethyl)carbamate (9.5 mg, 0.0156 mmol) in ethanol (0.12 ml) is added a 4N solution of hydrogen chloride in dioxane (0.12 ml), and the mixture is stirred at room temperature for 15 hours. After the reaction is completed, water (0.12 ml) is added to the mixture, and the solvent is evaporated under reduced pressure. The obtained residue is purified by preparative high performance liquid chromato-

graphy (0.035 % trifluoroacetic acid/acetonitrile-0.05 % trifluoroacetic acid/water) to give 6-((2R)-2-(((2R)-2-hydroxy-2-pyridin-3-ylethyl)-amino)propyl)-2-propyl[1,4]oxazino[2,3,4-hi]indol-3(2H)-one · 2 trifluoroacetate (3.7 mg).

5 2-((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)pentanoic · 2 trifluoroacetate

¹H-NMR (CD₃OD) δ: 1.00 (3H, t, J=7.3Hz), 1.34 (3H, t, J=5.5Hz), 1.55-1.68 (2H, m), 1.96-2.09 (2H, m), 3.01 (1H, dt, J=14, 8.3Hz), 3.24-3.41 (m, 2H), 3.64-3.67 (1H, m), 4.84-4.99 (2H, m), 5.20 (1H, t, J=9.4Hz), 6.54 (1H, d, J=7.7Hz), 6.94 (1H, t, J=7.7Hz), 7.18 (1H, s), 7.20 (1H, d, J=7.8Hz), 7.92 (1H, s), 8.40 (1H, s), 8.75 (1H, s), 8.82 (1H, s).

6-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-2-propyl-[1,4]oxazino[2,3,4-hi]indol-3(2H)-one · 2 trifluoroacetate

15 ¹H-NMR (CD₃OD) δ: 0.99 (3H, t, J=7.4Hz), 1.35 (3H, d, J=6.5Hz), 1.54-1.64 (2H, m), 1.96-2.04 (2H, m), 3.04 (1H, dd, J=14, 9.7Hz), 3.30-3.48 (3H, m), 3.76-3.81 (1H, m), 5.09 (1H, dd, J=6.9, 5.1Hz), 5.20 (1H, dd, J=9.7, 3.0Hz), 6.86 (1H, d, J=7.7Hz), 7.20 (1H, dd, J=7.9, 7.7Hz), 7.26 (1H, d, J=7.9Hz), 7.65 (1H, s), 7.79 (1H, dd, J=7.4, 5.6Hz), 8.31 (1H, d, J=8.0Hz), 8.70 (1H, brs), 8.81 (1H, brs).

Example 132

6-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-2-isopropyl-[1,4]oxazino[2,3,4-hi]indol-3(2H)-one · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 131.

25 ¹H-NMR (CD₃OD) δ: 0.97 (3H, d, J=6.8Hz), 1.17 (3H, d, J=7.0Hz), 1.35, 1.36 (total 3H, d, J=6.5Hz), 2.49-2.56 (1H, m), 3.04 (1H,

dd, J=14, 10Hz), 3.30-3.40 (2H, m), 3.50 (1H, dt, J=12, 3.1Hz), 3.37-3.83 (2H, m), 4.94 (1H, dd, J=3.6, 0.9Hz), 5.30 (1H, dd, J=9.8, 1.8Hz), 6.86 (2H, t, J=7.6Hz), 7.19 (1H, t, J=7.8Hz), 7.25 (1H, d, J=7.8Hz), 7.67 (1H, s), 8.01 (1H, dd, J=7.8, 5.7Hz), 8.58 (1H, d, J=8.1Hz), 8.80 (1H, d, J=5.2Hz), 8.92 (1H, s).

Reference Example 68

3-Methyl-1-morpholin-4-yl-1-oxobutan-2-ol

To a solution of 2-hydroxy-i-valeric acid (501 mg, 4.24 mmol) in tetrahydrofuran (20 mL) are added 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (976 mg, 5.09 mmol), 7-hydroxybenzotriazole (691 mg, 5.11 mmol) and morpholine (0.44 mL, 5.05 mmol), and the mixture is stirred at room temperature for 9 hours. After the reaction is completed, the solvent is evaporated under reduced pressure, and the resulting residue is purified by silica gel column chromatography (n-hexane/ethyl acetate = 2/1 → 1/1) to give the title compound (448 mg, yield: 56 %).

¹H-NMR (CDCl₃) δ: 0.82 (3H, d, J=6.8Hz), 1.08 (3H, d, J=6.9Hz), 1.77-1.88 (1H, m), 3.38-3.48 (2H, m), 3.59-3.75 (6H, m), 4.23 (1H, d, J=2.8Hz).

Reference Example 69

2-Methyl-1-(morpholin-4-ylcarbonyl)propyl 4-methylbenzenesulfonate

Under ice-cooling, to a solution of 60 % sodium hydride (136 mg, 3.41 mmol) in tetrahydrofuran (5 mL) is added dropwise a solution of 3-methyl-1-morpholin-4-yl-1-oxobutan-2-ol (426 mg, 2.27 mmol) in tetrahydrofuran (15 mL), and the mixture is stirred at room temperature for 2 hours. To the reaction solution is added p-toluenesulfonyl chloride (651 mg, 3.42 mmol), and the mixture is stirred at

room temperature for 2 hours. After the reaction is completed, water is added to the mixture, and the mixture is extracted with ethyl acetate. This organic layer is washed with water, a saturated aqueous sodium hydrogen carbonate solution, and a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is removed by distillation, and the obtained residue is purified by silica gel column chromatography (chloroform) to give the title compound (701 mg, yield: 90 %).

$^1\text{H-NMR}$ (CDCl_3) δ : 0.92 (3H, d, $J=6.8\text{Hz}$), 0.97 (3H, d, $J=6.6\text{Hz}$), 2.07-2.16 (1H, m), 2.45 (3H, s), 3.41-3.65 (8H, m), 4.74 (1H, d, $J=8.4\text{Hz}$), 7.35 (2H, d, $J=8.0\text{Hz}$), 7.81 (2H, d, $J=8.4\text{Hz}$).

Example 133

(1R)-2-(((1R)-1-Methyl-2-(7-(2-methyl-1-(morpholin-4-ylcarbonyl)propoxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 119.

$^1\text{H-NMR}$ (CD_3OD) δ : 1.13 and 1.10 (total 3H, d, $J=6.4\text{Hz}$), 1.24 and 1.22 (total 3H, d, $J=5.6\text{Hz}$), 2.30-2.35 (1H, m), 2.73-2.92 (4H, m), 3.04-3.07 (1H, m), 3.24-3.63 (8H, m), 3.76-3.77 (1H, m), 3.85-3.88 (1H, m), 4.69-4.74 (1H, m), 4.77-4.80 (1H, m), 6.55 and 6.54 (total 1H, d, $J=7.6\text{Hz}$), 6.89 (1H, t, $J=7.8\text{Hz}$), 7.02 (d, 1H, $J=5.3\text{Hz}$), 7.16 (1H, d, $J=7.9\text{Hz}$), 7.25-7.31 (1H, m), 7.67 and 7.65 (total 1H, $J_d=8.4\text{Hz}$), 8.37 and 8.36 (total 1H, d, $J=5.2\text{Hz}$), 8.45 (1H, brs).

Example 134

2-((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)-3-methylbutanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example

120.

¹H-NMR (CD₃OD) δ: 1.19, 1.15 (total 3H, d, J=6.8Hz), 1.33 and 1.34 (total 3H, d, J=6.5Hz), 2.33-2.41 (1H, m), 2.98-3.06 (1H, m), 3.24-3.35 (2H, m), 3.37-3.44 (1H, m), 3.65-3.70 (1H, m), 4.61-4.62 (1H, m), 5.20-5.26 (1H, m), 6.54 (1H, d, J=7.7Hz), 6.94 (1H, t, J=7.9Hz), 7.19 (1H, s), 7.21 (1H, d, J=8.0Hz), 7.97-8.00 (1H, m), 8.48 (1H, d, J=7.9Hz), 8.78 (1H, d, J=4.8Hz), 8.86 (1H, brs).

Reference Example 70

(2S)-3-Methyl-1-morpholin-4-yl-1-oxobutan-2-ol

The title compound is obtained in a similar manner to Reference Example 68.

¹H-NMR (CDCl₃) δ: 0.82 (3H, d, J=6.8Hz), 1.07 (3H, d, J=6.9Hz), 1.78-1.86 (1H, m), 3.42-3.44 (2H, m), 3.56 (1H, d, J=7.4Hz), 3.61-3.75 (6H, m), 4.23 (1H, dd, J=7.4, 2.8Hz).

Reference Example 71

(1S)-2-Methyl-1-(morpholin-4-ylcarbonyl)propyl 4-methylbenzenesulfonate

The title compound is obtained in a similar manner to Reference Example 69.

¹H-NMR (CDCl₃) δ: 0.92 (3H, d, J=6.8Hz), 0.97 (3H, d, J=6.6Hz), 2.04-2.15 (1H, m), 2.45 (3H, s), 3.43-3.65 (8H, m), 4.74 (1H, d, J=8.4Hz), 7.35 (2H, d, J=8.0Hz), 7.81 (2H, d, J=8.3Hz).

Example 135

(1R)-2-(((1R)-1-Methyl-2-(7-(((1R)-2-methyl-1-(morpholin-4-ylcarbonyl)propyl)oxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.11 (3H, d, J=6.8Hz), 1.18 (3H, d, J=6.4Hz), 1.20 (3H, d, J=6.7Hz), 2.24-2.32 (1H, m), 2.71-2.78 (1H, m), 2.89-2.97 (4H, m), 3.14-3.19 (1H, m), 3.36 (2H, t, J=4.4Hz), 3.48-3.67 (m, 6H), 4.65 (1H, d, J=7.6Hz), 4.70 (1H, dd, J=9.3, 3.0Hz), 6.64 (1H, d, J=7.7Hz), 6.98 (1H, t, J=7.9Hz), 7.03 (1H, d, J=1.7Hz), 7.22-7.25 (1H, m), 7.67 (dt, J=7.9, 1.8Hz), 8.53 (1H, d, 1H, J=1.9Hz), 8.49 (1H, dd, J=4.8, 1.5Hz), 8.81 (1H, brs).

Example 136

(2R)-2-((3-(((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)-3-methylbutanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CD₃OD) δ: 1.15 (3H, d, J=6.9Hz), 1.20 (3H, d, J=6.8Hz), 1.35 (3H, d, J=6.5Hz), 2.33-2.41 (1H, m), 3.03 (1H, dd, J=14, 8.8Hz), 3.27-3.37 (3H, m), 3.66-3.74 (1H, m), 4.62 (1H, d, J=5.1Hz), 5.15 (1H, dd, J=9.7, 2.9Hz), 6.54 (1H, d, J=7.7Hz), 6.94 (1H, t, J=7.9Hz), 7.19 (1H, s), 7.21 (2H, d, J=8.0Hz), 7.85 (1H, dd, J=7.4, 5.8Hz), 8.30 (1H, d, J=8.0Hz), 8.72 (1H, m), 8.78 (1H, brs).

Reference Example 72

(2R)-3-Methyl-1-morpholin-4-yl-1-oxobutan-2-ol

The title compound is obtained in a similar manner to Reference Example 68.

¹H-NMR (CDCl₃) δ: 0.82 (3H, d, J=6.8Hz), 1.08 (3H, d, J=6.9Hz), 1.78-1.86 (1H, m), 3.42-3.44 (2H, m), 3.55 (1H, d, J=7.4Hz), 3.62-3.75 (m, 6H), 4.23 (1H, dd, J=7.4, 2.8Hz).

Reference Example 73

(1R)-2-Methyl-1-(morpholin-4-ylcarbonyl)propyl 4-methylbenzene-

sulfonate

The title compound is obtained in a similar manner to Reference Example 69.

¹H-NMR (CDCl₃) δ: 0.92 (3H, d, J=6.8Hz), 0.97 (3H, d, J=6.6Hz),
 2.06-2.18 (1H, m), 2.45 (3H, s), 3.43-3.69 (8H, m), 4.74 (1H, d, J=8.4Hz),
 7.35 (2H, d, J=8.0Hz), 7.81 (2H, d, J=8.3Hz).

Example 137

(1R)-2-(((1R)-1-Methyl-2-(7-(((1S)-2-methyl-1-(morpholin-4-ylcarbonyl)-propyl)oxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.11 (3H, d, J=6.8Hz), 1.16 (3H, d, J=6.2Hz),
 1.20 (3H, d, J=6.6Hz), 2.25-2.31 (1H, m), 2.70 (1H, dd, J=12, 9.5Hz),
 2.85 (2H, m), 2.91 (1H, dd, J=12, 3.4Hz), 3.08-3.13 (1H, m), 3.34 (2H,
 m), 3.49-3.62 (6H, m), 4.61-4.64 (2H, m), 6.67 (1H, d, J=7.7Hz), 6.99
 (1H, t, J=7.8Hz), 7.03 (1H, s), 7.22-7.26 (1H, m), 8.49 (1H, d, J=4.1Hz),
 8.53 (1H, d, J=1.2Hz), 8.74 (1H, brs).

Example 138

(2S)-2-((3-(((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)-3-methylbutanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CDCl₃) δ: 1.15 (3H, d, J=6.9Hz), 1.19 (3H, d, J=6.8Hz),
 1.33 (3H, d, J=6.5Hz), 2.33-2.41 (1H, m), 3.01 (1H, dd, J=14, 9.0Hz),
 3.26-3.35 (2H, m), 3.42 (1H, dd, J=14, 3.2Hz), 3.65-3.70 (1H, m), 4.61
 (1H, d, J=5.1Hz), 5.24 (1H, dd, J=9.0, 3.2Hz), 6.54 (1H, d, J=7.7Hz),
 6.94 (1H, dd, J=7.8, 7.7Hz), 7.19 (1H, s), 7.22 (1H, d, J=7.8Hz), 7.99

(1H, dd, J=7.9, 5.7Hz), 8.48 (1H, d, J=7.9Hz), 8.79 (1H, d, J=5.7Hz), 8.87 (1H, brs).

Example 139

Ethyl (2S)-2-((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)-propyl)-1H-indol-7-yl)oxy)-3-methylbutanoate · 2 hydrochloride

The title compound is obtained in a similar manner to Example 125.

¹H-NMR (CDCl₃) δ: 1.13 (3H, d, J=6.9Hz), 1.18 (3H, d, J=6.8Hz), 1.23 (3H, t, J=7.1Hz), 1.34 (3H, d, J=6.1Hz), 2.31-2.40 (1H, m), 3.02 (1H, dd, J=13.8, 9.2Hz), 3.31-3.37 (2H, m), 3.47-3.50 (1H, m), 3.70 (1H, m), 4.21 (2H, q, J=7.1Hz), 4.63 (1H, d, J=5.4Hz), 5.38 (1H, m), 6.49 (1H, d, J=7.7Hz), 6.93 (1H, dd, J=7.9, 7.7Hz), 7.22 (1H, s), 7.26 (1H, d, J=7.9Hz), 8.11 (1H, dd, J=7.7, 5.0Hz), 8.68 (1H, d, J=7.7Hz), 8.84 (1H, d, J=5.0Hz), 8.99 (1H, brs).

Example 140

1-((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)cyclobutanecarboxylic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 109.

¹H-NMR (CD₃OD) δ: 1.34 (3H, d, J=6.5Hz), 1.97-2.14 (2H, m), 2.51-2.59 (2H, m), 2.77-2.86 (2H, m), 3.01 (1H, dd, J=14, 8.9Hz), 3.24-3.35 (3H, m), 3.38 (1H, dd, J=13, 2.9Hz), 3.63-3.71 (1H, m), 5.20 (1H, dd, J=9.6, 2.4Hz), 6.21 (1H, d, J=7.7Hz), 6.86 (1H, t, J=7.8Hz), 7.17 (1H, s), 7.18 (1H, d, J=7.4Hz), 7.93 (1H, m), 8.39 (1H, d, J=7.2Hz), 8.75 (1H, brs), 8.82 (1H, brs).

Reference Example 74

(2R)-1-Morpholin-4-yl-1-oxo-3-phenylpropan-2-ol

The title compound is obtained in a similar manner to Reference Example 68.

¹H-NMR (CDCl₃) δ: 2.90 (3H, dd, J=14, 6.2Hz), 2.96 (3H, dd, J=14, 6.8Hz), 3.26-3.32 (2H, m), 3.51-3.74 (6H, m), 4.59 (1H, dt, J=8.4, 6.4Hz), 7.21-7.33 (5H, m).

Reference Example 75

(1R)-1-Benzyl-2-morpholin-4-yl-2-oxoethyl 4-methylbenzenesulfonate

The title compound is obtained in a similar manner to Reference Example 69.

¹H-NMR (CDCl₃) δ: 2.43 (3H, s), 3.00-3.56 (10H, m), 5.26 (1H, dd, J=8.1, 6.8Hz), 7.11-7.15 (2H, m), 7.23-7.29 (5H, m), 7.68 (2H, dt, J=8.3, 1.7Hz).

Example 141

(1R)-2-(((1R)-1-Methyl-2-(7-(((1S)-1-benzyl-2-morpholin-4-yl-2-oxoethyl)-oxy)-1H-indol-3-yl)-1-methylethyl)amino)-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.14 (3H, d, J=6.2Hz), 2.69 (1H, dd, J=12, 9.5Hz), 2.84 (2H, d, J=6.5Hz), 2.91 (1H, dd, J=12, 3.5Hz), 3.06-3.11 (2H, m), 3.27-3.62 (9H, m), 4.63 (1H, dd, J=9.2, 3.1Hz), 5.16 (1H, t, J=6.8Hz), 6.55 (1H, d, J=7.7Hz), 6.95 (1H, t, J=7.9Hz), 6.98 (1H, d, J=1.4Hz), 7.21-7.26 (2H, m), 7.28-7.37 (5H, m), 7.66 (1H, d, J=7.8Hz), 8.48 (1H, dd, J=4.7, 1.2Hz), 8.53 (1H, d, J=1.6Hz), 8.63 (1H, brs).

Example 142

(2S)-2-(((3-(((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)-3-phenylpropanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example

120.

¹H-NMR (CD₃OD) δ: 1.33 (d, 3H, J=6.5Hz), 3.01 (1H, dd, J=14, 8.9Hz), 3.20-3.42 (5H, m), 3.64-3.69 (1H, m), 5.06 (1H, dd, J=7.5, 4.6Hz), 5.22 (1H, dd, J=9.8, 2.9Hz), 6.55 (1H, d, J=7.7Hz), 6.93 (1H, t, J=7.9Hz), 7.12-7.28 (4H, m), 7.35 (2H, d, J=7.2Hz), 7.97 (1H, dd, J=7.9, 5.7Hz), 8.47 (1H, d, J=8.1Hz), 8.77 (1H, d, J=5.3Hz), 8.86 (1H, brs).

Reference Example 76

(1R)-2-Morpholin-4-yl-2-oxo-1-phenylethanol

The title compound is obtained in a similar manner to

Reference Example 68.

¹H-NMR (CDCl₃) δ: 3.05-3.10 (1H, m), 3.14-3.19 (1H, m), 3.28-3.32 (1H, m), 3.45-3.50 (1H, m), 3.55-3.82 (4H, m), 4.71 (1H, brs), 5.19 (1H, s), 7.29-7.41 (5H, m).

Reference Example 77

(1R)-2-Morpholin-4-yl-2-oxo-1-phenylethyl 4-methylbenzenesulfonate

The title compound is obtained in a similar manner to

Reference Example 69.

¹H-NMR (CDCl₃) δ: 2.43 (3H, s), 3.35-3.61 (8H, m), 6.11 (1H, s), 7.31 (2H, d, J=8.1Hz), 7.34 (5H, s), 7.81 (2H, d, J=8.1Hz).

Example 143

(1R)-2-(((1R)-1-Methyl-2-(7-(((1S)-2-morpholin-4-yl-2-oxo-1-phenylethyl)oxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.16 (3H, d, J=6.2Hz), 2.71 (1H, dd, J=12.2, 9.4Hz), 2.86 (2H, m), 2.91 (1H, dd, J=12.2, 3.4Hz), 3.07-3.15 (1H, m), 3.19-3.22 (1H, m), 3.34-3.36 (1H, m), 3.50-3.63 (6H, m), 4.64 (1H, dd,

J=9.4, 3.4Hz), 6.03 (1H, s), 6.74 (1H, d, J=7.8Hz), 6.98 (1H, t, J=7.8Hz), 7.03 (1H, s), 7.21-7.26 (2H, m), 7.36-7.45 (3H, m), 7.56-7.58 (2H, m), 7.66 (1H, dt, J=6.2, 1.7Hz), 8.48 (1H, dd, J=4.7, 1.3Hz), 8.53 (1H, d, J=1.7Hz), 8.90 (1H, brs).

5 Example 144

(2S)-((3-((2R)-2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl)oxy)(phenyl)acetic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

10 ¹H-NMR (CD₃OD) δ: 1.33 (3H, d, J=6.5Hz), 3.02 (1H, dd, J=14.3, 8.8Hz), 3.23-3.37 (3H, m), 3.62-3.69 (1H, m), 5.13 (1H, dd, J=9.9, 3.1Hz), 5.93 (1H, s), 6.59 (1H, d, J=7.7Hz), 6.89 (1H, dd, J=8.0, 7.7Hz), 7.20 (1H, d, J=8.0Hz), 7.21 (1H, s), 7.33-7.43 (3H, m), 7.64-7.66 (2H, m), 7.78 (1H, dd, J=8.0, 4.3Hz), 8.24 (1H, d, J=8.0Hz), 8.68 (1H, d, J=4.3Hz), 8.75 (1H, s).

15 Reference Example 78

3-(2-Azidethyl)-7-(benzyloxy)-1H-indole

20 To a solution of 2-(7-(benzyloxy)-1H-indol-3-yl)ethanol (1.65 g, 6.17 mmol) in methylene chloride (20 mL) are added triethylamine (1.72 mL, 12.3 mmol) and methanesulfonyl chloride (0.6 mL, 7.78 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for 16 hours. The reaction solution is poured into water, and the mixture is extracted with chloroform. The organic layer is washed successively with a 1N aqueous hydrochloric acid solution, a
25 saturated aqueous sodium hydrogen carbonate solution and a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated, and the obtained crude product is dissolved in

N,N-dimethylformamide (20 mL), and thereto is added sodium azide (1.00 g, 15.4 mmol), and the mixture is reacted at 60°C for 3 hours. The reaction solution is poured into water, and the mixture is extracted with ethyl acetate. The organic layer is washed successively with a 1N aqueous hydrochloric acid solution, a saturated aqueous sodium hydrogen carbonate solution, and a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is removed by distillation, and the obtained crude product is purified by silica gel column chromatography (n-hexane:ethyl acetate = 10:1 → 3:1 → 1:1) to give the title compound (1.52 g, yield: 84 %).

¹H-NMR (CDCl₃) δ: 3.05 (2H, t, J=7.2Hz), 3.56 (2H, t, J=7.2Hz), 5.20 (2H, s), 6.74 (1H, d, J=7.7Hz), 7.02-7.06 (2H, m), 7.21 (1H, d, J=8.0Hz), 7.34-7.43 (3H, m), 7.47-7.49 (2H, m), 8.42 (1H, brs).

Reference Example 79

2-(7-(Benzyloxy)-1H-indol-3-yl)ethylamine

To a solution of 3-(2-azidethyl)-7-(benzyloxy)-1H-indole (5.10 g, 17.4 mmol) in pyridine (100 mL)-water (100 mL) is added triphenylphosphine (5.02 g, 19.1 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for 18 hours. The reaction solution is poured into water, and the mixture is extracted with chloroform. The organic layer is washed with a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated under reduced pressure, and the obtained crude product is purified by silica gel column chromatography (a saturated ammonia solution in chloroform → a saturated ammonia solution in chloroform-methanol = 50:1 → 10:1) to give the title compound (4.38 g, yield: 95 %).

¹H-NMR (CDCl₃) δ: 2.89 (2H, t, J=6.8Hz), 3.02 (2H, t, J=6.8Hz),

5.21 (2H, s), 6.73 (1H, d, J=7.7Hz), 7.01 (1H, d, J=2.1Hz), 7.02 (1H, dd, J=8.0, 7.7Hz), 7.24 (1H, d, J=8.0Hz), 7.34-7.43 (3H, m), 7.47-7.49 (2H, m), 8.29 (1H, brs).

Reference Example 80

5 (2R)-N-(2-(7-(Benzyloxy)-1H-indol-3-yl)ethyl)-2-hydroxy-2-pyridin-3-ylacetamide

The title compound is obtained in a similar manner to Reference Example 61.

10 $^1\text{H-NMR}$ (CDCl_3) δ : 2.82-2.97 (2H, m), 3.54-3.65 (2H, m), 4.99 (1H, s), 5.20 (2H, s), 6.37 (1H, t, J=5.3Hz), 6.73 (1H, d, J=7.6Hz), 6.80 (1H, d, J=2.2Hz), 7.02 (1H, dd, J=8.0, 7.6Hz), 7.16 (1H, d, J=8.0Hz), 7.22 (1H, dd, J=7.9, 4.8Hz), 7.33-7.42 (3H, m), 7.47-7.49 (2H, m), 7.65 (1H, dt, J=7.9, 1.8Hz), 8.44 (1H, brs), 8.47 (1H, dd, J=4.8, 1.6Hz), 8.52 (1H, d, J=2.1Hz).

15 Reference Example 81

tert-Butyl 2-(7-(benzyloxy)-1H-indol-3-yl)ethyl((2R)-2-hydroxy-2-pyridin-3-ylethyl)carbamate

The title compound is obtained in a similar manner to Reference Example 62 and Reference Example 63.

20 IR (ATR (total reflection absorption method)/FT-IR, cm^{-1}): 3320, 1670, 1577, 1411, 1365, 1257, 1226, 1161, 1045, 1026.

Reference Example 82

tert-Butyl 2-(7-(benzyloxy)-1H-indol-3-yl)ethyl((2R)-2-pyridin-3-yl-2-((triethylsilyl)oxy)ethyl)carbamate

25 The title compound is obtained in a similar manner to Reference Example 64.

IR (ATR (total reflection absorption method)/FT-IR, cm^{-1}): 1685,

1577, 1454, 1408, 1365, 1230, 1164, 1087.

Reference Example 83

tert-Butyl 2-(7-hydroxy-1H-indol-3-yl)ethyl((2R)-2-pyridin-3-yl-2-
((triethylsilyl)oxy)ethyl)carbamate

5 The title compound is obtained in a similar manner to
Reference Example 65.

IR (ATR (total reflection absorption method)/FT-IR, cm^{-1}): 3340,
1670, 1577, 1473, 1457, 1411, 1365, 1238, 1161, 1088.

Example 145

10 (1R)-2-((2-(7-(((1S)-1-Methyl-2-morpholin-4-yl-2-oxoethyl)oxy)-1H-indol-
3-yl)ethyl)amino)-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example
119.

15 $^1\text{H-NMR}$ (CDCl_3) δ : 1.65 (3H, d, $J=6.6\text{Hz}$), 2.69 (1H, dd, $J=12.2$,
9.3Hz), 2.93 (1H, dd, $J=12.2$, 3.4Hz), 2.94-3.08 (4H, m), 3.41-3.61 (8H,
m), 4.70 (1H, dd, $J=9.3$, 3.4Hz), 5.15 (1H, q, $J=6.6\text{Hz}$), 6.63 (1H, d,
 $J=7.7\text{Hz}$), 6.99 (1H, dd, $J=8.0$, 7.7Hz), 7.01 (1H, d, $J=1.8\text{Hz}$), 7.23-7.26
(2H, m), 7.69 (1H, d, $J=7.9\text{Hz}$), 8.49 (1H, dd, $J=4.8$, 1.3Hz), 8.55 (1H, d,
 $J=1.8\text{Hz}$), 8.83 (1H, brs).

20 Example 146

(2S)-2-((3-(2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)ethyl)-1H-indol-
7-yl)oxy)propanoic acid

The title compound is obtained in a similar manner to Example
120.

25 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.55 (3H, d, $J=6.6\text{Hz}$), 2.60-2.90 (5H, m),
3.02 (1H, dd, $J=12.3$, 3.3Hz), 4.60 (1H, q, $J=6.6\text{Hz}$), 4.93 (1H, dd, $J=9.9$,
3.3Hz), 6.50 (1H, d, $J=7.7\text{Hz}$), 6.70 (1H, dd, $J=7.9$, 7.7Hz), 6.78 (1H, d,

J=1.8Hz), 6.91 (1H, d, J=7.9Hz), 7.38 (1H, dd, J=7.7, 4.7Hz), 7.79 (1H, ddd, J=7.7, 1.8, 1.5Hz), 8.49 (1H, dd, J=4.7, 1.5Hz), 8.59 (1H, d, J=1.8Hz), 10.87 (1H, d, J=1.6Hz).

Example 147

5 (1R)-2-((2-(7-(((1S)-2-Methyl-1-(morpholin-4-ylcarbonyl)propyl)oxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

10 ¹H-NMR (CDCl₃) δ: 1.11 (3H, d, J=6.8Hz), 1.20 (3H, d, J=6.6Hz), 2.23-2.32 (1H, m), 2.70 (1H, dd, J=12.3, 9.5Hz), 2.95 (1H, dd, J=12.3, 3.5Hz), 2.96-3.07 (3H, m), 3.36-3.37 (2H, m), 3.48-3.67 (6H, m), 3.76-3.79 (1H, m), 4.62 (1H, d, J=7.8Hz), 4.73 (1H, dd, J=9.5, 3.5Hz), 6.67 (1H, d, J=7.7Hz), 6.99 (1H, dd, J=8.0, 7.7Hz), 7.03 (1H, d, J=1.9Hz), 7.25 (1H, dd, J=7.9, 4.8Hz), 7.25 (1H, d, J=8.0Hz), 7.68 (1H, ddd, J=7.9, 2.0, 1.6Hz), 8.49 (1H, dd, J=4.8, 1.6Hz), 8.55 (1H, d, J=2.0Hz), 8.69 (1H, brs).

Example 148

(2S)-2-((3-(2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)ethyl)-1H-indol-7-yl)oxy)-3-methylbutanoic acid · 2 trifluoroacetate

20 The title compound is obtained in a similar manner to Example 120.

25 ¹H-NMR (CD₃OD) δ: 1.19 (3H, d, J=6.9Hz), 1.24 (3H, d, J=6.9Hz), 2.35-2.45 (1H, m), 3.23-3.31 (3H, m), 3.41-3.47 (3H, m), 4.66 (1H, d, J=5.1Hz), 5.28 (1H, dd, J=9.9, 3.0Hz), 6.58 (1H, d, J=7.7Hz), 6.99 (1H, dd, J=7.9, 7.7Hz), 7.22 (1H, s), 7.26 (1H, d, J=7.9Hz), 8.04 (1H, brt, J=6Hz), 8.50 (1H, d, J=8.1Hz), 8.84-8.90 (2H, m).

Reference Example 84

2-(7-(((1S)-1-Methyl-2-morpholin-4-yl-2-oxoethyl)oxy)-1H-indol-3-yl)-ethanol

To a solution of 3-(2-hydroxyethyl)-1H-indol-7-ol (1.05 g, 5.93 mmol) in N,N-dimethylformamide (20 mL) are added (1R)-1-methyl-2-morpholin-4-yl-2-oxoethyl 4-methylbenzenesulfonate (2.79 g, 8.90 mmol) and potassium carbonate (1.3 g, 9.41 mmol), and the mixture is stirred at 60°C for 5 hours. The reaction solution is cooled to room temperature, and the mixture is poured into water. The mixture is extracted with ethyl acetate, and the organic layer is washed with a saturated aqueous sodium hydrogen carbonate solution, and dried over anhydrous sodium sulfate. The solvent is removed by distillation, and the obtained crude product is purified by silica gel column chromatography (chloroform → chloroform:methanol = 50:1 → 30:1) to give the title compound (1.29 g, yield: 68 %).

¹H-NMR (DMSO-d₆) δ: 1.63 (1H, t, J=6.1Hz), 1.66 (3H, d, J=6.7Hz), 3.02 (2H, td, J=6.3, 0.4Hz), 3.43-3.74 (8H, m), 3.90 (1H, td, J=6.3, 6.1Hz), 5.15 (1H, q, J=6.7Hz), 6.64 (1H, d, J=7.7Hz), 7.00 (1H, dd, J=7.9, 7.7Hz), 7.07 (1H, d, J=2.3Hz), 7.29 (1H, d, J=7.9Hz), 8.73 (1H, brs).

Reference Example 85

(7-(((1S)-1-Methyl-2-morpholin-4-yl-2-oxoethyl)oxy)-1H-indol-3-yl)acetaldehyde

The title compound is obtained in a similar manner to Reference Example 24.

¹H-NMR (CDCl₃) δ: 1.65 (3H, d, J=6.7Hz), 3.42-3.62 (8H, m), 3.78 (2H, dd, J=2.5, 0.7Hz), 5.15 (1H, q, J=6.7Hz), 7.02 (1H, dd, J=8.0, 7.7Hz), 7.12 (1H, d, J=2.4Hz), 7.18 (1H, d, J=8.0Hz), 9.00 (1H, brs),

9.74 (1H, t, J=2.5Hz).

Example 149

4-((1R,2S)-1-Hydroxy-2-{{2-(7-(((1S)-1-methyl-2-morpholin-4-yl-2-oxoethyl)oxy)-1H-indol-3-yl)ethyl}amino}propyl)phenol

5 The title compound is obtained in a similar manner to Example 30.

¹H-NMR (CDCl₃) δ: 1.18 (3H, d, J=6.3Hz), 1.68 (3H, d, J=6.7Hz), 2.59-2.65 (2H, m), 2.74-2.81 (1H, m), 2.86-2.92 (1H, m), 3.00-3.05 (1H, m), 3.61-3.83 (8H, m), 4.19 (1H, d, J=7.8Hz), 5.19 (1H, q, J=6.7Hz),
10 6.30 (2H, d, J=8.5Hz), 6.32 (1H, d, J=7.5Hz), 6.64 (1H, d, J=2.2Hz), 6.75 (2H, d, J=8.5Hz), 6.90 (1H, dd, J=8.0, 7.6Hz), 7.18 (1H, d, J=8.0Hz), 8.59 (1H, brs).

Example 150

15 (2S)-2-{{3-(2-(((1S,2R)-2-Hydroxy-2-(4-hydroxyphenyl)-1-methylethyl)-amino)ethyl)-1H-indol-7-yl}oxy}propanoic acid

 The title compound is obtained in a similar manner to Example 120.

¹H-NMR (DMSO-d₆) δ: 0.89 (3H, d, J=6.6Hz), 1.54 (3H, d, J=6.8Hz), 2.49-2.53 (1H, m), 2.62-2.75 (1H, m), 2.94-3.04 (2H, m), 3.16
20 (1H, brs), 4.58 (1H, q, J=6.8Hz), 4.99 (1H, brs), 6.51 (1H, d, J=7.6Hz), 6.67 (2H, d, J=8.3Hz), 6.73 (1H, s), 6.75 (1H, dd, J=7.9, 7.6Hz), 6.90 (1H, d, J=7.9Hz), 7.08 (2H, d, J=8.3Hz), 9.49 (1H, brs), 10.82 (1H, s).

Reference Example 86

25 2-(7-(((1R)-1-Methyl-2-morpholin-4-yl-2-oxoethyl)oxy)-1H-indol-3-yl)ethanol

 The title compound is obtained in a similar manner to Reference Example 84.

¹H-NMR (CDCl₃) δ: 1.63 (1H, t, J=6.1Hz), 1.65 (3H, d, J=6.7Hz), 3.02 (2H, t, J=6.2Hz), 3.43-3.63 (m, 8H), 3.90 (2H, td, J=6.2, 6.1Hz), 5.14 (1H, q, J=6.7Hz), 6.64 (1H, d, J=7.7Hz), 7.00 (1H, t, J=7.8Hz), 7.06 (1H, d, J=2.2Hz), 7.27 (1H, d, J=7.4Hz), 8.70 (brs, 1H).

5 Reference Example 87

(7-(((1R)-1-Methyl-2-morpholin-4-yl-2-oxoethyl)oxy)-1H-indol-3-yl)acetaldehyde

The title compound is obtained in a similar manner to Reference Example 24.

10 ¹H-NMR (CDCl₃) δ: 1.66 (3H, d, J=6.7Hz), 3.43-3.63 (8H, m), 3.79 (2H, dd, J=2.4, 0.65Hz), 5.15 (1H, q, J=6.7Hz), 6.67 (1H, d, J=7.7Hz), 7.03 (1H, t, J=7.8Hz), 7.16 (1H, d, J=2.4Hz), 7.19 (1H, d, J=8.0Hz), 9.76 (t, 1H, J=2.5Hz).

Example 151

15 4-(((1R,2S)-1-Hydroxy-2-((2-(7-(((1R)-1-methyl-2-morpholin-4-yl-2-oxoethyl)oxy)-1H-indol-3-yl)ethyl)amino)propyl)phenol

The title compound is obtained in a similar manner to Example 30.

20 ¹H-NMR (CDCl₃) δ: 1.24 (3H, d, J=6.1Hz), 1.69 (3H, d, J=6.7Hz), 2.47-2.54 (1H, m), 2.61-2.73 (2H, m), 2.61-2.73 (2H, m), 2.97-3.03 (2H, m), 3.65-3.91 (8H, m), 4.12 (1H, d, J=8.5Hz), 5.22 (1H, q, J=6.7Hz), 6.25 (1H, d, J=7.6Hz), 6.27-6.31 (2H, m), 6.60 (1H, d, J=7.6Hz), 6.66-6.70 (2H, m), 6.91 (1H, t, J=7.8Hz), 7.13 (1H, d, J=8.0Hz), 8.26 (brs, 1H).

25 Example 152

(2R)-2-((3-(2-(((1S,2R)-2-Hydroxy-2-(4-hydroxyphenyl)-1-methylethyl)amino)ethyl)-1H-indol-7-yl)oxy)propanoic acid

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CD₃OD) δ: 1.07 (3H, d, J=6.7Hz), 1.68 (3H, d, J=6.8Hz), 3.17-3.20 (2H, m), 3.35-3.42 (3H, m), 4.91-5.00 (2H, m), 6.59 (1H, d, J=7.7Hz), 6.76 (2H, d, J=8.6Hz), 6.96 (1H, t, J=7.9Hz), 7.11 (2H, d, J=8.6Hz), 7.18 (1H, s), 7.23 (1H, d, J=7.9Hz).

Reference Example 88

tert-Butyl (1R)-2-[7-(benzyloxy)-1H-indol-3-yl]-1-methylethylcarbamate

To a solution of (1R)-2-[7-(benzyloxy)-1H-indol-3-yl]-1-methylethylamine (3.09 g, 11.0 mmol) in tetrahydrofuran (100 mL) is added di-tert-butyl bicarbonate (3.80 mL, 16.5 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for 60 hours. The reaction solution is poured into a mixture of a saturated brine and water (1:1), and the mixture is extracted with ethyl acetate. The organic layer is washed with a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated, and the obtained crude product is purified by silica gel column chromatography (n-hexane:ethyl acetate = 3:1) to give the title compound (3.70 g, yield: 88 %).

¹H-NMR (CDCl₃) δ: 1.11 (3H, d, J=6.4Hz), 1.43 (9H, s), 2.84 (1H, dd, J=14.8, 7.2Hz), 2.95 (1H, dd, J=14.8, 5.2Hz), 4.01 (1H, brs), 4.45 (1H, brs), 5.20 (2H, s), 6.72 (1H, d, J=7.2Hz), 6.99 (1H, d, J=2.4Hz), 7.02 (1H, dd, J=7.6, 7.2Hz), 7.34-7.43 (3H, m), 7.46-7.49 (2H, m), 8.28 (1H, brs).

Reference Example 89

tert-Butyl (1R)-1-methyl-2-(7-([(1S)-1-methyl-2-morpholin-4-yl-2-oxoethyl]oxy)-1H-indol-3-yl)ethylcarbamate

To a solution of tert-butyl (1R)-2-[7-(benzyloxy)-1H-indol-3-yl]-1-methylethylcarbamate (1.0 g, 2.63 mmol) in methanol (30 mL) is added a 10 % palladium on carbon (50 % wet, 1.0 g), and the mixture is stirred under hydrogen atmosphere for one hour. The reaction solution is filtered through celite, and the filtrate is evaporated under reduced pressure to remove the solvent. The resulting residue is dissolved in N,N-dimethylformamide (5 mL), and thereto are added potassium carbonate (581 mg, 4.21 mmol) and (1R)-1-methyl-2-morpholin-4-yl-2-oxoethyl 4-methylbenzenesulfonate (1.24 g, 3.95 mmol), and the mixture is stirred at 55°C. The reaction solution is cooled to room temperature, and the mixture is poured into a mixture of a saturated brine and water (1:1), and extracted with ethyl acetate. The organic layer is washed with a saturated aqueous sodium hydrogen carbonate solution, and dried over anhydrous potassium carbonate. The solvent is distilled off, and the obtained crude product is purified by silica gel column chromatography (chloroform → chloroform:methanol = 100:1) to give the title compound (1.03 g, yield: 63 %).

¹H-NMR (CDCl₃) δ: 1.11 (3H, d, J=6.4Hz), 1.43 (9H, s), 1.66 (3H, d, J=6.8Hz), 2.83 (1H, dd, J=14, 6.8Hz), 2.94 (1H, dd, J=5.2Hz), 3.39-3.51 (2H, m), 3.55-3.65 (6H, m), 4.00 (1H, brs), 4.45 (1H, brs), 5.13 (1H, q, J=6.8Hz), 6.63 (1H, d, J=7.6Hz), 6.99 (1H, dd, J=8.0, 7.6Hz), 7.01 (1H, s), 7.29 (1H, d, J=8Hz), 8.55 (1H, brs).

Reference Example 90

(1R)-1-Methyl-2-(7-(((1S)-1-methyl-2-morpholin-4-yl-2-oxoethyl)oxy))-1H-indol-3-yl)ethylamine

To a solution of tert-butyl (1R)-1-methyl-2-(7-(((1S)-1-methyl-2-morpholin-4-yl-2-oxoethyl)oxy))-1H-indol-3-yl)ethylcarbamate (1.03 g,

2.39 mmol) in acetonitrile (10 mL) is added oxalic acid (860 mg, 9.55 mmol), and the mixture is refluxed for 5 hours. The reaction solution is evaporated under reduced pressure to remove the solvent, and to the residue are added a 10 % aqueous potassium carbonate solution and a mixture of chloroform:methanol (10:1) for separation. The organic layer is dried over anhydrous potassium carbonate, and the solvent is removed by distillation to give the title compound (626 mg, yield: 79 %).

¹H-NMR (CDCl₃) δ: 1.16 (3H, d, J=6.4Hz), 1.66 (3H, d, J=6.8Hz), 2.63 (1H, dd, J=8.4, 14.8Hz), 2.86 (1H, ddd, J=0.8, 4.8, 14.4Hz), 3.25-3.30 (1H, m), 3.40-3.52 (2H, m), 3.58-3.62 (6H, m), 5.13 (1H, q, J=6.8Hz), 6.64 (1H, d, J=7.6Hz), 6.99 (1H, dd, J=8.0, 8.0Hz), 7.27 (1H, d, J=8.0Hz), 8.55 (1H, brs).

Example 153

(1R)-1-(3-Chlorophenyl)-2-{{{(1R)-1-methyl-2-(7-{{{(1S)-1-methyl-2-morpholin-4-yl-2-oxoethyl}oxy}-1H-indol-3-yl)ethyl}amino}ethanol · hydrochloride

The title compound is obtained in a similar manner to Example 108.

¹H-NMR (CD₃OD) δ: 1.31 (3H, d, J=6.5Hz), 1.63 (3H, d, J=6.6Hz), 3.00 (1H, dd, J=9.0, 14.3Hz), 3.16-3.27 (3H, m), 3.40-3.43 (1H, m), 3.57-3.72 (8H, m), 4.93 (1H, dd, J=3.3, 9.9Hz), 5.35 (1H, q, J=6.6Hz), 6.62 (1H, d, J=8.0Hz), 6.96 (1H, dd, J=7.9, 7.9Hz), 7.18 (1H, s), 7.23 (1H, d, J=8.0Hz), 7.29-7.40 (3H, m), 7.45 (1H, s).

Example 154

(2S)-2-{{[3-{{(2R)-2-{{(2R)-2-(3-Chlorophenyl)-2-hydroxyethyl}amino}propyl)-1H-indol-7-yl]oxy}propanoic acid

The title compound is obtained in a similar manner to Example

120.

¹H-NMR (CD₃OD) δ: 1.16 (3H, d, J=6.4Hz), 1.60 (3H, d, J=6.7Hz), 2.78-2.98 (4H, m), 3.21-3.26 (1H, m), 4.70-4.76 (2H, m), 6.57 (1H, d, J=7.7Hz), 6.86 (1H, dd, J=7.8, 7.9Hz), 7.00 (1H, s), 7.07 (1H, d, J=7.7Hz), 7.14-7.16 (1H, m), 7.22-7.25 (2H, m), 7.34 (1H, s).

Reference Example 91

tert-Butyl (1R)-1-methyl-2-(7-(((1R)-1-methyl-2-morpholin-4-yl-2-oxoethyl]oxy)-1H-indol-3-yl)ethylcarbamate

The title compound is obtained in a similar manner to

Reference Example 89.

¹H-NMR (CDCl₃) δ: 1.11 (3H, d, J=6.6Hz), 1.44 (9H, s), 1.66 (3H, d, J=6.7Hz), 2.83 (1H, dd, J=14, 6.9Hz), 2.94 (1H, dd, J=14, 5.2Hz), 3.48-3.57 (8H, m), 4.40 (1H, m), 4.38-4.48 (1H, m), 5.14 (1H, q, J=6.7Hz), 6.63 (1H, d, J=7.7Hz), 6.99 (1H, t, J=7.9Hz), 7.01 (1H, d, J=2.1Hz), 7.29 (1H, d, J=8.0Hz), 8.57 (1H, brs).

Reference Example 92

(1R)-1-Methyl-2-(7-(((1R)-1-methyl-2-morpholin-4-yl-2-oxoethyl]oxy)-1H-indol-3-yl)ethylamine

The title compound is obtained in a similar manner to

Reference Example 90.

¹H-NMR (CDCl₃) δ: 1.16 (3H, d, J=6.3Hz), 1.66 (3H, d, J=6.7Hz), 2.64 (1H, dd, J=14, 8.2Hz), 2.85 (1H, dd, J=14, 5.0Hz), 3.23-3.31 (1H, m), 3.41-3.62 (8H, m), 5.14 (1H, q, J=6.7Hz), 6.64 (1H, d, J=7.7Hz), 6.99 (1H, t, J=7.8Hz), 7.04 (1H, d, J=2.1Hz), 7.27 (1H, d, J=7.9Hz), 8.55 (1H, s).

Example 155

(1R)-1-(3-Chlorophenyl)-2-(((1R)-1-methyl-2-(7-(((1R)-1-methyl-2-

morpholin-4-yl-2-oxoethyl]oxy}-1H-indol-3-yl)ethyl]amino}ethanol

The title compound is obtained in a similar manner to Example 108.

¹H-NMR (CDCl₃) δ: 1.11 (3H, d, J=6.2Hz), 1.63 (3H, d, J=6.6Hz),
 2.66 (1H, dd, J=12, 9.1Hz), 2.75-2.84 (2H, m), 2.85 (1H, dd, J=12,
 3.6Hz), 3.00-3.05 (1H, m), 3.39-3.59 (8H, m), 4.51 (1H, dd, J=9.0,
 3.4Hz), 5.14 (1H, q, J=6.7Hz), 6.60 (1H, d, J=7.7Hz), 6.97 (1H, s), 6.98
 (1H, t, J=7.9Hz), 7.14-7.24 (4H, m), 7.32 (1H, s), 8.92 (brs, 1H).

Example 156

(2R)-2-{{3-((2R)-2-{{(2R)-2-(3-Chlorophenyl)-2-hydroxyethyl]amino}-
 propyl)-1H-indol-7-yl]oxy}propanoic acid

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (DMSO-d₆) δ: 0.91 (3H, d, J=6.2Hz), 1.55 (3H, d,
 J=6.7Hz), 2.50-2.55 (1H, m), 2.77-2.85 (2H, m), 2.94-2.97 (1H, m), 3.04
 (2H, m), 4.65-4.70 (1H, m), 4.79-4.82 (1H, m), 6.76 (1H, t, J=7.8Hz),
 6.91 (1H, s), 7.00 (1H, d, J=8.0Hz), 7.30-7.38 (3H, m), 7.43 (1H, s), 7.48
 (1H, d, J=7.7Hz), 10.9 (s, 1H).

Reference Example 93

(1S)-2-Morpholin-4-yl-2-oxo-1-phenylethanol

The title compound is obtained in a similar manner to Reference Example 68.

¹H-NMR (CDCl₃) δ: 3.05-3.10 (1H, m), 3.14-3.19 (1H, m), 3.28-
 3.33 (1H, m), 3.45-3.50 (1H, m), 3.54-3.83 (4H, m), 4.70 (1H, brs), 5.19
 (1H, s), 7.30-7.41 (5H, m).

Reference Example 94

(1S)-2-Morpholin-4-yl-2-oxo-1-phenylethyl 4-methylbenzenesulfonate

The title compound is obtained in a similar manner to Reference Example 69.

¹H-NMR (CDCl₃) δ: 2.43 (3H, s), 3.35-3.61 (8H, m), 6.12 (1H, s), 7.31 (2H, d, J=8.4Hz), 7.34 (5H, s), 7.81 (2H, d, J=8.4Hz).

5 Example 157

(2R)-{[3-((2R)-2-((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl]oxy}(phenyl)acetic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 119 and Example 120.

10 ¹H-NMR (CD₃OD) δ: 1.33 (3H, d, J=6.5Hz), 3.20 (1H, dd, J=14, 9.0Hz), 3.25-3.31 (2H, m), 3.39 (1H, ddd, J=13, 4.1, 3.4Hz), 3.62-3.70 (1H, m), 5.21 (1H, dd, J=9.8, 2.9Hz), 5.93 (1H, s), 6.59 (1H, d, J=7.9Hz), 6.87 (1H, t, J=7.9Hz), 7.19-7.21 (1H, m), 7.21 (1H, s), 7.33-7.45 (3H, m), 7.64-7.66 (2H, m), 7.93 (1H, dd, J=7.4, 6.0Hz), 8.43 (1H, d, J=7.9Hz),
15 8.76 (1H, d, J=4.2Hz), 8.84 (1H, s).

Reference Example 95

(2S)-1-Morpholin-4-yl-1-oxo-3-phenylpropan-2-ol

The title compound is obtained in a similar manner to Reference Example 68.

20 ¹H-NMR (CDCl₃) δ: 2.90 (1H, dd, J=14, 6.7Hz), 2.96 (1H, dd, J=14, 6.7Hz), 3.27-3.30 (2H, m), 3.51-3.68 (6H, m), 4.59 (1H, dd, J=14, 6.7Hz), 7.21-7.33 (5H, m).

Reference Example 96

(1S)-1-Benzyl-2-morpholine-4-yl-2-oxoethyl 4-methylbenzenesulfonate

25 The title compound is obtained in a similar manner to Reference Example 69.

¹H-NMR (CDCl₃) δ: 2.43 (3H, s), 2.97-3.02 (1H, m), 3.07-3.13

(2H, m), 3.16-3.27 (2H, m), 3.30-3.43 (3H, m), 3.48-3.59 (2H, m), 5.26 (1H, dd, J=8.1, 6.9Hz), 7.12-7.14 (2H, m), 7.24-7.29 (5H, m), 7.68 (2H, d, J=8.3Hz).

Example 158

5 (1R)-2-[[[(1R)-2-(7-[[[(1R)-1-Benzyl-2-morpholin-4-yl-2-oxoethyl]oxy]-1H-indol-3-yl)-1-methylethyl]amino]-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

10 ¹H-NMR (CD₃OD) δ: 1.14 (3H, d, J=6.2Hz), 2.75-2.92 (5H, m), 3.02-3.12 (1H, m), 3.24-3.56 (9H, m), 4.71 (1H, dd, J=8.1, 4.7Hz), 5.43 (1H, t, J=7.0Hz), 6.53 (1H, d, J=7.7Hz), 6.88 (1H, dd, J=8.0, 7.7Hz), 7.02 (1H, s), 7.15 (1H, d, J=8.0Hz), 7.23-7.27 (2H, m), 7.30-7.34 (2H, m), 7.36-7.38 (2H, m), 7.64 (1H, d, J=7.9Hz), 7.88 (1H, s), 8.33 (1H, d, J=4.8Hz), 8.44 (1H, d, J=1.8Hz).

15 Reference Example 97

(1R)-1-Cyclohexyl-2-morpholin-4-yl-2-oxoethanol

The title compound is obtained in a similar manner to Reference Example 68.

20 ¹H-NMR (CDCl₃) δ: 1.14-1.22 (5H, m), 1.41-1.44 (3H, m), 1.64-1.67 (1H, m), 1.78-1.81 (2H, m), 3.43-3.44 (2H, m), 3.63-3.74 (6H, m), 4.19 (1H, brs).

Reference Example 98

(1R)-1-Cyclohexyl-2-morpholin-4-yl-2-oxoethyl 4-methylbenzene-sulfonate

25 The title compound is obtained in a similar manner to Reference Example 69.

¹H-NMR (CDCl₃) δ: 0.90-1.27 (5H, m), 1.49-1.91 (6H, m), 2.45

(3H, s), 3.43-3.70 (8H, m), 4.79 (1H, d, J=8.7Hz), 7.34 (2H, d, J=8.2Hz), 7.80 (2H, d, J=8.2Hz).

Example 159

(1R)-2-(((1R)-2-(7-(((1S)-1-Cyclohexyl-2-morpholin-4-yl-2-oxoethyl)oxy)-1H-indol-3-yl)-1-methylethyl)amino)-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.17 (3H, d, J=6.1Hz), 1.10-1.31 (6H, m), 1.71-1.74 (2H, m), 1.81-1.84 (2H, m), 1.97-2.05 (1H, m), 2.14-2.21 (1H, m), 2.70-2.75 (1H, m), 2.87 (1H, d, J=6.2Hz), 2.93 (1H, dd, J=12, 2.9Hz), 3.11-3.14 (1H, m), 3.35 (2H, m), 3.51-3.70 (6H, m), 4.63 (1H, d, J=6.9Hz), 4.68 (1H, d, J=8.0Hz), 6.66 (1H, d, J=7.7Hz), 6.98 (1H, t, J=7.8Hz), 7.03 (1H, s), 7.22 (1H, d, J=8.2Hz), 7.23 (1H, d, J=7.4Hz), 7.66 (1H, d, J=7.7Hz), 8.49 (1H, d, J=3.9Hz), 8.52 (1H, s), 8.63 (1H, brs).

Example 160

(2S)-Cyclohexyl{[3-((2R)-2-(((2R)-2-hydroxy-2-pyridin-3-ylethyl)amino)-propyl)-1H-indol-7-yl]oxy}acetic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CD₃OD) δ: 1.33 (3H, d, J=6.5Hz), 1.29-1.49 (6H, m), 1.71-1.74 (1H, m), 1.83 (3H, m), 1.95-2.05 (2H, m), 3.01 (1H, dd, J=14, 9.0Hz), 3.25-3.32 (1H, m), 3.37 (1H, dd, J=13, 3.3Hz), 3.66-3.68 (1H, m), 4.60 (1H, d, J=5.4Hz), 5.16 (1H, dd, J=9.9, 3.1Hz), 6.53 (1H, d, J=7.7Hz), 6.94 (1H, t, J=7.9Hz), 7.18 (1H, s), 7.21 (1H, d, J=7.8Hz), 7.83 (1H, dd, J=7.9, 5.5Hz), 8.29 (1H, d, J=8.1Hz), 8.71 (1H, d, J=4.4Hz), 8.78 (1H, brs).

Reference Example 99

(2R)-1-Morpholin-4-yl-1-oxobutan-2-ol

The title compound is obtained in a similar manner to Reference Example 68.

¹H-NMR (CDCl₃) δ: 1.00 (3H, t, J=7.4Hz), 1.46-1.56 (1H, m),
 1.65-1.73 (1H, m), 3.41-3.43 (2H, m), 3.63-3.74 (6H, m), 4.28-4.32 (1H, m).

Reference Example 100

(1R)-1-(Morpholin-4-ylcarbonyl)propyl 4-methylbenzenesulfonate

The title compound is obtained in a similar manner to Reference Example 69.

¹H-NMR (CDCl₃) δ: 0.93 (3H, t, J=7.4Hz), 1.81-1.88 (2H, m),
 2.45 (3H, s), 3.43-3.64 (8H, m), 5.00 (1H, t, J=6.7Hz), 7.35 (2H, d, J=8.0Hz), 7.81 (2H, d, J=8.3Hz).

Example 161

(1R)-2-{{{(1R)-1-Methyl-2-(7-(((1S)-1-(morpholin-4-ylcarbonyl)propyl)oxy)-1H-indol-3-yl)ethyl)amino}-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.14-1.18 (6H, m), 2.02-2.07 (2H, m), 2.65 (1H, dd, J=9.4, 12.2Hz), 2.81-2.92 (2H, m), 3.04-3.07 (1H, m), 3.39-3.49 (2H, m), 3.57-3.63 (6H, m), 4.55 (1H, dd, J=3.5, 9.3Hz), 4.88 (1H, dd, J=5.8, 7.6Hz), 6.65 (1H, d, J=7.7Hz), 7.00 (1H, dd, J=7.8, 7.9Hz), 7.03 (1H, d, J=2.0Hz), 7.23-7.26 (2H, m), 7.66-7.68 (1H, m), 8.50 (1H, dd, J=1.6, 4.8Hz), 8.54 (1H, d, J=2.0Hz), 8.58 (1H, brs).

Example 162

(2S)-2-[[3-(((2R)-2-((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl]oxy]butanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CD₃OD) δ: 1.13 (3H, t, J=7.5Hz), 1.32 (3H, d, J=6.5Hz), 2.01-2.13 (2H, m), 3.00 (1H, dd, J=9.1, 14.3Hz), 3.25-3.32 (2H, m), 3.63-3.68 (1H, m), 4.80 (1H, dd, J=5.2, 6.7Hz), 5.26 (1H, dd, J=3.0, 9.7Hz), 6.55 (1H, d, J=7.7Hz), 6.94 (1H, dd, J=7.9, 7.9Hz), 7.18 (1H, s), 7.22 (1H, d, J=7.8Hz), 8.00 (1H, dd, J=5.7, 8.0Hz), 8.51 (1H, d, J=8.2Hz), 8.78 (1H, d, J=4.9Hz), 8.88 (1H, brs).

Reference Example 101

(2R)-1-Morpholin-4-yl-1-oxopentane-2-ol

The title compound is obtained in a similar manner to Reference Example 68.

¹H-NMR (CDCl₃) δ: 0.95 (3H, t, J=7.2Hz), 1.47-1.58 (4H, m), 3.40-3.42 (2H, m), 3.61-3.74 (6H, m), 4.31-4.36 (1H, m).

Reference Example 102

(1R)-1-(Morpholin-4-ylcarbonyl)butyl 4-methylbenzenesulfonate

The title compound is obtained in a similar manner to Reference Example 69.

¹H-NMR (CDCl₃) δ: 0.87 (3H, t, J=7.4Hz), 1.24-1.33 (1H, m), 1.39-1.48 (1H, m), 1.69-1.87 (2H, m), 2.45 (3H, s), 3.41-3.64 (8H, m), 5.06 (1H, dd, J=5.3, 8.7Hz), 7.35 (2H, d, J=8.4Hz), 7.81 (2H, d, J=8.3Hz).

Example 163

(1R)-2-(((1R)-1-Methyl-2-(7-(((1S)-1-(morpholin-4-ylcarbonyl)butyl)oxy)-1H-indol-3-yl)ethyl)amino)-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.03 (3H, t, J=7.4Hz), 1.14 (3H, d, J=6.2Hz), 1.58-1.72 (2H, m), 1.89-2.05 (2H, m), 2.65 (1H, dd, J=9.3, 12.2Hz), 2.79-2.92 (3H, m), 3.03-3.08 (1H, m), 3.38-3.47 (2H, m), 3.58-3.62 (6H, m), 4.55 (1H, dd, J=3.5, 9.2Hz), 4.95 (1H, dd, J=4.8, 8.6Hz), 6.64 (1H, d, J=7.6Hz), 6.99 (1H, dd, J=7.8, 7.9Hz), 7.02 (1H, d, J=1.9Hz), 7.23-7.26 (2H, m), 7.65-7.67 (1H, m), 8.50 (1H, dd, J=1.6, 4.8Hz), 8.54 (2H, m).

Example 164

(2S)-2-([3-((2R)-2-((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)propyl)-1H-indol-7-yl]oxy}pentanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CD₃OD) δ: 0.99 (3H, t, J=7.4Hz), 1.32 (3H, d, J=6.5Hz), 1.57-1.67 (2H, m), 1.98-2.05 (2H, m), 3.99 (1H, dd, J=9.0, 14.3Hz), 3.25-3.32 (2H, m), 3.42 (1H, dd, J=3.2, 12.7Hz), 3.63-3.69 (1H, m), 4.84 (1H, dd, J=5.3, 7.0Hz), 5.25 (1H, dd, J=3.0, 9.7Hz), 6.54 (1H, d, J=7.7Hz), 6.94 (1H, dd, J=7.9, 7.9Hz), 7.17 (1H, s), 7.21 (1H, d, J=7.9Hz), 7.99 (1H, dd, J=5.7, 8.0Hz), 8.50 (1H, d, J=8.2Hz), 8.78 (1H, d, J=5.2Hz), 8.87 (1H, brs).

Example 165

(1R)-2-([2-(7-([(1R)-1-Methyl-2-morpholin-4-yl-2-oxoethyl]oxy)-1H-indol-3-yl)ethyl]amino)-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.64 (3H, d, J=6.7Hz), 2.67-2.72 (1H, m), 2.89-3.04 (5H, m), 3.42-3.66 (8H, m), 4.70 (1H, m), 5.13 (1H, q, J=6.7Hz), 6.62 (1H, d, J=7.6Hz), 6.97-7.01 (2H, m), 7.23-7.26 (2H, m), 7.68 (1H, d, J=7.8Hz), 8.48 (1H, d, J=4.6Hz), 8.55 (1H, s), 8.91 (1H, brs).

Example 166

(2R)-2-[[3-(2-[[[(2R)-2-Hydroxy-2-pyridin-3-ylethyl]amino]ethyl]-1H-indol-7-yl]oxy}propanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CD₃OD) δ: 1.67 (3H, d, J=6.8Hz), 3.18-3.28 (3H, m), 3.35-3.41 (3H, m), 4.93-5.00 (1H, m), 5.23 (1H, dd, J=9.7, 2.7Hz), 6.58 (1H, d, J=7.7Hz), 6.95 (1H, dd, J=7.9, 7.7Hz), 7.17 (1H, s), 7.72 (1H, d, J=7.9Hz), 7.96 (1H, brs), 8.45 (1H, d, J=8.1Hz), 8.77 (1H, brs), 8.85 (1H, brs).

Example 167

(1R)-2-[[2-(7-[[[(1R)-2-Methyl-1-(morpholin-4-ylcarbonyl)propyl]oxy]-1H-indol-3-yl]ethyl]amino]-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.10 (3H, d, J=6.8Hz), 1.19 (3H, d, J=6.8Hz), 2.23-2.32 (1H, m), 2.68 (1H, dd, J=12.3, 9.3Hz), 2.88-3.06 (5H, m), 3.33-3.36 (2H, m), 3.47-3.68 (6H, m), 4.62 (1H, d, J=7.8Hz), 4.69 (1H, dd, J=9.3, 3.5Hz), 6.66 (1H, d, J=7.7Hz), 6.97-7.00 (2H, m), 7.23-7.26 (2H, m), 7.68 (1H, ddd, J=7.9, 2.0, 1.6Hz), 8.48 (1H, dd, J=4.8, 1.6Hz), 8.55 (1H, d, J=2.0Hz), 8.78 (1H, brs).

Example 168

(2R)-2-[[3-(2-[[[(2R)-2-Hydroxy-2-pyridin-3-ylethyl]amino]ethyl]-1H-indol-7-yl]oxy}-3-methylbutanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CD₃OD) δ: 1.14 (3H, d, J=6.8Hz), 1.18 (3H, d,

J=6.8Hz), 2.32-2.40 (1H, m), 3.18-3.27 (3H, m), 3.32-3.40 (3H, m), 4.60 (1H, d, J=5.1Hz), 5.21-5.23 (1H, m), 6.53 (1H, d, J=7.7Hz), 6.93 (1H, dd, J=7.9, 7.7Hz), 7.18 (1H, s), 7.20 (1H, d, J=7.9Hz), 7.96 (1H, brs), 8.44 (1H, brs), 8.77 (1H, brs), 8.85 (1H, brs).

5 Example 169

(1R)-2-{{2-(7-{{(1S)-1-Benzyl-2-morpholin-4-yl-2-oxoethyl}oxy}-1H-indol-3-yl)ethyl}amino}-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

10 ¹H-NMR (CDCl₃) δ: 2.68 (1H, dd, J=12.2, 9.3Hz), 2.89-3.09 (6H, m), 3.28-3.36 (5H, m), 3.51-3.60 (4H, m), 4.70 (1H, dd, J=9.3, 3.4Hz), 5.15 (1H, t, J=6.9Hz), 6.56 (1H, d, J=7.6Hz), 6.93-6.97 (2H, m), 7.22-7.36 (6H, m), 7.68 (1H, ddd, J=7.9, 2.1, 1.6Hz), 8.49 (1H, dd, J=4.8, 1.6Hz), 8.55 (1H, d, J=2.1Hz), 8.60 (1H, brs).

15 Example 170

(2S)-2-{{3-(2-{{(2R)-2-Hydroxy-2-pyridin-3-ylethyl}amino}ethyl)-1H-indol-7-yl}oxy}-3-phenylpropanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

20 ¹H-NMR (CD₃OD) δ: 3.17-3.48 (8H, m), 5.06 (1H, dd, J=7.5, 4.6Hz), 5.24 (1H, dd, J=9.8, 2.9Hz), 6.55 (1H, d, J=7.9Hz), 6.92 (1H, t, J=7.9Hz), 7.17-7.27 (5H, m), 7.34-7.36 (2H, m), 7.99-8.02 (1H, m), 8.50 (1H, d, J=7.8Hz), 8.79 (1H, brs), 8.87 (1H, brs).

Example 171

25 (1R)-2-{{2-(7-{{(1S)-2-Morpholin-4-yl-2-oxo-1-phenylethyl}oxy}-1H-indol-3-yl)ethyl}amino}-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example

119.

¹H-NMR (CDCl₃) δ: 2.68 (1H, dd, J=12.3, 9.3Hz), 2.88-3.05 (5H, m), 3.19-3.22 (1H, m), 3.33-3.36 (1H, m), 3.49-3.63 (6H, m), 4.70 (1H, dd, J=9.3, 3.5Hz), 6.02 (1H, s), 6.74 (1H, dd, J=7.7, 1.6Hz), 6.98 (1H, dd, J=7.8, 7.7Hz), 6.99 (1H, d, J=1.6Hz), 7.24 (1H, dd, J=7.9, 4.8Hz), 7.26 (1H, d, J=7.8Hz), 7.36-7.45 (3H, m), 7.55-7.57 (2H, m), 7.68 (1H, ddd, J=7.9, 1.9, 1.6Hz), 8.48 (1H, dd, J=4.8, 1.6Hz), 8.55 (1H, d, J=1.9Hz), 8.94 (1H, brs).

Example 172

(2S)-{[3-(2-([(2R)-2-Hydroxy-2-pyridin-3-ylethyl]amino)ethyl)-1H-indol-7-yl]oxy}(phenyl)acetic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CD₃OD) δ: 3.17-3.27 (3H, m), 3.37-3.40 (3H, m), 5.23 (1H, d, J=9.7Hz), 5.92 (1H, s), 6.58 (1H, d, J=7.8Hz), 6.88 (1H, t, J=7.8Hz), 7.18-7.20 (2H, m), 7.31-7.40 (3H, m), 7.63-7.64 (2H, m), 7.98 (1H, brs), 8.47-8.50 (1H, m), 8.78 (1H, brs), 8.86 (1H, brs).

Example 173

(1R)-2-[[2-(7-([(1S)-1-(Morpholin-4-ylcarbonyl)propyl]oxy)-1H-indol-3-yl)ethyl]amino]-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CD₃OD) δ: 1.14 (3H, t, J=7.4Hz), 2.00-2.08 (2H, m), 2.83 (2H, d, J=6.5Hz), 2.94-3.03 (4H, m), 3.31-3.35 (1H, m), 3.47-3.61 (5H, m), 3.68-3.74 (1H, m), 3.77-3.83 (1H, m), 4.81 (1H, t, J=6.5Hz), 5.07 (1H, t, J=6.1Hz), 6.55 (1H, d, J=7.7Hz), 6.90 (1H, dd, J=8.0, 7.7Hz), 7.06 (1H, s), 7.19 (1H, d, J=8.0Hz), 7.36 (1H, dd, J=7.8, 4.9Hz), 7.76

(1H, ddd, J=7.8, 2.1, 1.6Hz), 8.41 (1H, dd, J=4.9, 1.6Hz), 8.51 (1H, d, J=2.1Hz).

Example 174

(1R)-2-{{2-(7-(((1S)-1-(Morpholin-4-ylcarbonyl)butyl)oxy)-1H-indol-3-yl)-ethyl}amino}-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.02 (3H, t, J=7.4Hz), 1.52-1.58 (1H, m), 1.65-1.70 (1H, m), 1.87-1.94 (1H, m), 1.98-2.08 (1H, m), 2.68 (1H, dd, J=12.3, 9.4Hz), 2.92-3.06 (5H, m), 3.37-3.49 (2H, m), 3.58-3.62 (6H, m), 4.70 (1H, dd, J=9.4, 3.5Hz), 4.94 (1H, dd, J=8.6, 4.8Hz), 6.64 (1H, d, J=7.4Hz), 6.97-7.02 (2H, m), 7.23-7.27 (2H, m), 7.69 (1H, ddd, J=7.9, 2.1, 1.6Hz), 8.51 (1H, dd, J=4.8, 1.6Hz), 8.56 (1H, d, J=2.1Hz), 8.56 (1H, brs).

Example 175

6-(2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)ethyl)-2,2-dimethyl-[1,4]oxazino[2,3,4-hi]indol-3(2H)-one · 2 hydrochloride

The title compound is obtained in a similar manner to Example 131.

¹H-NMR (CD₃OD) δ: 1.65 (6H, s), 3.23-3.36 (3H, m), 3.47-3.53 (3H, m), 6.38 (1H, dd, J=9.9, 2.8Hz), 6.84 (1H, d, J=7.8Hz), 7.20 (1H, t, J=7.8Hz), 7.30 (1H, d, J=7.8Hz), 7.63 (1H, s), 8.14 (1H, dd, J=8.1, 5.8Hz), 8.75 (1H, d, J=8.1Hz), 8.86 (1H, d, J=5.8Hz), 9.01 (1H, s).

Example 176

(2S)-2-{{3-(2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)ethyl)-1H-indol-7-yl}oxy}butanoic acid · 2 trifluoroacetate

The title compound is obtained in a similar manner to Example

120.

¹H-NMR (CD₃OD) δ: 1.13 (3H, t, J=7.4Hz), 2.02-2.11 (2H, m),
3.18-3.24 (3H, m), 3.30-3.35 (1H, m), 3.38-3.42 (2H, m), 4.81 (1H, dd,
J=6.7, 5.2Hz), 5.11 (1H, dd, J=10, 3.1Hz), 6.56 (1H, d, J=7.7Hz), 6.95
5 (1H, t, J=7.9Hz), 7.17 (1H, s), 7.21 (1H, d, J=7.9Hz), 7.72 (1H, dd, J=7.7,
5.3Hz), 8.15 (1H, d, J=8.0Hz), 8.64 (1H, d, J=4.7Hz), 8.70 (1H, brs).

Example 177

(2S)-2-([3-(2-([(2R)-2-Hydroxy-2-pyridin-3-ylethyl]amino)ethyl)-1H-indol-7-yl]oxy)pentanoic acid · 2 trifluoroacetate

10 The title compound is obtained in a similar manner to Example
120.

¹H-NMR (CD₃OD) δ: 1.00 (3H, t, J=7.4Hz), 1.59-1.65 (2H, m),
1.99-2.05 (2H, m), 3.18-3.25 (3H, m), 3.32-3.42 (3H, m), 4.83-4.89 (1H,
m), 5.14 (1H, dd, J=10, 3.1Hz), 6.55 (1H, d, J=7.9Hz), 6.95 (1H, t,
15 J=7.9Hz), 7.17 (1H, s), 7.21 (1H, d, J=8.0Hz), 7.78 (1H, dd, J=7.7,
5.5Hz), 8.21 (1H, d, J=8.1Hz), 8.67 (1H, d, J=5.1Hz), 8.73 (1H, brs).

Example 178

2-([3-(2-([(2R)-2-Hydroxy-2-pyridin-3-ylethyl]amino)ethyl)-1H-indol-7-yl]oxy)-2-methylpropanoic acid · 2 trifluoroacetate

20 The title compound is obtained in a similar manner to Example
120.

¹H-NMR (CD₃OD) δ: 1.62 (6H, s), 3.17-3.25 (3H, m), 3.32-3.41
(3H, m), 5.09 (1H, dd, J=10, 3.0Hz), 6.65 (1H, d, J=7.7Hz), 6.92 (1H, t,
J=7.9Hz), 7.14 (1H, s), 7.26 (1H, d, J=7.9Hz), 7.65-7.68 (1H, m), 8.10
25 (1H, d, J=7.3Hz), 8.61 (1H, d, J=4.1Hz), 8.68 (1H, brs).

Example 179

(1R)-2-[(2-{7-[(1S)-1-Cyclohexyl-2-morpholin-4-yl-2-oxoethoxy]-1H-

indol-3-yl}ethyl)amino]-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ: 1.17-1.34 (5H, m), 1.71-1.74 (2H, m), 1.81-1.84 (2H, m), 1.97-1.99 (1H, m), 2.14-2.17 (1H, m), 2.66 (1H, dd, J=9.4, 12.3Hz), 2.92-3.05 (5H, m), 3.34 (2H, m), 3.49-3.67 (6H, m), 4.65-4.69 (2H, m), 6.68 (1H, d, J=7.6Hz), 6.99 (1H, dd, J=7.8, 7.9Hz), 7.01 (2H, d, J=1.9Hz), 7.24-7.27 (2H, m), 7.68-7.71 (1H, m), 8.48 (1H, brs), 8.51 (1H, dd, J=1.6, 4.8Hz), 8.57 (1H, d, J=2.1Hz).

Example 180

(2S)-Cyclohexyl{[3-(2-(((2R)-2-hydroxy-2-pyridin-3-ylethyl)amino)ethyl)-1H-indol-7-yl]oxy}acetic acid

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (CD₃OD) δ: 1.23-1.41 (5H, m), 1.69-1.71 (1H, m), 1.81 (3H, m), 1.95-1.98 (1H, m), 2.07-2.09 (1H, m), 2.93-3.14 (6H, m), 4.35 (1H, d, J=6.3Hz), 4.99 (1H, dd, J=3.2, 10.3Hz), 6.58 (1H, d, J=7.7Hz), 6.79 (1H, dd, J=7.8, 7.9Hz), 6.98 (1H, s), 7.01 (1H, d, J=7.7Hz), 7.45 (1H, dd, J=4.8, 7.9Hz), 7.84-7.87 (1H, m), 8.49 (1H, dd, J=1.5, 4.9Hz), 8.56 (1H, d, J=2.0Hz).

Example 181

Ethyl 1-([3-(2-(((2R)-2-hydroxy-2-pyridin-3-ylethyl)amino)ethyl)-1H-indol-7-yl]oxy)cyclobutanecarboxylate

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CD₃OD) δ: 1.13 (3H, t, J=7.1Hz), 1.99-2.07 (2H, m), 2.51-2.59 (2H, m), 2.75-2.82 (2H, m), 3.17-3.26 (3H, m), 3.31-3.48 (3H,

m), 4.17 (2H, q, J=7.1Hz), 5.13 (1H, dd, J=10, 3.1Hz), 6.15 (1H, d, J=7.7Hz), 6.86 (1H, t, J=7.9Hz), 7.17 (1H, s), 7.19 (1H, d, J=7.9Hz), 7.74 (1H, dd, J=7.9, 5.3Hz), 8.20 (1H, d, J=7.9Hz), 8.66 (1H, d, J=4.7Hz), 8.74 (1H, brs).

5 Example 182

1-[[3-(2-[[[(2R)-2-Hydroxy-2-pyridin-3-ylethyl]amino]ethyl)-1H-indol-7-yl]oxy]cyclobutanecarboxylic acid

The title compound is obtained in a similar manner to Example 120.

10 ¹H-NMR (CD₃OD) δ: 1.97-2.13 (2H, m), 2.50-2.58 (2H, m), 2.76-2.83 (2H, m), 3.17-3.25 (3H, m), 3.31-3.41 (3H, m), 5.14 (1H, dd, J=10, 3.1Hz), 6.21 (1H, d, J=7.7Hz), 6.87 (1H, t, J=7.9Hz), 7.17 (1H, s), 7.18 (1H, d, J=8.0Hz), 7.78 (1H, dd, J=7.9, 5.4Hz), 8.23 (1H, d, J=8.0Hz), 8.67 (1H, d, J=4.8Hz), 8.74 (1H, brs).

15 Reference Example 103

1-[7-(Benzyloxy)-1H-indol-3-yl]-2-methylpropan-2-amine

The title compound is obtained in a similar manner to Reference Example 4.

20 ¹H-NMR (CDCl₃) δ: 1.17 (6H, s), 2.80 (2H, s), 5.21 (2H, s), 6.72 (1H, d, J=7.6Hz), 7.02 (1H, dd, J=8.2, 7.6Hz), 7.02 (1H, d, J=2.7Hz), 7.25 (1H, d, J=8.2Hz), 7.34-7.43 (3H, m), 7.48-7.50 (2H, m), 8.37 (1H, brs).

Reference Example 104

tert-Butyl {2-[7-(benzyloxy)-1H-indol-3-yl]-1,1-dimethylethyl}carbamate

25 The title compound is obtained in a similar manner to Reference Example 88.

¹H-NMR (CDCl₃) δ: 1.32 (6H, s), 1.46 (9H, s), 3.09 (2H, s), 4.43

(1H, s), 5.20 (2H, s), 6.71 (1H, d, J=7.6Hz), 6.97 (1H, d, J=2.3Hz), 7.01 (1H, dd, J=8.1, 7.6Hz), 7.25 (1H, d, J=8.1Hz), 7.34 (3H, m), 7.47-7.49 (2H, m), 8.32 (1H, brs).

Reference Example 105

5 tert-Butyl (1,1-dimethyl-2-{7-[(1S)-1-methyl-2-morpholin-4-yl-2-oxoethoxy]-1H-indol-3-yl}ethyl)carbamate

The title compound is obtained in a similar manner to Reference Example 89.

10 ¹H-NMR (CDCl₃) δ: 1.31 (3H, s), 1.32 (3H, s), 1.47 (9H, s), 1.66 (3H, d, J=6.7Hz), 3.09 (2H, s), 3.40-3.66 (8H, m), 4.41 (1H, s), 5.13 (1H, q, J=6.7Hz), 6.62 (1H, d, J=7.7Hz), 6.98 (1H, dd, J=7.9, 7.7Hz), 6.99 (1H, d, J=2.5Hz), 7.27 (1H, d, J=7.9Hz), 8.50 (1H, brs).

Reference Example 106

15 2-Methyl-1-{7-[(1S)-1-methyl-2-morpholin-4-yl-2-oxoethoxy]-1H-indol-3-yl}propan-2-amine

To a solution of tert-butyl (1,1-dimethyl-2-{7-[(1S)-1-methyl-2-morpholin-4-yl-2-oxoethoxy]-1H-indol-3-yl}ethyl)carbamate (166 mg, 0.373 mmol) in methylene chloride (2 mL) is added a 4N hydrogen chloride solution in dioxane (0.5 mL), and the mixture is stirred at room temperature for 3 hours. The reaction solution is poured into a saturated aqueous sodium hydrogen carbonate solution, and the mixture is extracted with chloroform. The organic layer is dried over anhydrous magnesium sulfate, and the solvent is evaporated. The obtained crude product is purified by silica gel column chromatography (a saturated ammonia solution in chloroform → a saturated ammonia solution in chloroform - methanol = 50:1) to give the title compound (68.4 mg, yield: 53 %).

20

25

¹H-NMR (CDCl₃) δ: 1.17 (6H, s), 1.67 (3H, d, J=6.7Hz), 2.76 (2H, s), 3.40-3.70 (8H, m), 5.14 (1H, q, J=6.7Hz), 6.62 (1H, d, J=7.7Hz), 6.99 (1H, dd, J=8.1, 7.7Hz), 7.04 (1H, d, J=2.1Hz), 7.27 (1H, d, J=8.1Hz), 8.61 (1H, brs).

5 Example 183

(2S)-2-([3-(2-([(2R)-2-Hydroxy-2-pyridin-3-ylethyl]amino)-2-methylpropyl)-1H-indol-7-yl]oxy)propanoic acid · 2 trifluoroacetate

To a solution of 2-methyl-1-{7-[(1S)-1-methyl-2-morpholin-4-yl-2-oxoethoxy]-1H-indol-3-yl}propan-2-amine (68.4 mg, 0.198 mmol) in
 10 ethanol (4 mL) - water (0.4 mL) is added (R)-(pyridin-3-yl)oxirane (36.0 mg, 0.297 mmol), and the mixture is refluxed for 8 hours. To the reaction solution is added (R)-(pyridin-3-yl)oxirane (144.0 mg, 1.19 mmol) in 4 portions, and the mixture is heated with stirring at 100 to 110°C in a sealed tube for 18 hours. After the reaction is completed,
 15 the solvent is evaporated under reduced pressure, and the obtained residue is purified by silica gel column chromatography (chloroform/methanol = 20/1 → 10/1 → chloroform/a saturated ammonia methanol = 10/1) to give (1R)-2-[(1,1-dimethyl-2-{7-[(1S)-1-methyl-2-morpholin-4-yl-2-oxoethoxy]-1H-indol-3-yl}ethyl)amino]-1-pyridin-3-ylethanol (23.1
 20 mg).

To a solution of (1R)-2-[(1,1-dimethyl-2-{7-[(1S)-1-methyl-2-morpholin-4-yl-2-oxoethoxy]-1H-indol-3-yl}ethyl)amino]-1-pyridin-3-ylethanol (23.1 mg, 0.0495 mmol) in methanol (0.3 mL)-tetrahydrofuran (0.3 mL) is added a 2M aqueous lithium hydroxide solution (0.3 mL),
 25 and the mixture is stirred at room temperature overnight. After the reaction is completed, the mixture is neutralized with a 1N hydrochloric acid solution, and the solvent is evaporated under reduced pressure.

The obtained residue is purified by preparative high performance liquid chromatography (0.035 % trifluoroacetic acid/acetonitrile - 0.05 % trifluoroacetic acid/water) to give the title compound (12.6 mg, yield: 10 %).

¹H-NMR (CD₃OD) δ: 1.40 (s, 3H), 1.41 (s, 3H), 1.68 (s, 3H), 3.13-3.17 (m, 2H), 3.21-3.31 (m, 1H), 3.45 (dd, 1H, J=13, 2.7Hz), 4.98-5.01 (m, 1H), 5.21 (dd, 1H, J=10, 2.5Hz), 6.58 (d, 1H, J=7.8Hz), 6.97 (dd, 1H, J=8.0, 7.8Hz), 7.22 (s, 1H), 7.24 (d, 1H, J=8.0Hz), 8.03 (dd, 1H, J=6.9, 6.4Hz), 8.50 (d, 1H, J=8.1Hz), 8.81 (m, 1H), 8.88 (brs, 1H).

Reference Example 107

4-{{tert-Butyl(dimethyl)silyl}oxy}-1H-indole

To a suspension of 4-hydroxyindole (5.65 g, 42.4 mmol) in chloroform (200 mL) are added imidazole (4.33 g, 63.6 mmol) and tert-butylchlorodimethylsilane (7.03 g, 46.6 mmol), and the mixture is stirred at room temperature for 2.5 hours. The reaction mixture is washed with a mixture of a saturated brine and water (1:1, 100 mL), and the organic layer is dried over anhydrous potassium carbonate, filtered, and the filtrate is concentrated to about 50 mL under reduced pressure. The resultant is purified by silica gel column (300 g, ethyl acetate:hexane = 1:5 → 1:1) to give the title compound as a white solid (3.90 g, yield: 37 %), and the starting compound (2.62g, recovery rate: 46 %).

¹H-NMR (CDCl₃) δ: 0.24 (6H, s), 1.06 (9H, s), 6.52 (1H, dd, J=6.0, 2.0Hz), 6.58-6.59 (1H, m), 7.00-7.03 (2H, m), 7.09 (1H, dd, J=3.2, 2.4Hz), 8.10 (1H, brs).

Reference Example 108

(4-{{tert-Butyl(dimethyl)silyl}oxy}-1H-indol-1-yl)acetonitrile

Under nitrogen atmosphere, to a suspension of sodium hydride (60 % dispersion in mineral oil, 750 mg, 18.8 mmol) in dimethylformamide (30 mL) is added dropwise a solution of 4-{{tert-butyl-(dimethyl)silyl}oxy}-1H-indole (2.32 g, 9.38 mmol) in dimethylformamide (10 mL) with stirring under ice-cooling, and the mixture is stirred at the same temperature for 15 minutes. To the mixture is added dropwise acetonitrile bromide (1.96 mL, 28.1 mmol), and the mixture is stirred at room temperature for 15 hours. The reaction mixture is diluted with a mixture of a saturated brine and water (1:1, 400 mL), and the mixture is extracted with ethyl acetate (2 x 100 mL). The combined organic layer is washed with a mixture of a saturated brine and water (1:1, 100 mL), dried over anhydrous potassium carbonate, and filtered. The filtrate is concentrated under reduced pressure, and the obtained residue is purified by silica gel column (ethyl acetate:hexane = 10:1 → 5:1) to give the title compound as a pale brown solid (803 mg, yield: 39 %).

¹H-NMR (CDCl₃) δ: 0.23 (6H, s), 1.06 (9H, s), 4.98 (2H, s), 6.60 (1H, dd, J=8, 0.8Hz), 6.64 (1H, dd, J=3.2, 0.8Hz), 6.98 (1H, d, J=8.4Hz), 7.00 (1H, d, J=3.2Hz), 7.16 (1H, dd, J=8.4, 8.0Hz).

Reference Example 109

2-(4-{{tert-Butyl(dimethyl)silyl}oxy}-1H-indol-1-yl)ethylamine

To a solution of (4-{{tert-butyl(dimethyl)silyl}oxy}-1H-indol-1-yl)acetonitrile (259 mg, 0.904 mmol) in tetrahydrofuran (20 mL) are added a 2.0M hydrochloric acid in ethanol (1 mL) and active Raney-nickel (manufactured by Kawaken Fine Chemicals Co., Ltd., about 1 mL), and the mixture is vigorously stirred under hydrogen atmosphere under pressure (0.5 Pa) at room temperature for 1 hour. To the

mixture is further added active Raney-nickel (about 1.5 mL), and the mixture is vigorously stirred under the same conditions for 4 hours. The catalyst is removed by filtration through celite, and the filtrate is concentrated under reduced pressure. The obtained residue is purified
 5 by silica gel column (ammonia - chloroform:methanol = 100:1 → 50:1 → 30:1) to give the title compound as pale brown oil (207 mg, yield: 79 %).

¹H-NMR (CDCl₃) δ: 0.24 (6H, s), 1.06 (9H, s), 3.12 (2H, t, J=6.0Hz), 4.17 (2H, t, J=6.0Hz), 6.51 (1H, d, J=7.2Hz), 6.54 (1H, d, J=3.2Hz), 6.96-7.07 (3H, m).

10 Reference Example 110

(2R)-N-[2-(4-{{tert-Butyl(dimethyl)silyl}oxy}-1H-indol-1-yl)ethyl]-2-hydroxy-2-pyridin-3-ylacetamide

The title compound is obtained in a similar manner to Reference Example 61.

15 ¹H-NMR (CDCl₃) δ: 0.24 (6H, s), 1.06 (9H, s), 3.59-3.73 (3H, m), 4.22-4.25 (2H, m), 6.40 (1H, t, J=6Hz), 6.50-6.53 (2H, m), 6.80 (1H, d, J=3.6Hz), 6.91 (1H, d, J=8.0Hz), 7.04 (1H, t, J=8.0Hz), 7.25 (1H, ddd, J=0.8, 4.8, 8.0Hz), 7.60 (1H, dt, J=7.6, 2.0Hz), 8.51-8.52 (2H, m).

Reference Example 111

20 tert-Butyl [2-(4-{{tert-butyl(dimethyl)silyl}oxy}-1H-indol-1-yl)ethyl][(2R)-2-hydroxy-2-pyridin-3-ylethyl]carbamate

To a solution of (2R)-N-[2-(4-{{tert-butyl(dimethyl)silyl}oxy}-1H-indol-1-yl)ethyl]-2-hydroxy-2-pyridin-3-ylacetamide (498 mg, 1.17 mmol) in tetrahydrofuran (5 mL) is added a 2.0 M solution of borane dimethylsulfide complex in tetrahydrofuran (1.76 mL, 3.51 mmol), and
 25 the mixture is refluxed for 3.5 hours. To the reaction solution is added piperidine (1.0 mL), and the mixture is further refluxed for 2 hours.

The reaction mixture is separated into a mixture of a saturated brine and water (1:1, 30 mL) and chloroform (30 mL). The aqueous layer is extracted with ethyl acetate (2 x 30 mL), and the combined organic layer is washed with a mixture of a saturated brine and water (1:1, 30 mL), dried over anhydrous potassium carbonate, and filtered. The filtrate is concentrated under reduced pressure, and the obtained residue is subjected to azeotropic distillation with toluene to remove piperidine. To a solution of the yellow residue thus obtained in tetrahydrofuran (5.0 mL) is added di-tert-butyl bicarbonate (296 μ L, 1.29 mmol), and the mixture is stirred at room temperature for 3.5 hours. The reaction mixture is concentrated under reduced pressure, and the obtained orange residue is purified by silica gel column (ethyl acetate:hexane = 1:1) to give the title compound as oil (316 mg, yield: 53 %).

$^1\text{H-NMR}$ (CDCl_3) δ : 0.18 (6H, s), 1.02 (9H, s), 1.40 (9H, s), 2.55-2.64 (1H, m), 3.06-3.16 (1H, m), 3.32-3.52 (2H, m), 4.09-4.34 (2H, m), 4.72 (1H, m), 6.85-7.00 (2H, m), 7.05 (1H, dd, $J=8.0, 7.6\text{Hz}$), 7.21-7.24 (1H, m), 7.42-7.55 (1H, m), 8.35-8.42 (1H, m), 8.46-8.47 (1H, m).

Reference Example 112

tert-Butyl [2-(4-hydroxy-1H-indol-1-yl)ethyl][(2R)-2-hydroxy-2-pyridin-3-ylethyl]carbamate

To a solution of tert-butyl [2-(4-[[tert-butyl(dimethyl)silyl]oxy]-1H-indol-1-yl)ethyl][(2R)-2-hydroxy-2-pyridin-3-ylethyl]carbamate (316 mg, 0.618 mmol) in tetrahydrofuran (5 mL) is added a 1.0M solution of tetrabutylammonium fluoride in tetrahydrofuran (927 μ L, 1.5 equivalent), and the mixture is stirred at room temperature for 3 hours. Further, to the mixture is added a 1.0 M solution of tetrabutylammonium fluoride in tetrahydrofuran (927 μ L, 1.5 equivalent), and the

mixture is stirred for 15 hours. The reaction mixture is concentrated under reduced pressure, and the obtained brown residue is purified by silica gel column (ammonia - chloroform:methanol = 100:1 → 50:1 → 20:1) to give the title compound as oil (239 mg, yield: 97 %).

LC/MS M+1: 398, retention time: 2.89 min.

Conditions for analysis:

Body: API 150EX (Applied Biosystems)

Method for ionization: ESI

Column: CombiScreen Hydrosphere C18 S-5 μ m (4.6 x 50 mm) (YMC)

Mobile phase: Solution A: 0.05 % aqueous trifluoroacetic acid

Solution B: 0.035 % trifluoroacetic acid in acetonitrile

Flow rate: 3.5 mL/min.

Conditions for HPLC: 0.0 min → 0.5 min: 90 % Solution A constant

0.5 min → 4.2 min: 90 % Solution A → 1 %

4.2 min → 4.4 min: 1 % Solution A constant

Example 184

(1R)-2-[(2-{4-[(1S)-1-Methyl-2-morpholin-4-yl-2-oxoethoxy]-1H-indol-1-yl}ethyl)amino]-1-pyridin-3-ylethanol

The title compound is obtained in a similar manner to Example 119.

¹H-NMR (CDCl₃) δ : 1.71 (3H, d, J=6.8Hz), 2.64 (1H, dd, J=12, 9.2Hz), 2.88 (1H, dd, J=12, 3.6Hz), 3.04-3.17 (2H, m), 3.49-3.68 (8H, m), 4.26 (2H, t, J=6.0Hz), 4.63 (1H, dd, J=9.2, 3.6Hz), 5.14 (1H, q, J=6.8Hz), 6.55 (1H, d, J=8.0Hz), 6.59 (1H, d, J=3.2Hz), 7.02 (1H, d, J=8.0Hz), 7.04 (1H, d, J=3.2Hz), 7.11 (1H, t, J=8.0Hz), 7.23-7.26 (1H, m), 7.64 (1H, ddd, J=7.6, 2.0, 1.6Hz), 8.51 (1H, dd, J=4.8, 1.2Hz), 8.53 (1H, d, J=2.4Hz).

Example 185

(2S)-2-([1-(2-(((2R)-2-Hydroxy-2-pyridin-3-ylethyl)amino)ethyl)-1H-indol-4-yl]oxy)propanoic acid

The title compound is obtained in a similar manner to Example 120.

¹H-NMR (DMSO-d₆) δ: 1.57 (3H, d, J=6.8Hz), 3.04-3.35 (4H, m), 4.57 (2H, t, J=6.8Hz), 4.89 (1H, q, J=6.8Hz), 5.04 (1H, d, J=9.6Hz), 6.41 (1H, d, J=8.0Hz), 6.41 (1H, brs), 6.51 (1H, dd, J=3.2, 0.8Hz), 7.07 (1H, t, J=8.0Hz), 7.22 (1H, d, J=8.0Hz), 7.31 (1H, d, J=2.8Hz), 7.46 (1H, dd, J=8.0, 4.8Hz), 7.80 (1H, ddd, J=8.0, 2.0, 1.6Hz), 8.55 (1H, dd, J=4.8, 1.6Hz), 8.60 (1H, d, J=2.0Hz), 9.14 (1H, brs), 9.44 (1H, brs).

Reference Example 113

N-(2-Methoxy-6-methylphenyl)acetamide

To a solution of 3-methyl-2-nitroanisole (10.00 g, 59.8 mmol) in ethanol (60 mL)-tetrahydrofuran (60 mL) is added a 10 % palladium on carbon (50 % wet, 2.00 g), and the mixture is subjected to hydrogenation, and stirred at room temperature for 1.5 hour. After the reaction is completed, the reaction solution is filtered through celite, and the solvent is evaporated under reduced pressure. To a solution of the residue in ethyl acetate (110 mL) is added acetic anhydride (9.0 mL, 95.2 mmol), and the mixture is refluxed for 5 hours. The reaction solution is cooled with ice, and the precipitated solid is collected by filtration, and washed with hexane to give the title compound (9.28 g, yield: 87 %).

¹H-NMR (CDCl₃) δ: 2.21 (3H, s), 2.24 (3H, s), 3.81 (3H, s), 6.75 (1H, d, J=8.0Hz), 6.85 (1H, d, J=8.0Hz), 7.13 (1H, t, J=8.0Hz).

Reference Example 114

7-Methoxy-1H-indazole

To a solution of N-(2-methoxy-6-methylphenyl)acetamide (9.28 g, 51.8 mmol) in ethyl acetate (100 ml) are added acetic anhydride (14.7 mL, 156 mmol), tetrabutylammonium bromide (0.84 g, 2.61 mmol), potassium acetate (10.16 g, 104 mmol), and isoamyl nitrite (9.1 mL, 67.7 mmol), and the mixture is refluxed for 9 hours. After the reaction is completed, the solvent is evaporated under reduced pressure, and to the obtained residue is added 6N aqueous sodium hydroxide solution (100 mL). The mixture is stirred at room temperature for 1 hour. After the reaction is completed, the pH value of the mixture is adjusted to pH 7-8 with 3N hydrochloric acid, and the mixture is extracted with chloroform. The organic layer is dried over anhydrous magnesium sulfate, and the solvent is evaporated under reduced pressure. The obtained residue is purified by silica gel column chromatography (n-hexane/ethyl acetate = 3/1 → 1/1) to give the title compound (3.95 g, yield: 51 %).

¹H-NMR (DMSO-d₆) δ: 3.94 (3H, s), 6.82 (1H, d, J=7.5Hz), 7.02 (1H, dd, J=8.0, 7.5Hz), 7.30 (1H, d, J=8.0Hz), 8.01 (1H, d, J=1.5Hz), 13.27 (brs, 1H).

Reference Example 115

3-Iodo-7-methoxy-1H-indazole

To a solution of 7-methoxy-1H-indazole (2.60 g, 17.5 mmol) in N,N-dimethylformamide (50 mL) are added iodine (6.68 g, 26.3 mmol) and potassium hydroxide (2.79 g, 49.7 mmol), and the mixture is stirred at room temperature for 0.5 hour. The reaction solution is poured into a 10 % aqueous sodium hydrogen sulfite solution (200 mL), and the mixture is extracted with diethyl ether. The organic layer is

washed with water and a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated under reduced pressure, and the obtained residue is purified by silica gel column chromatography (n-hexane/ethyl acetate = 5/1) to give the title compound (3.97 g, yield: 83 %).

$^1\text{H-NMR}$ (CDCl_3) δ : 3.99 (3H, s), 6.80 (1H, d, $J=7.4\text{Hz}$), 7.09 (1H, d, $J=8.1\text{Hz}$), 7.15 (1H, dd, $J=8.1, 7.4\text{Hz}$), 10.24 (1H, brs).

Reference Example 116

tert-Butyl 3-iodo-7-methoxy-1H-indazole-1-carboxylate

To a solution of 3-iodo-7-methoxy-1H-indazole (1.33 g, 4.86 mmol) in acetonitrile (20 mL) are added di-t-butyl bicarbonate (1.34 mL, 5.83 mmol), triethylamine (0.81 mL, 5.81 mmol) and N,N-dimethylaminopyridine (60.0 mg, 0.491 mmol), and the mixture is stirred at room temperature for 3 hours. To the reaction solution is added water, and the mixture is extracted with ethyl acetate. The organic layer is washed with a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated under reduced pressure, and the obtained residue is purified by silica gel column chromatography (n-hexane/ethyl acetate = 5/1) to give the title compound (1.79 g, yield: 98 %).

$^1\text{H-NMR}$ (CDCl_3) δ : 1.68 (9H, s), 3.99 (3H, s), 7.01 (1H, d, $J=7.8\text{Hz}$), 7.10 (1H, dd, $J=8.0, 0.7\text{Hz}$), 7.30 (1H, dd, $J=8.0, 7.8\text{Hz}$).

Reference Example 117

tert-Butyl 7-methoxy-3-[(1E)-3-methoxy-3-oxopropyl-1-en-1-yl]-1H-indazole-1-carboxylate

To a solution of tert-butyl 3-iodo-7-methoxy-1H-indazole-1-carboxylate (484 mg, 1.29 mmol) in N,N-dimethylformamide:water (11:1,

76 mL) are added triethylamine (1.76 mL, 12.6 mmol), methyl acrylate (1.17 mL, 13.0 mmol), dichloro[1,1'-bis(diphenylphosphono)ferrocene]-palladium (190 mg, 0.259 mmol) and tetra-n-butylammonium iodide (954 mg, 2.58 mmol), and the mixture is refluxed at 50°C for 4 hours.

5 After the reaction is completed, the excess methyl acrylate is removed by distillation under reduced pressure, and to the residue is added water (100 mL). The precipitates are collected by filtration, dissolved in ethyl acetate, and filtered through celite. The solvent is evaporated under reduced pressure, and the obtained residue is purified by silica
10 gel column chromatography (n-hexane/ethyl acetate = 3/1) to give the title compound (145 mg, yield: 38 %).

¹H-NMR (CDCl₃) δ: 1.69 (9H, s), 3.84 (3H, s), 3.99 (3H, s), 6.99 (1H, d, J=16.3Hz), 6.98 (1H, d, J=7.7Hz), 7.32 (1H, dd, J=8.0, 7.7Hz), 7.49 (1H, dd, J=8.0, 0.6Hz), 7.96 (1H, d, J=16.3Hz).

15 Reference Example 118

Methyl (2E)-3-(7-methoxy-1H-indazol-3-yl)acrylate

To a solution of tert-butyl 7-methoxy-3-[(1E)-3-methoxy-3-oxopropyl-1-en-1-yl]-1H-indazole-1-carboxylate (147 mg, 0.442 mmol) in tetrahydrofuran (2 mL) is added a solution of sodium methoxide in
20 methanol (0.2 mL), and the mixture is stirred at room temperature overnight. To the reaction solution is added water, and the mixture is extracted with diethyl ether. The organic layer is washed with a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated under reduced pressure, and the obtained residue
25 is purified by silica gel column chromatography (n-hexane/ethyl acetate = 3/1) to give the title compound (98.5 mg, yield: 96 %).

¹H-NMR (CDCl₃) δ: 3.85 (3H, s), 4.01 (3H, s), 6.79 (1H, d,

J=16.3Hz), 6.80 (1H, d, J=7.6Hz), 7.20 (1H, dd, J=8.1, 7.6Hz), 7.50 (1H, d, J=8.1Hz), 8.02 (1H, d, J=16.3Hz), 10.46 (1H, brs).

Reference Example 119

Methyl 3-(7-methoxy-1H-indazol-3-yl)propanoate

5 To a solution of methyl (2E)-3-(7-methoxy-1H-indazol-3-yl)-acrylate (143 mg, 0.614 mmol) in methanol (10 mL) is added nickel chloride · 6 hydrate (73.0 mg, 0.307 mmol), and further thereto is added sodium borohydride (93 mg, 2.46 mmol) under ice-cooling over a period of 0.5 hour. The mixture is stirred at room temperature for 4.5 hours.
10 After the reaction is completed, the solvent is evaporated under reduced pressure. Water is added to the residue, and the mixture is extracted with ethyl acetate. The organic layer is washed with a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated under reduced pressure, and the obtained residue is
15 purified by silica gel column chromatography (n-hexane/ethyl acetate = 3/1) to give the title compound (122 mg, yield: 85 %).

¹H-NMR (CDCl₃) δ: 2.89 (2H, dd, J=8.0, 7.4Hz), 3.30 (2H, dd, J=8.0, 7.4Hz), 3.70 (3H, s), 3.97 (3H, s), 6.73 (1H, d, J=7.5Hz), 7.06 (1H, dd, J=8.1, 7.5Hz), 7.28 (1H, dd, J=8.1, 0.3Hz), 10.06 (1H, brs)

20 Reference Example 120

3-(7-Methoxy-1H-indazol-3-yl)propanoic acid

 To a solution of methyl 3-(7-methoxy-1H-indazol-3-yl)-propanoate (140 mg, 0.596 mmol) in tetrahydrofuran (2 mL) is added a 6N aqueous sodium hydroxide solution (2 ml), and the mixture is
25 stirred at room temperature overnight. The reaction solution is washed with diethyl ether, and the aqueous layer is acidified with 6N hydrochloric acid, and the mixture is extracted with diethyl ether. The

organic layer is dried over anhydrous magnesium sulfate, and the solvent is evaporated under reduced pressure to give the title compound (123 mg, yield: 94 %).

¹H-NMR (CDCl₃) δ: 2.89 (dd, 2H, J=8.0, 7.4Hz), 3.30 (dd, 2H, J=8.0, 7.4Hz), 3.70 (s, 3H), 3.97 (s, 3H), 6.73 (d, 1H, J=7.5Hz), 7.06 (dd, 1H, J=8.1, 7.5Hz), 7.28 (dd, 1H, J=8.1, 0.3Hz), 10.06 (brs, 1H).

Reference Example 121

tert-Butyl [2-(7-methoxy-1H-indazol-3-yl)ethyl]carbamate

Under nitrogen atmosphere, to a solution of 3-(7-methoxy-1H-indazol-3-yl)propanoic acid (86.4 mg, 0.392 mmol) in toluene (5 mL)-tert-butanol (2.5 mL) are added triethylamine (80.0 μL, 0.574 mmol) and diphenylphosphoryl azide (110 μL, 0.510 mmol), and the mixture is stirred at room temperature for 2 hours. The mixture is stirred at 60°C for 1 hour, and further stirred at 100°C for 5 hours. The mixture is allowed to cool to room temperature, and thereto is added a saturated aqueous sodium hydrogen carbonate solution, and the mixture is extracted with ethyl acetate. The organic layer is dried over anhydrous magnesium sulfate, and the solvent is evaporated under reduced pressure. The obtained residue is purified by silica gel column chromatography (n-hexane/ethyl acetate = 3/1) to give the title compound (94.3 mg, yield: 82 %).

¹H-NMR (CDCl₃) δ: 1.43 (s, 9H), 3.17 (dd, 1H, J=12, 6.6Hz), 3.62 (dd, 1H, J=12, 5.9Hz), 3.98 (s, 3H), 5.08 (brs, 1H), 6.74 (d, 1H, J=7.5Hz), 7.04 (dd, 1H, J=8.1, 7.5Hz), 7.27 (d, 1H, J=8.1Hz), 10.19 (brs, 1H).

Reference Example 122

tert-Butyl [2-(7-hydroxy-1H-indazol-3-yl)ethyl]carbamate

To a solution of tert-butyl [2-(7-methoxy-1H-indazol-3-yl)ethyl]-carbamate (91.8 mg, 0.315 mmol) in dichloromethane (1 mL) is added a solution of boron tribromide (1.0 M dichloromethane solution, 0.95 mL, 0.95 mmol) under nitrogen atmosphere, and the mixture is refluxed for 2.5 hours. The reaction solution is cooled with ice, and thereto is added dropwise methanol (3 ml), and the mixture is stirred at room temperature for 2 hours. The solvent is concentrated under reduced pressure to give a crude phenol product.

To a solution of the crude phenol product in acetonitrile (3.5 mL) are added triethylamine (100 μ L, 0.717 mmol) and di-t-butyl bicarbonate (72.4 μ L, 0.315 mmol) under nitrogen atmosphere, and the mixture is stirred at room temperature for 0.5 hour. Water is added to the reaction solution, and the mixture is extracted with ethyl acetate. The organic layer is washed with a saturated brine, and dried over anhydrous magnesium sulfate. The solvent is evaporated under reduced pressure, and the obtained residue is purified by silica gel column chromatography (n-hexane/ethyl acetate = 2/1) to give the title compound (46.0 mg, yield: 54 % (2 steps)).

$^1\text{H-NMR}$ (DMSO-d_6) δ : 1.37 (s, 9H), 2.96 (dd, 2H, $J=8.1, 7.1\text{Hz}$), 3.24-3.30 (m, 2H), 6.63 (d, 1H, $J=7.2\text{Hz}$), 6.85 (dd, 1H, $J=8.0, 7.2\text{Hz}$), 6.88 (t, 1H, 5.7Hz), 7.10 (d, 1H, $J=8.0\text{Hz}$), 9.98 (brs, 1H), 12.65 (brs, 1H).

Reference Example 123

tert-Butyl (2-{7-[(1S)-1-methyl-2-morpholin-4-yl-2-oxoethoxy]-1H-indazol-3-yl}ethyl)carbamate

The title compound is obtained in a similar manner to Reference Example 84.

¹H-NMR (CDCl₃) δ: 1.41 (s, 9H), 1.68 (d, 3H, J=6.7Hz), 3.17 (t, 2H, J=6.4Hz), 3.43-3.47 (m, 1H), 3.56-3.64 (m, 9H), 5.28 (q, 1H, J=6.7Hz), 5.22 (m, 1H), 6.72 (d, 1H, J=7.6Hz), 7.01 (dd, 1H, J=8.1, 7.6Hz), 7.33 (d, 1H, J=8.1Hz).

5 Reference Example 124

(2-{7-[(1S)-1-Methyl-2-morpholin-4-yl-2-oxoethoxy]-1H-indazol-3-yl}-ethyl)amine

The title compound is obtained in a similar manner to Reference Example 106.

10 ¹H-NMR (CDCl₃) δ: 1.69 (d, 3H, J=6.7Hz), 3.10-3.18 (m, 4H), 3.42-3.46 (m, 1H), 3.56-3.64 (m, 7H), 5.15 (q, 1H, J=6.7Hz), 6.72 (d, 1H, J=7.6Hz), 7.02 (dd, 1H, J=8.1, 7.6Hz), 7.33 (d, 1H, J=8.1Hz).

Example 186

15 (2S)-2-[[3-(2-[[[(2R)-2-Hydroxy-2-pyridin-3-ylethyl]amino]ethyl)-1H-indazol-7-yl]oxy}propanoic acid

The title compound is obtained in a similar manner to Example 183.

20 ¹H-NMR (CD₃OD) δ: 1.71 (d, 3H, J=6.8Hz), 3.33-3.36 (m, 1H), 3.39-3.43 (m, 2H), 3.49 (dd, 1H, J=12.8, 3.1Hz), 3.53-3.63 (m, 2H), 5.02 (q, 1H, J=6.8Hz), 5.25 (dd, 1H, J=9.8, 3.1Hz), 6.75 (d, 1H, J=7.6Hz), 7.05 (dd, 1H, J=8.1, 7.6Hz), 7.33 (d, 1H, J=8.1Hz), 7.90 (dd, 1H, J=7.3, 6.1Hz), 8.43 (d, 1H, J=8.0Hz), 8.74 (d, 1H, J=5.3Hz), 8.85 (s, 1H).

Reference Example 125

25 2-[[tert-Butyl(dimethyl)silyl]oxy]-6-nitroaniline

The title compound is obtained in a similar manner to Reference Example 107.

¹H-NMR (CD₃OD) δ: 0.28 (s, 6H), 1.04 (s, 9H), 6.29 (brs, 2H),

6.53 (dd, 1H, J=8.8, 7.5Hz), 6.89 (dd, 1H, J=7.5, 1.3Hz), 7.75 (dd, 1H, J=8.8, 1.4Hz).

Reference Example 126

3-[[tert-Butyl(dimethyl)silyl]oxy]benzen-1,2-diamine

5 To a solution of 2-[[tert-butyl(dimethyl)silyl]oxy]-6-nitroaniline (8.5 g, 31.7 mmol) in tetrahydrofuran (100 mL) is added a 10 % palladium on carbon (50 % wet, 8.2 g), and the mixture is vigorously stirred under hydrogen atmosphere at room temperature for one hour. The catalyst is removed by filtration, and the filtrate is concentrated under reduced pressure. The obtained residue is purified by silica gel column chromatography (ethyl acetate/n-hexane = 1/5, 1/3 and 1/1) to give the title compound as brown oil (6.95 g, yield: 92 %).

¹H-NMR (CD₃OD) δ: 6.56 (t, 1H, J=8.0Hz), 6.37 (dd, 1H, J=8.0, 1.3Hz), 6.34 (dd, 1H, J=8.0, 1.3Hz), 3.15 (brs, 4H), 1.02 (s, 9H), 0.24 (s, 6H).

Reference Example 127

7-[[tert-Butyl(dimethyl)silyl]oxy]1H-benzimidazole

20 To 3-[[tert-butyl(dimethyl)silyl]oxy]benzene-1,2-diamine (4.27 g, 17.9 mmol) is added diethoxymethyl acetate (8.7 g, 53.6 mmol), and the mixture is stirred at room temperature for 5 hours. The precipitated white solid is collected by filtration, and the filtrate is further stirred at room temperature for 12 hours. The precipitated white solid is collected by filtration, and washed with a small amount of hexane. The filtrate is concentrated under reduced pressure, and the obtained residue is purified by silica gel column chromatography (ammoniacal chloroform) to give the title compound as white solid (1.89 g, yield: 43 %).

¹H-NMR (CD₃OD) δ: 9.16 (brs, 1H), 7.98 (s, 1H), 7.44 (brs, 1H), 7.13 (t, 1H, J=8.1Hz), 6.73 (d, 1H, J=8.6Hz), 1.05 (s, 9H), 0.29 (s, 6H).

Reference Example 128

(4-{{tert-Butyl(dimethyl)silyl}oxy}-1H-benzimidazol-1-yl)acetonitrile

5

The title compound is obtained in a similar manner to

Reference Example 108.

¹H-NMR (CD₃OD) δ: 7.85 (s, 1H), 7.25 (t, 1H, J=8.0Hz), 7.06 (dd, 1H, J=8.0, 0.8Hz), 6.80 (dd, 1H, J=8.0, 0.8Hz), 5.05 (s, 2H), 1.06 (s, 9H), 0.29 (s, 6H).

10

Reference Example 129

[2-(4-{{tert-Butyl(dimethyl)silyl}oxy}-1H-benzimidazol-1-yl)ethyl]amine

The title compound is obtained in a similar manner to

Reference Example 109 except that a 10 % palladium on carbon (50 % wet) is used instead of active Raney-nickel.

15

¹H-NMR (CD₃OD) δ: 7.87 (s, 1H), 7.14 (t, 1H, J=8.0Hz), 7.01 (dd, 1H, J=8.0, 0.7Hz), 6.72 (dd, 1H, J=8.0, 0.7Hz), 4.21 (t, 2H, J=5.9Hz), 3.15 (t, 2H, J=5.9Hz), 1.06 (s, 9H), 0.30 (s, 6H).

Example 187

(1R)-2-[(2-{4-[(1S)-1-Methyl-2-morpholin-4-yl-2-oxoethoxy]-1H-benzimidazol-1-yl}ethyl)amino]-1-pyridin-3-ylethanol

20

The title compound is obtained in a similar manner to

Reference Examples 61, 111 and 112 and Example 119.

¹H-NMR (CD₃OD) δ: 1.67 (d, 1H, J=6.4Hz), 3.40-3.45 (m, 1H), 3.58-3.98 (m, 11H), 5.04-5.06 (m, 2H), 5.46 (d, 1H, J=8.3Hz), 5.63 (q, 1H, J=6.4Hz), 7.11 (d, 1H, J=7.9Hz), 7.62 (t, 1H, J=8.0Hz), 7.71 (d, 1H, J=8.3Hz), 8.16 (dd, 1H, J=7.4, 6.1Hz), 8.80 (d, 1H, J=7.9Hz), 8.87 (d, 1H, J=5.4Hz), 9.04 (s, 1H), 9.66 (s, 1H).

25

Example 188

(2S)-2-([1-(2-([(2R)-2-Hydroxy-2-pyridin-3-ylethyl]amino)ethyl)-1H-benzimidazol-4-yl]oxy)propanoic acid

The title compound is obtained in a similar manner to Example

120.

¹H-NMR (CD₃OD) δ: 1.74 (d, 3H, J=6.8Hz), 3.30-3.37 (m, 3H), 3.52 (dd, 1H, J=12.6, 3.2Hz), 3.72-3.83 (m, 2H), 5.18 (q, 1H, J=6.8Hz), 5.29 (dd, 1H, J=9.7, 3.0Hz), 7.07 (dd, 1H, J=6.4, 2.3Hz), 7.53-7.58 (m, 2H), 7.97 (dd, 1H, J=7.9, 5.7Hz), 8.52 (d, 1H, J=8.1Hz), 8.77 (d, 1H, J=5.3Hz), 8.87 (s, 1H), 9.39 (s, 1H).

Experiment 1

Human β₃, β₂ and β₁-receptor stimulation test

Human β₃-receptor-stimulating activity was studied using SK-N-MC cell line, i.e., cells permanently expressing human β₃- and human β₁-receptors (SK-N-MC cells, purchased from Dainippon Pharmaceutical Co., Ltd.), and human β₂- and β₁-receptor-stimulating activities were studied using THP-1 cell line, i.e., cells expressing both human β₂- and β₁-receptors (THP-1 cells, purchased Dainippon Pharmaceutical Co., Ltd.).

The β₃-receptor stimulating activity of a test compound (10⁻⁹ to 10⁻⁵ M) was studied using as an index cyclic AMP (hereinafter, referred to as cAMP) producing activity of SK-N-MC cells at almost confluent in the presence of a β₁-receptor inhibitor ((±)-2-hydroxy-5-(2-((2-hydroxy-3-(4-(1-methyl-4-(trifluoromethyl)-1H-imidazol-2-yl)phenoxy)propyl)-amino)ethoxy)benzamide methanesulfonate, CGP-20712A (Sigma-Aldrich), 10⁻⁶M), wherein said SK-N-MC cells were seeded into a 96-well plate at a cell density of 2 x 10⁴ cells/well and had been cultured for 3

days. The β 2-receptor-stimulating activity of a test compound was studied by stimulating floating THP-1 cells (2.5×10^5 cells) with a test compound in the presence of a β 1-receptor inhibitor (the above-mentioned CGP-20712A, 10^{-6} M), and the β 1-receptor stimulating activity was studied by stimulating floating THP-1 cells (2.5×10^4 cells) with a test compound in the presence of a β 2-receptor inhibitor ((R*,S*)-(\pm)-1-((2,3-dihydro-7-methyl-1H-inden-4-yl)oxy)-3-((1-methylethyl)-amino)-2-butanol hydrochloride, ICI-118551 (Sigma-Aldrich), 10^{-8} M), and both activities were evaluated with using as an index cAMP producing activity thereof. The cAMP production in the cells was measured by ELISA method.

The action potency of each compound was evaluated by calculating EC_{50} value ($-\log_{10} EC_{50} = pD_2$) and intrinsic activity (%) from the dose-response curve thus obtained, wherein the responses to 10^{-6} M isoproterenol, 10^{-5} M salmeterol and 10^{-5} M isoproterenol were counted as 100 % for β 3-receptor-stimulating activity, β 2-receptor-stimulating activity and β 1-receptor-stimulating activity, respectively, and comparing them. It was confirmed that the present compound exhibits a selective stimulating activity on human β 3-receptor.

For example, the EC_{50} value of β 3-stimulating activity of the compound of Example 2 was 48 nM while the β 1- and β 2-stimulating activities thereof were not observed up to a maximum concentration of 10 μ M.

Experiment 2

Effect of a medicament on autonomic beat of the excised right atrium specimen based on β 1-adrenoceptor-stimulating activity (study on positive chronotropic action)

The heart of Hartley guinea pig was excised, and a right atrium specimen was prepared therefrom and the experiment was carried out thereon according to Magnus method. The specimen was hung at 34°C in Krebs-Henseleit solution to which a mixed gas of 95 % oxygen and 5 % carbon dioxide was aerated, and loaded so that the maximum resting tension became about 0.5 g. The autonomic beat of the right atrium was detected through a pressure transducer, and the tension and the beat were continuously recorded onto a linear recorder. A medicament was cumulatively added every time the beat became plateau. With counting the maximum increase in the beat by isoproterenol as 100 %, the ratio of the maximum increase in the beat by a medicament to that of isoproterenol was calculated, and the potency of a medicament was evaluated in terms of EC₅₀ value, which was a concentration of the medicament to be required for increasing the beat by 50 % of the maximum increase of the beat by said medicament.

In order to show the divergence between β 3-stimulating activity/ β 1-stimulating activity in the experimental results, the EC₅₀ value of positive chronotropic action of the test compound of Examples obtained in Experiment 2 was divided by the EC₅₀ value of human β 3-stimulating activity of said compound obtained in Experiment 1, which is the main activity in Experiment 1. The results are expressed by relative values, which are calculated by counting the value of the reference compound (AJ 9677) as 1.

As shown below, the compound of the feature [1] as defined herein before, especially the compound of the feature [2], exhibited a high divergence between β 3-stimulating activity and β 1-stimulating activity even in the evaluation system using tissues, as compared to the

reference compound (AJ 9677) which was disclosed in JP-A-11-255743.

EC₅₀ values of positive chronotropic effect of the compounds of Examples (relative values when the value of the reference compound (AJ 9677) is counted as 1)

5	The compound of Example 14:	68
	The compound of Example 73:	77
	The compound of Example 76:	300
	The compound of Example 114:	230
10	The compound of Example 121:	78
	The compound of Example 150:	470
	The compound of Example 154:	90
	Reference compound (AJ 9677):	1

15 INDUSTRIAL APPLICABILITY

The indole, indazole, and benzazole derivatives of the formula (I) and a pharmaceutically acceptable salt thereof exhibit excellent β 3-adrenoceptor-stimulating activity, and are useful as a medicament for treatment of obesity, hyperglycemia, diseases caused by increased intestinal motility, frequent urination, urinary incontinence, depression, 20 bilestone, or disease caused by increased biliary tract motility.